


# QUANTUM GRAVITY- THE EL DORADO – NAY A NE PLUS ULTRA --THE FINAL FINALE

<sup>1</sup>DR K N PRASANNA KUMAR, <sup>2</sup>PROF B S KIRANAGI AND <sup>3</sup>PROF C S BAGEWADI

**ABSTRACT:** *Motivation for quantizing gravity comes from the remarkable success of the quantum theories of the other three fundamental interactions, and from experimental evidence suggesting that gravity can be made to show quantum effects. Although some quantum gravity theories such as string theory and other unified field theories (or 'theories of everything') attempt to unify gravity with the other fundamental forces, others such as loop quantum gravity make no such attempt; they simply quantize the gravitational field while keeping it separate from the other forces. Observed physical phenomena can be described well by quantum mechanics or general relativity, without needing both. This can be thought of as due to an extreme separation of mass scales at which they are important. Quantum effects are usually important only for the "very small", that is, for objects no larger than typical molecules. General relativistic effects, on the other hand, show up mainly for the "very large" bodies such as collapsed stars. (Planets' gravitational fields, as of 2011, are well-described by linearised except for Mercury's perihelion precession; so strong-field effects—any effects of gravity beyond lowest nonvanishing order in  $\varphi/c^2$ —have not been observed even in the gravitational fields of planets and main sequence stars). There is a lack of experimental evidence relating to quantum gravity, and classical physics adequately describes the observed effects of gravity over a range of 50 orders of magnitude of mass, i.e., for masses of objects from about 10–23 to 1030 kg. We present a complete Model which probably explains the positivities and discrepancies and inadequacies of each model. Physics is certainly moving in to the subterranean realm and ceratoid dualism of consciousness and subject object duality (Freud vouchsafed only at the mother's breast shall the subject and object shall be one), like a maverick trying to transcend the boundaries of space time, standing on the threshold of infinity trying to ponder what lies beyond the veil which separates the scene from unseen?*



**INTRODUCTION:**

The following figurative representation is explains in best possible words the model that is proposed. A consummate model encompassing all the theories is presented. The theories are there to be applied to various physical systems which have different parametric representationalitiesof. Concept of “Theory”is explained in previous examples. And the bank’s example of conservativeness of individual debits and credits and the holistic conservativeness of assets and Liability is pronouncedly predominant in this case also. We shall not repeat in the following the same argument. One more factor that is to be remarked is that there are possibilities of concatenation of same theory with different theories. That the name appeared twice in the Model should not foreclose its option for its relationship with others.

**CLASSICAL MECHANICS AND NEWTONIAN GRAVITY:**

**MODULE NUMBERED ONE**

**NOTATION :**

$G_{13}$  : CATEGORY ONE OF CLASSICAL MECHANICS

$G_{14}$  : CATEGORY TWO OF CLASICAL MECHANICS

$G_{15}$  : CATEGORY THREE OF CLASSICAL MECHANICS

$T_{13}$  : CATEGORY ONE OF NEWTONIAN GRAVITY

$T_{14}$  : CATEGORY TWO OF NEWTONIAN GRAVITY

$T_{15}$  : CATEGORY THREE OF NEWTONIAN GRAVITY (WE ARE TALKING OF SYSTEMS; LAW IS THERE BUT IS APPLICABLE TO VARIOUS SYSTEMS) INVARIANT SU(3), THE PHYSICAL PARAMETER STATES

---

---

**QUANTUM MECHANICS AND QUANTUM FIELD THEORY:**

**MODULE NUMBERED TWO:**

=====

===

$G_{16}$  : CATEGORY ONE OF QUANTUM MECHANICS

$G_{17}$  : CATEGORY TWO OF QUANTUM MECHANICS

$G_{18}$  : CATEGORY THREE OF QUANTUM MECHANICS

$T_{16}$  : CATEGORY ONE OF QUANTUM FIELD THEORY

$T_{17}$  : CATEGORY TWO OF QUANTUM FIELD THEORY

$T_{18}$  : CATEGORY THREE OF QUANTUM FIELD THEORY

---

---

**ELECTROMAGNETISM AND STR (SPECIAL THEORY OF RELATIVITY):**

**MODULE NUMBERED THREE:**

=====

===

$G_{20}$  : CATEGORY ONE OF ELECTROMAGNETISM

$G_{21}$  : CATEGORY TWO OF ELECTROMAGNETIC THEORY

$G_{22}$  : CATEGORY THREE OF ELECTROMAGNETIC THEORY

$T_{20}$  : CATEGORY ONE OF STR

$T_{21}$  : CATEGORY TWO OF STR

$T_{22}$  : CATEGORY THREE OF STR

---

---

**GTR (GENERAL THEORY OF RELATIVITY) AND QFT (QUANTUM FIELD THEORY) IN CURVED SPACE TIME (BASED ON CERTAIN VARIABLES OF THE SYSTEM WHICH CONSEQUENTIALLY CLASSIFIABLE ON PARAMETERS)**

**: MODULE NUMBERED FOUR:**

=====

===

$G_{24}$  : CATEGORY ONE OFGTREVALUATIVE PARAMETRICIZATION OF SITUATIONAL ORIENTATIONS AND ESSENTIAL COGNITIVE ORIENTATION AND CHOICE VARIABLES OF THE SYSTEM TO WHICH QFT IS APPLICABLE)

$G_{25}$  : CATEGORY TWO OF GTR

$G_{26}$  : CATEGORY THREE OF GTR

$T_{24}$  :CATEGORY ONE OF QFT IN CURVED SPACE TIME

$T_{25}$  :CATEGORY TWO OF QFT(SYSTEMIC INSTRUMENTAL CHARACTERISATIONS AND ACTION ORIENTATIONS AND FUYNCTIONAL IMPERATIVES OF CHANGE MANIFESTED THEREIN )

$T_{26}$  : CATEGORY THREE OF QUANTUM FIELD THEORY

---

---

**GTR(GENERAL THEORY OF RELATIVITY(THERE ARE MANY OBSERVES AND GTR IS APPLICABLE TO BILLION SYSTEMS NOTWITHSTANDING THE GENERALISATIONAL NATURE OF THE THEORY) AND QUANTUM GRAVITY**

**MODULE NUMBERED FIVE:**

=====

===

$G_{28}$  : CATEGORY ONE OF GTR

$G_{29}$  : CATEGORY TWO OF QUANTUM GRAVITY

$G_{30}$  :CATEGORY THREE OFQUANTUM GRAVITY(THE FINAL THEORY MUST POSSESS THE SAME CHARACTERSTICS OF ITS CONSTITUENTS-IT CANNOT SIT IN IVORY TOWER WITHOUT APPLICABILITY TO VARIOUS SYSTEMS)

$T_{28}$  :CATEGORY ONE OF QUANTUM GRAVITY

$T_{29}$  :CATEGORY TWO OFQUANTUM GRAVITY

$T_{30}$  :CATEGORY THREE OF QUANTUM GRAVITY

=====

**QFT IN CURVED SPACE TIME AND QUANTUM GRAVITY:**

**MODULE NUMBERED SIX:**

---

---

=====

===

$G_{32}$  : CATEGORY ONE OF QFT IN CURVED SPACE AND TIME

$G_{33}$  : CATEGORY TWO OF QFT IN SPACE AND TIME

$G_{34}$  : CATEGORY THREE OF QFT IN CURVED SPACE AND TIME

$T_{32}$  : CATEGORY ONE OF QUANTUM GRAVITY

$T_{33}$  : CATEGORY TWO OF QUANTUM GRAVITY

$T_{34}$  : CATEGORY THREE OF QUANTUM GRAVITY

**GTR AND QFT IN CURVED SPACE TIME**

**MODULE NUMBERED SEVEN**

$G_{36}$  : CATEGORY ONE OF GTR

$G_{37}$  : CATEGORY TWO OF GTR

$G_{38}$  : CATEGORY THREE OF GTR

$T_{36}$  : CATEGORY ONE OF QFT IN CURVED SPACE TIME

$T_{37}$  : CATEGORY TWO OF QFT IN CURVED SPACE TIME

$T_{38}$  : CATEGORY THREE OF QFT IN CURVED SPACE AND TIME

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$   
 $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}, (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$   
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$   
 $(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$   
 $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$   
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$   
 $(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

are Dissipation coefficients

**CLASSICAL MECHANICS AND NEWTONIAN GRAVITY:**

1

**MODULE NUMBERED ONE**

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13}$$

2

$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14}$	3
$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15}$	4
$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13}$	5
$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14}$	6
$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15}$	7
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor	8
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor	

**QUANTUM MECHANICS AND QUANTUM FIELD THEORY:**

9

**MODULE NUMBERED TWO:**

The differential system of this model is now ( Module numbered two)

$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16}$	10
$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17}$	11
$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18}$	12
$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16}$	13
$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17}$	14
$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18}$	15
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor	16
$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor	17

**ELECTROMAGNETISM AND STR(SPECIAL THEORY OF RELATIVITY):**

18

**MODULE NUMBERED THREE:**

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20}$	19
$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21}$	20
$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22}$	21
$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20}$	22

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 23$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 24$$

$+(a''_{20})^{(3)}(T_{21}, t) =$  First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$  First detritions factor 25

**GTR(GENERAL THEORY OF RELATIVITY )ANDQFT(QUANTUM FIELD THEORY)IN CURVED SPACE TIME(BASED ON CERTAIN VARAIBLES OF THE SYSTEM WHICH CONSEQUENTIALLY CLSSIFIABLE ON PARAMETERS)** 26

**: MODULE NUMBERED FOUR)**

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 27$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 28$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 29$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 30$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 31$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 32$$

$+(a''_{24})^{(4)}(T_{25}, t) =$  First augmentation factor 33

$-(b''_{24})^{(4)}((G_{27}), t) =$  First detritions factor 34

**GTR(GENERAL THEORY OF RELATIVITY(THERE ARE MANY OBSERVES AND GTR IS APPLICABLE TO BILLION SYSTEMS NOTWITHSTANDING THE GENERALISATIONAL NATURE OF THE THEORY) AND QUANTUM GRAVITY** 35

**MODULE NUMBERED FIVE**

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 36$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 37$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 38$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 39$$

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29}$	40
$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30}$	41
$+(a''_{28})^{(5)}(T_{29}, t) =$ <b>First augmentation factor</b>	42
$-(b''_{28})^{(5)}((G_{31}), t) =$ <b>First detritions factor</b>	43
	44
<b><u>QFT IN CURVED SPACE TIME AND QUANTUM GRAVITY:</u></b>	45

**MODULE NUMBERED SIX:**

The differential system of this model is now (Module numbered Six)

$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32}$	46
$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33}$	47
$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34}$	48
$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32}$	49
$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33}$	50
$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34}$	51
$+(a''_{32})^{(6)}(T_{33}, t) =$ <b>First augmentation factor</b>	52

**GTR AND QFT IN CURVED SPACE TIME**

**MODULE NUMBERED SEVEN:**

The differential system of this model is now (SEVENTH MODULE)

$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36}$	54
$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37}$	55
$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38}$	56
$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36}$	57
$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37}$	58



59

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38}$$

60

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$$

61

$$-(b''_{36})^{(7)}((G_{39}), t) = \text{First detritions factor}$$

62

### FIRST MODULE CONCATENATION:

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[ \begin{array}{c} (a'_{13})^{(1)} \boxed{+(a''_{13})^{(1)}(T_{14}, t)} \boxed{+(a'_{16})^{(2,2)}(T_{17}, t)} \boxed{+(a'_{20})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7)}(T_{37}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[ \begin{array}{c} (a'_{14})^{(1)} \boxed{+(a''_{14})^{(1)}(T_{14}, t)} \boxed{+(a'_{17})^{(2,2)}(T_{17}, t)} \boxed{+(a'_{21})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7)}(T_{37}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[ \begin{array}{c} (a'_{15})^{(1)} \boxed{+(a''_{15})^{(1)}(T_{14}, t)} \boxed{+(a'_{18})^{(2,2)}(T_{17}, t)} \boxed{+(a'_{22})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7)}(T_{37}, t)} \end{array} \right] G_{15}$$

Where  $\boxed{(a''_{13})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{14})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{15})^{(1)}(T_{14}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7)}(T_{37}, t)}$   $\boxed{+(a''_{37})^{(7)}(T_{37}, t)}$   $\boxed{+(a''_{38})^{(7)}(T_{37}, t)}$  ARE SEVENTH AUGMENTATION COEFFICIENTS

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{c} (b'_{13})^{(1)} \boxed{-(b''_{16})^{(1)}(G, t)} \boxed{-(b''_{36})^{(7)}(G_{39}, t)} \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7)}(G_{39}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{ccc} (b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} & \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} & & \end{array} \right] T_{14}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{ccc} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} & \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} & & \end{array} \right] T_{15}$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$  are fifth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$  are sixth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$  ARE SEVENTH DETRITION COEFFICIENTS

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{ccc} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} & \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{15} \tag{63}$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detrition coefficients for category 1, 2 and 3 64  
 $\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$  are fifth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$  are sixth detritions coefficients for category 1, 2 and 3

**SECOND MODULE CONCATENATION:** 65

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[ \begin{array}{ccc} (a'_{16})^{(2)} \boxed{+(a''_{16})^{(2)}(T_{17}, t)} & \boxed{+(a''_{13})^{(1,1)}(T_{14}, t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)} & & \end{array} \right] G_{16} \tag{66}$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[ \begin{array}{c} (a'_{17})^{(2)} \boxed{+(a''_{17})^{(2)}(T_{17}, t)} \boxed{+(a''_{14})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{21})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37}, t)} \end{array} \right] G_{17} \quad 67$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[ \begin{array}{c} (a'_{18})^{(2)} \boxed{+(a''_{18})^{(2)}(T_{17}, t)} \boxed{+(a''_{15})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{22})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37}, t)} \end{array} \right] G_{18} \quad 68$$

Where  $\boxed{+(a''_{16})^{(2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2)}(T_{17}, t)}$  are first augmentation coefficients for category 1, 2 and 3 69

$\boxed{+(a''_{13})^{(1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1)}(T_{14}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3 70

$\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$  ARE SEVENTH DETRITION COEFFICIENTS 71

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[ \begin{array}{c} (b'_{16})^{(2)} \boxed{-(b''_{16})^{(2)}(G_{19}, t)} \boxed{-(b''_{13})^{(1,1)}(G, t)} \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \end{array} \right] T_{16} \quad 72$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[ \begin{array}{c} (b'_{17})^{(2)} \boxed{-(b''_{17})^{(2)}(G_{19}, t)} \boxed{-(b''_{14})^{(1,1)}(G, t)} \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \end{array} \right] T_{17} \quad 73$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[ \begin{array}{c} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} \boxed{-(b''_{15})^{(1,1)}(G, t)} \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \end{array} \right] T_{18} \quad 74$$

where  $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$  are first detrition coefficients for category 1, 2 and 3 75

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1)}(G, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detritions coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$  are sixth detritions coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7,7)}(G_{39}, t)$  are seventh detrition coefficients

### THIRD MODULE CONCATENATION:

$$\frac{dG_{20}}{dt} = \tag{76}$$

$$(a_{20})^{(3)} G_{21} - \left[ \begin{array}{c} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[ \begin{array}{c} (a'_{21})^{(3)} \boxed{+(a''_{21})^{(3)}(T_{21}, t)} \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{21} \tag{77}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[ \begin{array}{c} (a'_{22})^{(3)} \boxed{+(a''_{22})^{(3)}(T_{21}, t)} \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{22} \tag{78}$$

79

$\boxed{+(a''_{20})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3)}(T_{21}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2)}(T_{17}, t)}$  are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$  are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficients for category 1, 2 and 3

80

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficient

81

$$\frac{dT_{20}}{dt} = \tag{82}$$

$$(b_{20})^{(3)} T_{21} - \left[ \begin{array}{c} (b'_{20})^{(3)} \boxed{-(b''_{20})^{(3)}(G_{23}, t)} \boxed{-(b'_{36})^{(7,7,7)}(G_{19}, t)} \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{20}$$

$$\frac{dT_{21}}{dt} = \tag{83}$$

$$(b_{21})^{(3)}T_{20} - \left[ \begin{array}{ccc} (b'_{21})^{(3)} \boxed{-(b''_{21})^{(3)}(G_{23}, t)} & \boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ & \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \end{array} \right] T_{21}$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{ccc} (b'_{22})^{(3)} \boxed{-(b''_{22})^{(3)}(G_{23}, t)} & \boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ & \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \end{array} \right] T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b'_{21})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$  are first detritions coefficients for category 1, 2 and 3 85

$\boxed{-(b'_{16})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b'_{17})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b'_{18})^{(2,2,2)}(G_{19}, t)}$  are second detritions coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b'_{14})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b'_{15})^{(1,1,1)}(G, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$  are seventh detritions coefficients 86

=====

### FOURTH MODULE CONCATENATION:

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{ccc} (a'_{24})^{(4)} \boxed{+(a''_{24})^{(4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)} \\ & \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} & \end{array} \right] G_{24}$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{ccc} (a'_{25})^{(4)} \boxed{+(a''_{25})^{(4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)} \\ & \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} & \end{array} \right] G_{25}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{ccc} (a'_{26})^{(4)} \boxed{+(a''_{26})^{(4)}(T_{25}, t)} & \boxed{+(a''_{30})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{34})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)} \\ & \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)} & \end{array} \right] G_{26}$$

90

Where  $\boxed{(a''_{24})^{(4)}(T_{25}, t)}$ ,  $\boxed{(a''_{25})^{(4)}(T_{25}, t)}$ ,  $\boxed{(a''_{26})^{(4)}(T_{25}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5)}(T_{29}, t)}$  are second augmentation coefficient for category 1, 2 and 3 <sup>91</sup>

$\boxed{+(a''_{32})^{(6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$  ARE SEVENTH augmentation coefficients

92

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[ \begin{array}{ccc} \boxed{(b'_{24})^{(4)} \boxed{-(b''_{24})^{(4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}} & & \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)} & & \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} & & \end{array} \right] T_{24} \quad 93$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[ \begin{array}{ccc} \boxed{(b'_{25})^{(4)} \boxed{-(b''_{25})^{(4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}} & & \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)} & & \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} & & \end{array} \right] T_{25} \quad 94$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[ \begin{array}{ccc} \boxed{(b'_{26})^{(4)} \boxed{-(b''_{26})^{(4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}} & & \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)} & & \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} & & \end{array} \right] T_{26} \quad 95$$

Where  $\boxed{-(b''_{24})^{(4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4)}(G_{27}, t)}$  are first detrition coefficients for category 1, 2 and 3 <sup>96</sup>

$\boxed{-(b''_{28})^{(5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G, t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G, t)}$   
 are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)}$   
 are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)}$   
 are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$  ARE SEVENTH DETRITION

COEFFICIENTS

97

**FIFTH MODULE CONCATENATION:**

98

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[ \begin{array}{c} (a'_{28})^{(5)} \boxed{+(a''_{28})^{(5)}(T_{29}, t)} \quad \boxed{+(a''_{24})^{(4,4)}(T_{25}, t)} \quad \boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{28} \quad 99$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[ \begin{array}{c} (a'_{29})^{(5)} \boxed{+(a''_{29})^{(5)}(T_{29}, t)} \quad \boxed{+(a''_{25})^{(4,4)}(T_{25}, t)} \quad \boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{29} \quad 100$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[ \begin{array}{c} (a'_{30})^{(5)} \boxed{+(a''_{30})^{(5)}(T_{29}, t)} \quad \boxed{+(a''_{26})^{(4,4)}(T_{25}, t)} \quad \boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{30} \quad 101$$

Where  $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$  are first augmentation coefficients for category 1, 2 and : 102

And  $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$  are second augmentation coefficient for category 1, 2 a

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and :

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[ \begin{array}{c} (b'_{28})^{(5)} \boxed{-(b''_{28})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{24})^{(4,4)}(G_{23}, t)} \quad \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{28} \quad 103$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[ \begin{array}{c} (b'_{29})^{(5)} \boxed{-(b''_{29})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} \quad \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{29} \quad 104$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)} T_{29} - \left[ \begin{array}{c} (b'_{30})^{(5)} \boxed{-(b''_{30})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} \quad \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{30} \quad 105$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1, 2 and 3 106

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1,2, and 3

## SIXTH MODULE CONCATENATION

108

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[ \begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ & + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & \end{array} \right] G_{32}$$

109

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ & + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & \end{array} \right] G_{33}$$

110

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ & + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & \end{array} \right] G_{34}$$

111

$+(a''_{32})^{(6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6)}(T_{33}, t)$  are first augmentation coefficients for category 1, 2 and 3

112

$+(a''_{28})^{(5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5)}(T_{29}, t)$  are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4)}(T_{25}, t)$  are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$  - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$  - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$  sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$ ,  $+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$ ,  $+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)$  ARE SEVENTH AUGMENTATION COEFFICIENTS

113

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ & - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & \end{array} \right] T_{32}$$

114



$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{l} (b'_{33})^{(6)} \boxed{-(b''_{33})^{(6)}(G_{35}, t)} \quad \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)} \end{array} \right] T_{33} \quad 115$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{l} (b'_{34})^{(6)} \boxed{-(b''_{34})^{(6)}(G_{35}, t)} \quad \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} \end{array} \right] T_{34} \quad 116$$

$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6)}(G_{35}, t)}$  are first detrition coefficients for category 1, 2 and 3 117

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)}$  ARE SEVENTH DETRITION COEFFICIENTS

118

## SEVENTH MODULE CONCATENATION:

119

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[ \begin{array}{l} \boxed{(a'_{36})^{(7)}} + \boxed{(a''_{36})^{(7)}(T_{37}, t)} + \boxed{(a''_{16})^{(7)}(T_{17}, t)} + \boxed{(a''_{20})^{(7)}(T_{21}, t)} + \boxed{(a''_{24})^{(7)}(T_{23}, t)G_{36}} + \\ \boxed{(a''_{28})^{(7)}(T_{29}, t)} + \boxed{(a''_{32})^{(7)}(T_{33}, t)} + \boxed{(a''_{13})^{(7)}(T_{14}, t)} \end{array} \right] G_{36} \quad 120$$

121

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[ \begin{array}{l} \boxed{(a'_{37})^{(7)}} + \boxed{(a''_{37})^{(7)}(T_{37}, t)} + \boxed{(a''_{14})^{(7)}(T_{14}, t)} + \boxed{(a''_{21})^{(7)}(T_{21}, t)} + \\ \boxed{(a''_{17})^{(7)}(T_{17}, t)} + \boxed{(a''_{25})^{(7)}(T_{25}, t)} + \boxed{(a''_{33})^{(7)}(T_{33}, t)} + \boxed{(a''_{29})^{(7)}(T_{29}, t)} \end{array} \right] G_{37} \quad 122$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} -$$

123

$$\left[ \begin{array}{l} \boxed{(a'_{38})^{(7)}} + \boxed{(a''_{38})^{(7)}(T_{37}, t)} + \boxed{(a''_{15})^{(7)}(T_{14}, t)} + \boxed{(a''_{22})^{(7)}(T_{21}, t)} + \boxed{(a''_{18})^{(7)}(T_{17}, t)} \end{array} \right] +$$

125

$$\left[ (a''_{26})^{(7)}(T_{25}, t) + (a''_{34})^{(7)}(T_{33}, t) + (a''_{30})^{(7)}(T_{29}, t) \right] G_{38}$$

$$\frac{dT_{36}}{dt} = \left[ (b'_{36})^{(7)}T_{37} - \left[ (b'_{36})^{(7)} - \left[ (b''_{36})^{(7)}((G_{39}), t) - (b''_{16})^{(7)}((G_{19}), t) - (b''_{13})^{(7)}((G_{14}), t) - (b''_{20})^{(7)}((G_{231}), t) - (b''_{24})^{(7)}((G_{27}), t) - (b''_{28})^{(7)}((G_{31}), t) - (b''_{32})^{(7)}((G_{35}), t) \right] T_{36} \right] T_{36} \quad 126$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[ (b'_{36})^{(7)} - \left[ (b''_{37})^{(7)}((G_{39}), t) - (b''_{17})^{(7)}((G_{19}), t) - (b''_{19})^{(7)}((G_{14}), t) - (b''_{21})^{(7)}((G_{231}), t) - (b''_{25})^{(7)}((G_{27}), t) - (b''_{29})^{(7)}((G_{31}), t) - (b''_{33})^{(7)}((G_{35}), t) \right] T_{37} \right] T_{37} \quad 127$$

Where we suppose

- (A)  $(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0,$   
 $i, j = 13, 14, 15$
- (B) The functions  $(a''_i)^{(1)}, (b''_i)^{(1)}$  are positive continuous increasing and bounded.  
**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

- (C)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)}$   
 $\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants  
 and  $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} |G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(1)}(T'_{14}, t)$  and  $(a''_i)^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a''_i)^{(1)}(T_{14}, t)$

is uniformly continuous. In the eventuality of the fact, that if  $(\widehat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

**Definition of**  $(\widehat{M}_{13})^{(1)}, (\widehat{k}_{13})^{(1)}$  :

(D)  $(\widehat{M}_{13})^{(1)}, (\widehat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\widehat{P}_{13})^{(1)}, (\widehat{Q}_{13})^{(1)}$  :

(E) There exists two constants  $(\widehat{P}_{13})^{(1)}$  and  $(\widehat{Q}_{13})^{(1)}$  which together with  $(\widehat{M}_{13})^{(1)}, (\widehat{k}_{13})^{(1)}, (\widehat{A}_{13})^{(1)}$  and  $(\widehat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\widehat{M}_{13})^{(1)}} [ (a_i)^{(1)} + (a_i')^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)} ] < 1$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}} [ (b_i)^{(1)} + (b_i')^{(1)} + (\widehat{B}_{13})^{(1)} + (\widehat{Q}_{13})^{(1)} (\widehat{k}_{13})^{(1)} ] < 1$$



$$\frac{dT_{38}}{dt} = \left[ (b_{38})^{(7)} T_{37} - \left[ (b'_{38})^{(7)} - \left[ (b''_{38})^{(7)}((G_{39}), t) - \left[ (b''_{18})^{(7)}((G_{19}), t) - \left[ (b''_{20})^{(7)}((G_{14}), t) - \left[ (b''_{22})^{(7)}((G_{23}), t) - \left[ (b''_{26})^{(7)}((G_{27}), t) - \left[ (b''_{30})^{(7)}((G_{31}), t) - \left[ (b''_{34})^{(7)}((G_{35}), t) \right] T_{38} \right] \right] \right] \right] \right] T_{38} \quad 128$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \quad 134$$

$$(1)(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18 \quad 135$$

(F) (2) The functions  $(a''_i)^{(2)}, (b''_i)^{(2)}$  are positive continuous increasing and bounded. 136

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ : 137

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 138$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 139$$

$$(G) (3) \lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \quad 141$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$ : 142

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$

They satisfy Lipschitz condition: 143

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T'_{17} - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 144$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 145$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(2)}(T'_{17}, t)$  and  $(a''_i)^{(2)}(T_{17}, t)$ .  $(T'_{17}, t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a''_i)^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a''_i)^{(2)}(T_{17}, t)$ , the SECOND augmentation coefficient would be absolutely continuous.

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ : 147

(H) (4)  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants 148

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$ : 149

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$ ,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 150$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 151$$

Where we suppose 152

$$(I) \quad (5) \quad (a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 153$$

The functions  $(a''_i)^{(3)}, (b''_i)^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 154$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)} \quad 155$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$  : 156

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$

They satisfy Lipschitz condition: 157

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t} \quad 158$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23} - G'_{23}| e^{-(\hat{M}_{20})^{(3)}t} \quad 159$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(3)}(T'_{21}, t)$  and  $(a''_i)^{(3)}(T_{21}, t)$ .  $(T'_{21}, t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a''_i)^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a''_i)^{(3)}(T_{21}, t)$ , the THIRD augmentation coefficient, would be absolutely continuous. 160

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$  : 161

(J) (6)  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants 162

$(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22,$  163

satisfy the inequalities 164

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \quad 165$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \quad 166$$

167

Where we suppose 168

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 169$$

(L) (7) The functions  $(a''_i)^{(4)}, (b''_i)^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

170

(M) (8)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)}$   
 $\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$ :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

They satisfy Lipschitz condition:

171

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} \|(G_{27})' - (G_{27})\| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(4)}(T'_{25}, t)$  and  $(a''_i)^{(4)}(T_{25}, t) \cdot (T'_{25}, t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a''_i)^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 4$  then the function  $(a''_i)^{(4)}(T_{25}, t)$ , the **FOURTH augmentation coefficient WOULD** be absolutely continuous.

173

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ :

174

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$ :

175

(P) (9) There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose 176

$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30$  177  
 (R) (10) The functions  $(a''_i)^{(5)}, (b''_i)^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$ :

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}(G_{31}, t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

178

(S) (11)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)}$   
 $\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$  :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$

They satisfy Lipschitz condition: 179

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b''_i)^{(5)}(G'_{31}, t) - (b''_i)^{(5)}(G_{31}, t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G'_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(5)}(T'_{29}, t)$  180  
 and  $(a''_i)^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is  
 to be noted that  $(a''_i)^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  
 $(\hat{M}_{28})^{(5)} = 5$  then the function  $(a''_i)^{(5)}(T_{29}, t)$ , the FIFTH **augmentation coefficient** attributable  
 would be absolutely continuous.

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$  : 181

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$  are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$  : 182

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  
 $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  
 $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose 183

$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$  184  
 (12) The functions  $(a''_i)^{(6)}, (b''_i)^{(6)}$  are positive continuous increasing and bounded.



**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

185

(13)  $\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$   
 $\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$  :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

186

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} |(G_{35}) - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(6)}(T_{33}', t)$  and  $(a_i'')^{(6)}(T_{33}, t)$ .  $(T_{33}', t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a_i'')^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 6$  then the function  $(a_i'')^{(6)}(T_{33}, t)$ , the SIXTH **augmentation coefficient** would be absolutely continuous.

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$  :

188

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants  
 $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$  :

189

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

190

(V)  $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0,$   
 $i, j = 36, 37, 38$  191

(W) The functions  $(a_i'')^{(7)}, (b_i'')^{(7)}$  are positive continuous increasing and bounded.  
**Definition of**  $(p_i)^{(7)}, (r_i)^{(7)}$ :

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

192

(X)  $\lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$   
 $\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

**Definition of**  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$  :

Where  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$  are positive constants  
 and  $i = 36, 37, 38$

They satisfy Lipschitz condition:

193

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), (T_{39}))| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(7)}(T_{37}', t)$  and  $(a_i'')^{(7)}(T_{37}, t)$ .  $(T_{37}', t)$  and  $(T_{37}, t)$  are points belonging to the interval  $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$ . It is to be noted that  $(a_i'')^{(7)}(T_{37}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{36})^{(7)} = 7$  then the function  $(a_i'')^{(7)}(T_{37}, t)$ , the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous. 194

**Definition of**  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$  :

195

(Y)  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$ , are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

**Definition of**  $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$  :

196

(Z) There exists two constants  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  which together with  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$  and  $(\hat{B}_{36})^{(7)}$  and the constants  $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [ (a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)} ] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [ (b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)} ] < 1$$

197

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \boxed{T_i(0) = T_i^0 > 0}$$

198

**Definition of**  $G_i(0), T_i(0)$  :

199

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \boxed{T_i(0) = T_i^0 > 0}$$

=====

=

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions 200  
 $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 201$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 202$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 203$$

By 204

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t [(a_{13})^{(1)} G_{14}(s_{(13)}) - ((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})] G_{13}(s_{(13)})] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t [(a_{14})^{(1)} G_{13}(s_{(13)}) - ((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}))] G_{14}(s_{(13)})] ds_{(13)} \quad 205$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t [(a_{15})^{(1)} G_{14}(s_{(13)}) - ((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}))] G_{15}(s_{(13)})] ds_{(13)} \quad 206$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t [(b_{13})^{(1)} T_{14}(s_{(13)}) - ((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}))] T_{13}(s_{(13)})] ds_{(13)} \quad 207$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t [(b_{14})^{(1)} T_{13}(s_{(13)}) - ((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}))] T_{14}(s_{(13)})] ds_{(13)} \quad 208$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t [(b_{15})^{(1)} T_{14}(s_{(13)}) - ((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}))] T_{15}(s_{(13)})] ds_{(13)} \quad 209$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

210

if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

By

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36}(s_{(36)})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)} (G(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)} (G(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)} (G(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$



211

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)}, \quad 212$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad 213$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad 214$$

By 215

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)} \quad 216$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)} \quad 217$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)} \quad 218$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)} \quad 219$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)} \quad 220$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

221

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)}, \quad 222$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad 223$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad 224$$

By 225

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)} \quad 226$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)} \quad 227$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)}(G(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)} \quad 228$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)}(G(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)} \quad 229$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)}(G(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)} \quad 230$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 231

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)}, \quad 232$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad 233$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad 234$$

By 235

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)} \quad 236$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)} \quad 237$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)} \quad 238$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)}(G(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)} \quad 239$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)}(G(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)} \quad 240$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)}(G(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)} \quad 240$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 241

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)}, \quad 242$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad 243$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad 244$$

By 245

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)} \quad 246$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)} \quad 247$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)} \quad 248$$



$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)} \quad 249$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)} \quad 250$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)} \quad 251$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

252

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)}, \quad 253$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad 254$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad 255$$

By 256

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32}^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)} \quad 257$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)} \quad 258$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)} \quad 259$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)} \quad 260$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)} \quad 261$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

: if the conditions IN THE FOREGOING are fulfilled, there exists a solution satisfying the conditions 262

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:**

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$

which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)}, \quad 263$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad 264$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad 265$$

By 266

$$\begin{aligned} \bar{G}_{36}(t) &= G_{36}^0 + \\ &\int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36}(T_{37}(s_{(36)}), s_{(36)})) G_{36}(s_{(36)}) \right) \right] ds_{(36)} \end{aligned} \quad 267$$

$$\begin{aligned} \bar{G}_{37}(t) &= G_{37}^0 + \\ &\int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})) G_{37}(s_{(36)}) \right) \right] ds_{(36)} \end{aligned}$$

$$\begin{aligned} \bar{G}_{38}(t) &= G_{38}^0 + \\ &\int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})) G_{38}(s_{(36)}) \right) \right] ds_{(36)} \end{aligned} \quad 268$$

$$\begin{aligned} \bar{T}_{36}(t) &= T_{36}^0 + \\ &\int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)}(G(s_{(36)}), s_{(36)})) T_{36}(s_{(36)}) \right) \right] ds_{(36)} \end{aligned} \quad 269$$

$$\begin{aligned} \bar{T}_{37}(t) &= T_{37}^0 + \\ &\int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)}(G(s_{(36)}), s_{(36)})) T_{37}(s_{(36)}) \right) \right] ds_{(36)} \end{aligned} \quad 270$$

$$\begin{aligned} \bar{T}_{38}(t) &= T_{38}^0 + \\ &\int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)}(G(s_{(36)}), s_{(36)})) T_{38}(s_{(36)}) \right) \right] ds_{(36)} \end{aligned} \quad 271$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$  272

(a) The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself 273  
 .Indeed it is obvious that

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} \left( G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$\left( 1 + (a_{24})^{(4)} t \right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\bar{M}_{24})^{(4)}} \left( e^{(\bar{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 274

$$(G_{24}(t) - G_{24}^0) e^{-(\bar{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\bar{M}_{24})^{(4)}} \left[ \left( (\hat{P}_{24})^{(4)} + G_{25}^0 \right) e^{-\left( -\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0} \right)} + (\hat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

(b) The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself 275  
 .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} \left( G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$\left( 1 + (a_{28})^{(5)} t \right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\bar{M}_{28})^{(5)}} \left( e^{(\bar{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that 276

$$(G_{28}(t) - G_{28}^0) e^{-(\bar{M}_{28})^{(5)} t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}} \left[ \left( (\hat{P}_{28})^{(5)} + G_{29}^0 \right) e^{-\left( -\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0} \right)} + (\hat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

(c) The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself 277  
 .Indeed it is obvious that

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} \left( G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left( 1 + (a_{32})^{(6)} t \right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}} \left( e^{(\bar{M}_{32})^{(6)} t} - 1 \right)$$

From which it follows that 278

$$(G_{32}(t) - G_{32}^0) e^{-(\bar{M}_{32})^{(6)} t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ \left( (\hat{P}_{32})^{(6)} + G_{33}^0 \right) e^{-\left( -\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0} \right)} + (\hat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(d) The operator  $\mathcal{A}^{(7)}$  maps the space of functions satisfying 37,35,36 into itself .Indeed it is obvious that 279

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} \left( G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)} t) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left( e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

280

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[ ((\hat{P}_{36})^{(7)} + G_{37}^0) e^{-\left( \frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 7

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$  and to choose 281

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have 282

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\left( \frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{13})^{(1)} \quad 283$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ ((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 284$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying GLOBAL EQUATIONS into itself 285

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric 286

$$d \left( (G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \right\}$$

Indeed if we denote 287

**Definition of  $\tilde{G}, \tilde{T}$  :**

$$(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$$

It results

$$\begin{aligned}
 |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\
 &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\
 &(a''_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\
 &G_{13}^{(2)} |(a'_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) - (a'_{13})^{(1)}(T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}
 \end{aligned}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned}
 |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq & 288 \\
 \frac{1}{(\bar{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}))
 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{13})^{(1)}$  and  $(b''_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$  and  $(\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$  respectively of  $\mathbb{R}_+$ . 289

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a'_i)^{(1)}$  and  $(b'_i)^{(1)}$ ,  $i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  290

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0 \quad 291$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \quad \text{for } t > 0$$

**Definition of**  $((\bar{M}_{13})^{(1)})_1$ , and  $((\bar{M}_{13})^{(1)})_3$  : 292

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$$G_{13} < ((\bar{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\bar{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\bar{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\bar{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\bar{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\bar{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}$ ,  $G_{15}$  and  $G_{13}$ ,  $G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below. 293

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$  then  $T_{14} \rightarrow \infty$ . 294

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to 295

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_1}$  By taking now  $\varepsilon_1$  sufficiently small one sees that  $T_{14}$  is unbounded. The same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

296

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$  and to choose

297

$(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ (\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

298

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ ((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

299

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying

300

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric

301

$$d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

302

**Definition**  $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

303

$$|\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)} |(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$

304

From the hypotheses it follows

$$\begin{aligned} & |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)}t} \leq & 305 \\ & \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + \\ & (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d\left( ((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows 306

**Remark 1:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ . 307

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a'_i)^{(2)}$  and  $(b'_i)^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any t where  $G_i(t) = 0$  and  $T_i(t) = 0$  308

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{16})^{(2)})_1$ ,  $((\widehat{M}_{16})^{(2)})_2$  and  $((\widehat{M}_{16})^{(2)})_3$  : 309

**Remark 3:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$ . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)}) \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)} \quad 310$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$ ,  $G_{18}$  and  $G_{16}$ ,  $G_{17}$  respectively.

**Remark 4:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below. 311

**Remark 5:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$  then  $T_{17} \rightarrow \infty$ . 312

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b'_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to 313

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$ ,  $t = \log \frac{2}{\varepsilon_2}$  By taking now  $\varepsilon_2$  sufficiently small one sees that  $T_{17}$  is 314  
 unbounded. The same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b''_{18})^{(2)}((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}$ ,  $\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$  and to choose 315  
 316

$(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ (\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 317$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ ((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 318$$

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself 319

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric 320

$$d\left( ((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 321

**Definition of**  $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 322

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_{20}^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} \{ (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) \} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned} \quad 323$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ &\frac{1}{(\bar{M}_{20})^{(3)}} \left( (a_{20})^{(3)} + (a'_{20})^{(3)} + (\hat{A}_{20})^{(3)} + \right. \\ &\left. (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)} \right) d\left( ((G_{23})^{(1)}, (T_{23})^{(1)}); (G_{23})^{(2)}, (T_{23})^{(2)} \right) \end{aligned} \quad 324$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on t can be considered as 325



not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$  and  $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$ ,  $i = 20,21,22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$ (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  326

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$  and  $((\widehat{M}_{20})^{(3)})_3$  : 327

**Remark 3:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$  . indeed if

$$G_{20} < ((\widehat{M}_{20})^{(3)})_1 \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$  ,  $G_{22}$  and  $G_{20}$  ,  $G_{21}$  respectively.

**Remark 4:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below. 328

**Remark 5:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$  then  $T_{21} \rightarrow \infty$ . 329

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  : 330

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21}')^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to 331

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_3}$  By taking now  $\varepsilon_3$  sufficiently small one sees that  $T_{21}$  is unbounded. The same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

332

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$  ,  $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$  and to choose 333

$(\widehat{P}_{24})^{(4)}$  and  $(\widehat{Q}_{24})^{(4)}$  large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ (\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\widehat{P}_{24})^{(4)} \quad 334$$

$$\frac{(b_j)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ ((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)} \quad 335$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying IN to itself 336

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric 337

$$d\left( ((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{27}), (\widetilde{T}_{27}) : (\widetilde{G}_{27}), (\widetilde{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\widetilde{G}_{24}^{(1)} - \widetilde{G}_{24}^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)}(T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)}(T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)}(T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)}t} &\leq \quad 338 \\ \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + \\ (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) d\left( ((G_{27})^{(1)}, (T_{27})^{(1)}); (G_{27})^{(2)}, (T_{27})^{(2)} \right) &\quad 339 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  and  $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ . 340

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$ ,  $i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  341

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$  and  $((\widehat{M}_{24})^{(4)})_3$  : 342

**Remark 3:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$ . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})_1$  it follows  $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a_{25}')^{(4)}G_{25}$  and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25}')^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a_{25}')^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26}')^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a_{26}')^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$ ,  $G_{26}$  and  $G_{24}$ ,  $G_{25}$  respectively.

**Remark 4:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below. 343

**Remark 5:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$  then  $T_{25} \rightarrow \infty$ . 344

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25}')^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25}')^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to 345

$$T_{25} \geq \left( \frac{(a_{25}')^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$T_{25} \geq \left( \frac{(a_{25}')^{(4)}(m)^{(4)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_4}$  By taking now  $\varepsilon_4$  sufficiently small one sees that  $T_{25}$  is unbounded. The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b_{26}')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions ANALOGOUS inequalities hold also for  $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

346

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} < 1$  and to choose 347

$(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[ (\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 348$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[ ((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 349$$

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself 350

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric 351

$$d\left( (G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)} \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{31}), (\widetilde{T}_{31}) : ((\widetilde{G}_{31}), (\widetilde{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_{28}^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s} e^{(\bar{M}_{28})^{(5)}s} ds_{(28)} + \\ &\int_0^t ((a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s} e^{-(\bar{M}_{28})^{(5)}s} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s} e^{(\bar{M}_{28})^{(5)}s} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\bar{M}_{28})^{(5)}s} e^{(\bar{M}_{28})^{(5)}s} ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} &\leq \quad 353 \\ \frac{1}{(\bar{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + \\ (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}) d\left( (G_{31})^{(1)}, (T_{31})^{(1)}, (G_{31})^{(2)}, (T_{31})^{(2)} \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (35,35,36) the result follows

**Remark 1:** The fact that we supposed  $(a_{28}'')^{(5)}$  and  $(b_{28}'')^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ . 354

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$ ,  $i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  355

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t ((a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})) ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{28})^{(5)})_1$ ,  $((\widehat{M}_{28})^{(5)})_2$  and  $((\widehat{M}_{28})^{(5)})_3$  : 356

**Remark 3:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$$G_{28} < ((\widehat{M}_{28})^{(5)}) \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$ ,  $G_{30}$  and  $G_{28}$ ,  $G_{29}$  respectively.

**Remark 4:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below. 357

**Remark 5:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$  then  $T_{29} \rightarrow \infty$ . 358

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29}')^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)} \quad \text{359}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$  which leads to 360

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is}$$

unbounded. The same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

361

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} < 1$  and to choose

362

$(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ (\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)}$$

363

$$\frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ ((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)}$$

364

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

365

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric

366

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

$$\text{Definition of } (\widetilde{G_{35}}, \widetilde{T_{35}}) : (\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_{32}^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{-(\bar{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

367

From the hypotheses it follows

$$(1) \quad (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \\ i, j = 13, 14, 15$$

(2) The functions  $(a''_i)^{(1)}, (b''_i)^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$(3) \lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \\ \lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants  
 and  $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, T)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T'_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  :

(AA)  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  :

(BB) There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together with  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Analogous inequalities hold also for  $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$

368

It is now sufficient to take  $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 7$  and to choose

$(\widehat{P}_{36})^{(7)}$  and  $(\widehat{Q}_{36})^{(7)}$  large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ (\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 369$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ ((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 370$$

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 371  
 37,35,36 into itself

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric 372

$$d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right), \left((G_{39})^{(2)}, (T_{39})^{(2)}\right)\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of  $(\widetilde{G}_{39}), (\widetilde{T}_{39})$  :**

$$(\widetilde{G}_{39}), (\widetilde{T}_{39}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$$

It results

$$|\widetilde{G}_{36}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} e^{(\widehat{M}_{36})^{(7)}s_{(36)}} ds_{(36)} +$$

$$\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$$(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$$G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\} ds_{(36)}$$



Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

373

$$\begin{aligned} & |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widehat{M}_{36})^{(7)}t} \leq \\ & \frac{1}{(\widehat{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + \\ & (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}) d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned}$$

374

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (37,35,36) the result follows

**Remark 1:** The fact that we supposed  $(a''_{36})^{(7)}$  and  $(b''_{36})^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  and  $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 375

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(7)}$  and  $(b''_i)^{(7)}$ ,  $i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

376

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 79 to 36 it results

$$G_i(t) \geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i^{(7)})t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{36})^{(7)})_1$ ,  $((\widehat{M}_{36})^{(7)})_2$  and  $((\widehat{M}_{36})^{(7)})_3$  :

377

**Remark 3:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$  . indeed if

$G_{36} < ((\widehat{M}_{36})^{(7)})_1$  it follows  $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37}$  and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$  ,  $G_{38}$  and  $G_{36}$  ,  $G_{37}$  respectively.

**Remark 7:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below. 378

**Remark 5:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i^{(7)})^{(7)})((G_{39})(t), t) = (b'_{37})^{(7)}$  then  $T_{37} \rightarrow \infty$ . 379

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b_i^{(7)})^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 380

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$

If we take  $t$  such that  $e^{-\varepsilon_7 t} = \frac{1}{2}$  it results

$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_7}$  By taking now  $\varepsilon_7$  sufficiently small one sees that  $T_{37}$  is unbounded. The same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} (b_{38})''^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 72

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 381  
 GLOBAL EQUATIONS AND ITS CONCOMITANT CONDITIONALITIES into itself

382

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric 383

$$d\left( ((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G_{39}}, \widetilde{T_{39}})$  :

$$(\widetilde{G_{39}}, \widetilde{T_{39}}) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$$

It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_{36}^{(2)}| \leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} +$$

$$\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} +$$

$$(a''_{36})^{(7)}(T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} +$$

$$G_{36}^{(2)} |(a_{36}'')^{(7)}(T_{37}^{(1)}, s_{(36)}) - (a_{36}'')^{(7)}(T_{37}^{(2)}, s_{(36)})| e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} e^{(\widehat{M}_{36})^{(7)}s_{(36)}} ds_{(36)}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

384

$$\begin{aligned} & |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widehat{M}_{36})^{(7)}t} \leq \\ & \frac{1}{(\widehat{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a_{36}')^{(7)} + (\widehat{A}_{36})^{(7)} + \\ & (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a_{36}'')^{(7)}$  and  $(b_{36}'')^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  and  $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 385

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$ ,  $i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  386

From CONCATENATED GLOBAL EQUATIONS it results

$$\begin{aligned} G_i(t) & \geq G_i^0 e^{\left[ - \int_0^t \{ (a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)} \right]} \geq 0 \\ T_i(t) & \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$  and  $((\widehat{M}_{36})^{(7)})_3$  : 387

**Remark 3:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$ . indeed if

$G_{36} < ((\widehat{M}_{36})^{(7)})$  it follows  $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37}$  and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$ ,  $G_{38}$  and  $G_{36}$ ,  $G_{37}$  respectively.

**Remark 7:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$ . The proof is 388  
 analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below.

**Remark 5:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$  then 389  
 $T_{37} \rightarrow \infty$ .

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b'_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 390

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_7}$  By taking now  $\varepsilon_7$  sufficiently small one sees that  $T_{37}$  is  
 unbounded. The same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} (b''_{38})^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} \quad 391$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 392$$

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  : 393

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots 394

$$(a) \text{ of the equations } (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0 \quad 395$$

$$\text{and } (b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \text{ and} \quad 396$$

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  : 397

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the 398

$$\text{roots of the equations } (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0 \quad 399$$

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 400$$

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :- 401

(b) If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by 402

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \mathbf{if} (v_0)^{(2)} < (v_1)^{(2)} \quad 403$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \mathbf{if} (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 404$$

$$\text{and } \boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \mathbf{if} (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 405$$

and analogously 406

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \mathbf{if} (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \mathbf{if} (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } \boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \mathbf{if} (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 407$$

Then the solution satisfies the inequalities 408

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$  is defined 409

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 410$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[ e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 411$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[ e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 412$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 413$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[ e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 414$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[ e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :- 415

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)} \quad 416$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 417$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

418

**Behavior of the solutions**

419

If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

(a)  $\sigma_1^{(3)}, \sigma_2^{(3)}, \tau_1^{(3)}, \tau_2^{(3)}$  four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  :

420

(b) By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations  $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the

roots of the equations  $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :-

421

(c) If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

422

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution satisfies the inequalities

$$G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(s_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined

423

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(s_1)^{(3)}t}$$

424

$$\left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((s_1)^{(3)} - (p_{20})^{(3)} - (s_2)^{(3)})} \right) \left[ e^{((s_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(s_2)^{(3)}t} \right] + G_{22}^0 e^{-(s_2)^{(3)}t} \leq G_{22}(t) \leq$$

425

$$\frac{(a_{22})^{(3)}G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})} [e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 426$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 427$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} [e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 428$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} [e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :- 429

Where  $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

430

431

432

If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

(d)  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  : 433

(e) By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$  and  $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  : 434

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and  $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$  435

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :- 436

(f) If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$



$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

437  
438

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } \boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)}$  where  $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$  are defined respectively

Then the solution satisfies the inequalities

439  
440  
441  
442  
443  
444  
445

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

446  
447

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

448

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

449

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

450

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

451

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[ e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :-

452

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)} (\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

453

**Behavior of the solutions**

454

If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

(g)  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  :

455

(h) By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$  and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  :

456

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$  and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

(i) If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

457

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)}$  where  $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$  are defined respectively

Then the solution satisfies the inequalities

458

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t}$$

459

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \right. \\ \left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right) \quad \begin{matrix} 460 \\ 461 \end{matrix}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 462$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 463$$

$$\frac{(b_{30})^{(5)} T_{30}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 464$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :- 465

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions** 466

If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

(j)  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  : 467

(k) By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$  and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  : 468

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the

roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

(l) If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)} \quad 470$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and  $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 471

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and  $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)}$  where  $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$  are defined respectively

Then the solution satisfies the inequalities 472

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 473$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \right) \leq G_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[ e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \quad 474$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 475$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 476$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 477$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$ :- 478

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

479

If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

(m)  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying  
 $-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$   
 $-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  : 480

(n) By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$  481  
 and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  : 482

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

(o) If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 483

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

are defined respectively

Then the solution satisfies the inequalities

484

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined

485

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t}$$

486

$$\left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \right) \leq G_{38}(t) \leq \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

487

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

488

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

489

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq$$

490

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :-

491

$$\text{Where } (S_1)^{(7)} = (a_{36})^{(7)} (m_2)^{(7)} - (a'_{36})^{(7)}$$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

From GLOBAL EQUATIONS we obtain

492

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - \\ (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of**  $v^{(7)}$  :-

$$v^{(7)} = \frac{G_{36}}{G_{37}}$$

It follows

$$- \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq \\ - \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

(a) For  $0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}, \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

493

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

(b) If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case, 494

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

(c) If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$ , we obtain 495

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :-

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.



**Particular case :**

If  $(a''_{36})^{(7)} = (a''_{37})^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case .**

Analogously if  $(b''_{36})^{(7)} = (b''_{37})^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then

$(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

We can prove the following

If  $(a'_i)^{(7)}$  and  $(b'_i)^{(7)}$  are independent on  $t$ , and the conditions

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with  $(p_{36})^{(7)}, (r_{37})^{(7)}$  as defined are satisfied, then the system WITH THE SATISFACTION OF THE FOLLOWING PROPERTIES HAS A SOLUTION AS DERIVED BELOW.

.

**Particular case :**

If  $(a''_{16})^{(2)} = (a''_{17})^{(2)}$ , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if  $(b''_{16})^{(2)} = (b''_{17})^{(2)}$ , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of**  $v^{(3)}$  :- 501

$$v^{(3)} = \frac{G_{20}}{G_{21}}$$

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

502

From which one obtains

(a) For  $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

503

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of**  $(\bar{v}_1)^{(3)}$  :-

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

(b) If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case, 504

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

(c) If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$  , we obtain 505

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(3)}(t)$  :-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)} , \quad v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(3)}(t)$  :-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If  $(a''_{20})^{(3)} = (a''_{21})^{(3)}$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} = (v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b''_{20})^{(3)} = (b''_{21})^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$

506

: From GLOBAL EQUATIONS we obtain

507

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :-  $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

508

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$  :-

(d) For  $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_0)^{(4)}]t}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_0)^{(4)}]t}}, \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

509

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}}, \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

(e) If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case,

510

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_2)^{(4)}]t}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_2)^{(4)}]t}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

511

(f) If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$ , we obtain

512

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{24})^{(4)} = (a''_{25})^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$  **this also defines  $(v_0)^{(4)}$  for the special case .**

513

Analogously if  $(b''_{24})^{(4)} = (b''_{25})^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then  $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of  $(u_0)^{(4)}$ .**

514

From GLOBAL EQUATIONS we obtain

515

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \right) - (a''_{29})^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)}$  :-  $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left( (a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

(g) For  $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}, \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

516

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}, \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

(h) If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

517

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}} \leq (\bar{v}_1)^{(5)}$$

(i) If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$ , we obtain

518

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}} \leq (v_0)^{(5)}$$

519

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{28})^{(5)} = (a''_{29})^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b''_{28})^{(5)} = (b''_{29})^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of  $(u_0)^{(5)}$ .**

we obtain

520  
521

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of**  $v^{(6)}$  :- 
$$v^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-

(j) For  $0 < \left[ (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} \right] < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_0)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_0)^{(6)}]t}}, \quad (C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

522

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}, \quad (\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

523

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

(k) If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

524

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (\bar{v}_1)^{(6)}$$

(l) If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \left[ (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} \right]$ , we obtain

525

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{32})^{(6)} = (a''_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case .**

Analogously if  $(b''_{32})^{(6)} = (b''_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then  $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

526  
527

**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

(p)  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  :

528

(q) By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$  and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$  and

529

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  :

530.

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

(r) If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

531

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

are defined by 59 and 67 respectively

Then the solution of GLOBAL EQUATIONS satisfies the inequalities

532

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined



$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 533$$

$$\left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \right) \leq G_{38}(t) \leq \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t} \quad 534$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 535$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 536$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 537$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :- 538

Where  $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)} \quad 539$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

From CONCATENATED GLOBAL EQUATIONS we obtain 540

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of**  $v^{(7)}$  :-  $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq$$

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

(m) For  $0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}, \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

541

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}, \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

(n) If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case,

542

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (\bar{v}_1)^{(7)}$$

(o) If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$ , we obtain

543

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :-

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in CONCATENATED GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{36})^{(7)} = (a''_{37})^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case .**

Analogously if  $(b''_{36})^{(7)} = (b''_{37})^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then

$(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 544$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 545$$

has a unique positive solution , which is an equilibrium solution for the system 546

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 547$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 548$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 549$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 550$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 551$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 552$$

has a unique positive solution , which is an equilibrium solution for 553

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 554$$

$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0$	555
$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0$	556
$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0$	557
$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0$	558
$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0$	559
has a unique positive solution , which is an equilibrium solution	560
$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0$	561
$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0$	563
$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0$	564
$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27})]T_{24} = 0$	565
$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27})]T_{25} = 0$	566
$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27})]T_{26} = 0$	567
has a unique positive solution , which is an equilibrium solution for the system	568
$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0$	569
$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0$	570
$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0$	571
$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0$	572
$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0$	573
$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0$	574
has a unique positive solution , which is an equilibrium solution for the system	575
$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0$	576
$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0$	577
$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0$	578
$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0$	579

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 580$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 584$$

has a unique positive solution , which is an equilibrium solution for the system 582

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 583$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 584$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 585$$

586

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 587$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 588$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 589$$

has a unique positive solution , which is an equilibrium solution for the system 560

(a) Indeed the first two equations have a nontrivial solution  $G_{36}, G_{37}$  if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

**Definition and uniqueness of  $T_{37}^*$  :-** 561

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(7)}(T_{37})$  being increasing, it follows that there exists a unique  $T_{37}^*$  for which  $f(T_{37}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

(e) By the same argument, the equations( SOLUTIONAL) admit solutions  $G_{36}, G_{37}$  if

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in  $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{37}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{37}^*$  such that  $\varphi(G^*) = 0$  562

Finally we obtain the unique solution OF THE SYSTEM

$G_{37}^*$  given by  $\varphi((G_{39})^*) = 0, T_{37}^*$  given by  $f(T_{37}^*) = 0$  and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]}, \quad G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]}, \quad T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]} \quad 563$$

**Definition and uniqueness of  $T_{21}^*$  :-** 564

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]}, \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$
565

**Definition and uniqueness of  $T_{25}^*$  :-** 566

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]}, \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$
567

**Definition and uniqueness of  $T_{29}^*$  :-** 567

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]}, \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$
568

**Definition and uniqueness of  $T_{33}^*$  :-** 568

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

(f) By the same argument, the equations 92,93 admit solutions  $G_{13}, G_{14}$  if 569

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

(g) By the same argument, the equations 92,93 admit solutions  $G_{16}, G_{17}$  if 570

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi((G_{19})^*) = 0$  571

(a) By the same argument, the concatenated equations admit solutions  $G_{20}, G_{21}$  if 572

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$  573

(b) By the same argument, the equations of modules admit solutions  $G_{24}, G_{25}$  if 574

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

(c) By the same argument, the equations (modules) admit solutions  $G_{28}, G_{29}$  if 575

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

(d) By the same argument, the equations (modules) admit solutions  $G_{32}, G_{33}$  if 578

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - 579$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0 580$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G^*) = 0$  581

Finally we obtain the unique solution of 89 to 94 582

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]}, \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}, \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 583

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0, T_{17}^*$  given by  $f(T_{17}^*) = 0$  and 584

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]}, \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} 585$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]}, \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} 586$$

Obviously, these values represent an equilibrium solution 587

Finally we obtain the unique solution 588

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0, T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]}, \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]}, \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 589

$G_{25}^*$  given by  $\varphi(G_{27}) = 0, T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]}, \quad G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]}, \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} 590$$



Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 591

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$ ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^*)]} , T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^*)]}$$
592

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 593

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$ ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} , G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^*)]} , T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^*)]}$$
594

Obviously, these values represent an equilibrium solution

#### ASYMPTOTIC STABILITY ANALYSIS 595

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$
596

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 597

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14}$$
598

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14}$$
599

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14}$$
600

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*\mathbb{G}_j$$
601

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*\mathbb{G}_j$$
602

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*\mathbb{G}_j$$
603

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  604

Belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote 605

**Definition of  $G_i, T_i$  :-**

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i \quad \text{606}$$

$$\frac{\partial(a'_{17})^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial(b'_i)^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad \text{607}$$

taking into account equations (global) and neglecting the terms of power 2, we obtain 608

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad \text{609}$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad \text{610}$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad \text{611}$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*G_j \quad \text{612}$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*G_j \quad \text{613}$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*G_j \quad \text{614}$$

If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(3)}$  and  $(b'_i)^{(3)}$  belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable 615

Denote

**Definition of  $G_i, T_i$  :-**

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{21})^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \quad \frac{\partial(b'_i)^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij}$$

616

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 617

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad \text{618}$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad \text{619}$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad \text{6120}$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^*G_j \quad \text{621}$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^*G_j \quad \text{622}$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^*G_j \quad 623$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable 624

Denote

**Definition of**  $G_i, T_i$  :- 625

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 626

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^*T_{25} \quad 627$$

$$\frac{dG_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^*T_{25} \quad 628$$

$$\frac{dG_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^*T_{25} \quad 629$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*G_j \quad 630$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*G_j \quad 631$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*G_j \quad 632$$

633

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :- 634

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 635

$$\frac{dG_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})G_{28} + (a_{28})^{(5)}G_{29} - (q_{28})^{(5)}G_{28}^*T_{29} \quad 636$$

$$\frac{dG_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})G_{29} + (a_{29})^{(5)}G_{28} - (q_{29})^{(5)}G_{29}^*T_{29} \quad 637$$

$$\frac{dG_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^*T_{29} \quad 638$$

$$\frac{dT_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*G_j \quad 639$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*G_j \quad 640$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*G_j \quad 641$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable 642

Denote

**Definition of**  $G_i, T_i$  :- 643

$$G_i = G_i^* + G_i \quad , T_i = T_i^* + T_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}} (T_{33}^*) = (q_{33})^{(6)} \quad , \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j} ((G_{35})^*) = s_{ij}$$

Then taking into account equations(global) and neglecting the terms of power 2, we obtain 644

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 645$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 646$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 647$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^*G_j \quad 648$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^*G_j \quad 649$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^*G_j \quad 650$$

Obviously, these values represent an equilibrium solution of 79,20,36,22,23, 651

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  belong to  $C^{(7)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 652

$$G_i = G_i^* + G_i \quad , T_i = T_i^* + T_i$$

653

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations(SOLUTIONAL) and neglecting the terms of power 2, we obtain 654

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 656$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 657$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 658$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^*G_j \quad 659$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^*G_j \quad 660$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^*G_j \quad 661$$

2. 662

The characteristic equation of this system is

$$\begin{aligned} & ((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ & \left[ \left( ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right) \right] \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^* \right) \\ & + \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^* \right) \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^* \right) \\ & \left( ((\lambda)^{(1)})^2 + \left( (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)} \right) (\lambda)^{(1)} \right) \\ & \left( ((\lambda)^{(1)})^2 + \left( (b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)} \right) (\lambda)^{(1)} \right) \end{aligned}$$

$$\begin{aligned}
 &+ \left( (\lambda)^{(1)} \right)^2 + \left( (a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)} \right) (\lambda)^{(1)} (q_{15})^{(1)} G_{15} \\
 &+ \left( (\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)} \right) \left( (a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^* \right) \\
 &\left( \left( (\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)} \right) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &\left( (\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)} \right) \left\{ \left( (\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)} \right) \right. \\
 &\left[ \left( \left( (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \right] \\
 &\left( \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \right) \\
 &+ \left( \left( (\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)} \right) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \right) \\
 &\left( \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \right) \\
 &\left( \left( (\lambda)^{(2)} \right)^2 + \left( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \right) \\
 &\left( \left( (\lambda)^{(2)} \right)^2 + \left( (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \right) \\
 &+ \left( \left( (\lambda)^{(2)} \right)^2 + \left( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \right) (q_{18})^{(2)} G_{18} \\
 &+ \left( (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) \left( (a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \\
 &\left( \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right) \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 &\left( (\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \left\{ \left( (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \right) \right. \\
 &\left[ \left( \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right) \right] \\
 &\left( \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \right) \\
 &+ \left( \left( (\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \right)
 \end{aligned}$$

$$\left( \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \right)$$

$$\left( ((\lambda)^{(3)})^2 + \left( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \right)$$

$$\left( ((\lambda)^{(3)})^2 + \left( (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \right)$$

$$+ \left( ((\lambda)^{(3)})^2 + \left( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \right) (q_{22})^{(3)} G_{22}$$

$$+ \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) \left( (a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^* \right)$$

$$\left( \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0$$

+

$$\left( (\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)} \right) \left\{ \left( (\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \right) \right.$$

$$\left[ \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right]$$

$$\left( \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \right)$$

$$+ \left( (\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)} \right) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \right)$$

$$\left( \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \right)$$

$$\left( ((\lambda)^{(4)})^2 + \left( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \right)$$

$$\left( ((\lambda)^{(4)})^2 + \left( (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} \right) (\lambda)^{(4)} \right)$$

$$+ \left( ((\lambda)^{(4)})^2 + \left( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26}$$

$$+ \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) \left( (a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^* \right)$$

$$\left( \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \} = 0$$

+

$$\left( (\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)} \right) \left\{ \left( (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \right) \right.$$

$$\left[ \left( \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right) \right. \\
\left. \left( \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \right) \right. \\
+ \left( \left( (\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)} \right) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \right) \\
\left. \left( \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \right) \right. \\
\left( (\lambda)^{(5)} \right)^2 + \left( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} \right) (\lambda)^{(5)} \\
\left( (\lambda)^{(5)} \right)^2 + \left( (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} \right) (\lambda)^{(5)} \\
+ \left( (\lambda)^{(5)} \right)^2 + \left( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} \right) (\lambda)^{(5)} (q_{30})^{(5)} G_{30} \\
+ \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) \left( (a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^* \right) \\
\left. \left( \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \right\} = 0$$

+

$$\left( (\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)} \right) \left\{ \left( (\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)} \right) \right. \\
\left[ \left( \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \right. \\
\left. \left( \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \right. \\
+ \left( \left( (\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)} \right) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
\left. \left( \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \right. \\
\left( (\lambda)^{(6)} \right)^2 + \left( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} \\
\left( (\lambda)^{(6)} \right)^2 + \left( (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\
+ \left( (\lambda)^{(6)} \right)^2 + \left( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
+ \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left( (a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \left. \right\}$$



$$\left( \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0$$

+

$$\begin{aligned} & \left( (\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \{ (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \\ & \left[ \left( (\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \\ & \left( (\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)} \right) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \\ & + \left( (\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)} \right) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left( (\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)} \right) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \\ & \left( (\lambda)^{(7)} \right)^2 + \left( (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left( (\lambda)^{(7)} \right)^2 + \left( (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left( (\lambda)^{(7)} \right)^2 + \left( (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + \left( (\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) \left( (a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\ & \left( (\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)} \right) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \} = 0 \end{aligned}$$

#### REFERENCES

- =====
- (1) A HAIMOVICI: "On the growth of a two species ecological system divided on age groups". Tensor, Vol 37 (1982), Commemoration volume dedicated to Professor Akitsugu Kawaguchi on his 80<sup>th</sup> birthday
  - (2) FRTJOF CAPRA: "The web of life" Flamingo, Harper Collins See "Dissipative structures" pages 172-188
  - (3) HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [http://www.eolss.net
  - (4) MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204, doi:10.1029/2005JD006097
  - (5) STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" J. Atmos. Sci., 53, 980-

1006

(6)FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: [10.1038/nature09314](https://doi.org/10.1038/nature09314), Published 12-Aug 2010

(7)HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [<http://www.eolss.net>

(8)MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", *J. Geophys. Res.*, 111, D17204, doi:10.1029/2005JD006097

(8A)STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" *J. Atmos. Sci.*, 53, 980-1006

(8B)FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: [10.1038/nature09314](https://doi.org/10.1038/nature09314), Published 12-Aug 2010

(9)<sup>a b c</sup> Einstein, A. (1905), "Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?", *Annalen der Physik* **18**:  
639 Bibcode 1905AnP...323..639E, DOI:10.1002/andp.19053231314. See also the English translation.

(10)<sup>a b</sup> Paul Allen Tipler, Ralph A. Llewellyn (2003-01), *Modern Physics*, W. H. Freeman and Company, pp. 87–88, ISBN 0-7167-4345-0

(11)<sup>a b</sup> Rainville, S. et al. World Year of Physics: A direct test of  $E=mc^2$ . *Nature* 438, 1096-1097 (22 December 2005) | doi: 10.1038/4381096a; Published online 21 December 2005.

(12)<sup>a</sup> In F. Fernflores. The Equivalence of Mass and Energy. Stanford Encyclopedia of Philosophy

(13)<sup>a</sup> Note that the relativistic mass, in contrast to the rest mass  $m_0$ , is not a relativistic invariant, and that the velocity is not a Minkowski four-vector, in contrast to the quantity  $dx^\mu$ , where  $dx^\mu$  is the differential of the proper time. However, the energy-momentum four-vector is a genuine Minkowski four-vector, and the intrinsic origin of the square-root in the definition of the relativistic mass is the distinction between  $d\tau$  and  $dt$ .

(14)<sup>a</sup> Relativity DeMystified, D. McMahon, Mc Graw Hill (USA), 2006, ISBN 0-07-145545-0

(15)<sup>a</sup> Dynamics and Relativity, J.R. Forshaw, A.G. Smith, Wiley, 2009, ISBN 978-0-470-01460-8

(16)<sup>a</sup> Hans, H. S.; Puri, S. P. (2003). *Mechanics* (2 ed.). Tata McGraw-Hill. p. 433. ISBN 0-07-047360-9., Chapter 12 page 433

(17)<sup>a</sup> E. F. Taylor and J. A. Wheeler, **Spacetime Physics**, W.H. Freeman and Co., NY. 1992. ISBN 0-7167-2327-1, see pp. 248-9 for discussion of mass remaining constant after detonation of nuclear bombs, until heat is allowed to escape.

(18)<sup>a</sup> Mould, Richard A. (2002). *Basic relativity* (2 ed.). Springer. p. 126. ISBN 0-387-95210-1., Chapter 5 page 126

(19)<sup>a</sup> Chow, Tail L. (2006). *Introduction to electromagnetic theory: a modern perspective*. Jones & Bartlett Learning. p. 392. ISBN 0-7637-3827-1., Chapter 10 page 392

(20)<sup>a</sup> [2] Cockcroft-Walton experiment

(21)<sup>a b c</sup> Conversions used: 1956 International (Steam) Table (IT) values where one calorie

$\equiv 4.1868 \text{ J}$  and one BTU  $\equiv 1055.05585262 \text{ J}$ . Weapons designers' conversion value of one gram TNT  $\equiv 1000$  calories used.

(22)  $\triangle$  Assuming the dam is generating at its peak capacity of 6,809 MW.

(23)  $\triangle$  Assuming a 90/10 alloy of Pt/Ir by weight, a  $C_p$  of 25.9 for Pt and 25.1 for Ir, a Pt-dominated average  $C_p$  of 25.8, 5.134 moles of metal, and  $132 \text{ J.K}^{-1}$  for the prototype. A variation of  $\pm 1.5$  picograms is of course, much smaller than the actual uncertainty in the mass of the international prototype, which are  $\pm 2$  micrograms.

(24)  $\triangle$  [31] Article on Earth rotation energy. Divided by  $c^2$ .

(25)  $\wedge^{a b}$  Earth's gravitational self-energy is  $4.6 \times 10^{-10}$  that of Earth's total mass, or 2.7 trillion metric tons. Citation: *The Apache Point Observatory Lunar Laser-Ranging Operation (APOLLO)*, T. W. Murphy, Jr. *et al.* University of Washington, Dept. of Physics (132 kB PDF, [here.](#)).

(26)  $\triangle$  There is usually more than one possible way to define a field energy, because any field can be made to couple to gravity in many different ways. By general scaling arguments, the correct answer at everyday distances, which are long compared to the quantum gravity scale, should be *minimal coupling*, which means that no powers of the curvature tensor appear. Any non-minimal couplings, along with other higher order terms, are presumably only determined by a theory of quantum gravity, and within string theory, they only start to contribute to experiments at the string scale.

(27)  $\triangle$  G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Physical Review D* 14:3432–3450 (1976).

(28)  $\triangle$  A. Belavin, A. M. Polyakov, A. Schwarz, Yu. Tyupkin, "Pseudoparticle Solutions to Yang Mills Equations", *Physics Letters* 59B:85 (1975).

(29)  $\triangle$  F. Klinkhammer, N. Manton, "A Saddle Point Solution in the Weinberg Salam Theory", *Physical Review D* 30:2212.

(30)  $\triangle$  Rubakov V. A. "Monopole Catalysis of Proton Decay", *Reports on Progress in Physics* 51:189–241 (1988).

(31)  $\triangle$  S.W. Hawking "Black Holes Explosions?" *Nature* 248:30 (1974).

(32)  $\wedge$  Einstein, A. (1905), "Zur Elektrodynamik bewegter Körper." (PDF), *Annalen der Physik* 17: 891–921, Bibcode 1905AnP...322...891E, DOI:10.1002/andp.19053221004. English translation.

(33)<sup>△</sup> See e.g. Lev B. Okun, *The concept of Mass*, *Physics Today* **42** (6), June 1969, p. 31–36, [http://www.physicstoday.org/vol-42/iss-6/vol42no6p31\\_36.pdf](http://www.physicstoday.org/vol-42/iss-6/vol42no6p31_36.pdf)

(34)<sup>△</sup> Max Jammer (1999), *Concepts of mass in contemporary physics and philosophy*, Princeton University Press, p. 51, ISBN 0-691-01017-X

(35)<sup>△</sup> Eriksen, Erik; Vøyenli, Kjell (1976), "The classical and relativistic concepts of mass", *Foundations of Physics* (Springer) **6**: 115–124, Bibcode 1976FoPh....6..115E, DOI:10.1007/BF00708670

(36)<sup>△</sup> <sup>a b</sup> Janssen, M., Mecklenburg, M. (2007), *From classical to relativistic mechanics: Electromagnetic models of the electron.*, in V. F. Hendricks, et al., , *Interactions: Mathematics, Physics and Philosophy* (Dordrecht: Springer): 65–134

(37)<sup>△</sup> <sup>a b</sup> Whittaker, E.T. (1951–1953), 2. Edition: *A History of the theories of aether and electricity*, vol. 1: *The classical theories* / vol. 2: *The modern theories 1900–1926*, London: Nelson

(38)<sup>△</sup> Miller, Arthur I. (1981), *Albert Einstein's special theory of relativity. Emergence (1905) and early interpretation (1905–1911)*, Reading: Addison–Wesley, ISBN 0-201-04679-2

(39)<sup>△</sup> <sup>a b</sup> Darrigol, O. (2005), "The Genesis of the theory of relativity." (PDF), *Séminaire Poincaré* **1**: 1–22

(40)<sup>△</sup> Philip Ball (Aug 23, 2011). "Did Einstein discover  $E = mc^2$ ?" *Physics World*.

(41)<sup>△</sup> Ives, Herbert E. (1952), "Derivation of the mass-energy relation", *Journal of the Optical Society of America* **42** (8): 540–543, DOI:10.1364/JOSA.42.000540

(42)<sup>△</sup> Jammer, Max (1961/1997). *Concepts of Mass in Classical and Modern Physics*. New York: Dover. ISBN 0-486-29998-8.

(43)<sup>△</sup> Stachel, John; Torretti, Roberto (1982), "Einstein's first derivation of mass-energy equivalence", *American Journal of Physics* **50** (8): 760–763, Bibcode1982AmJPh..50..760S, DOI:10.1119/1.12764

(44)<sup>△</sup> Ohanian, Hans (2008), "Did Einstein prove  $E=mc^2$ ?", *Studies In History and Philosophy of Science Part B* **40** (2): 167–173, arXiv:0805.1400, DOI:10.1016/j.shpsb.2009.03.002

---

(45)^ Hecht, Eugene (2011), "How Einstein confirmed  $E=mc^2$ ", *American Journal of Physics* **79** (6): 591–600, Bibcode 2011AmJPh..79..591H, DOI:10.1119/1.3549223

(46)^ Rohrlich, Fritz (1990), "An elementary derivation of  $E=mc^2$ ", *American Journal of Physics* **58** (4): 348–349, Bibcode 1990AmJPh..58..348R, DOI:10.1119/1.16168

(47) (1996). *Lise Meitner: A Life in Physics*. California Studies in the History of Science. **13**. Berkeley: University of California Press. pp. 236–237. ISBN 0-520-20860-

---

(48)^ UIBK.ac.at

(49)^ J. J. L. Morton; *et al.* (2008). "Solid-state quantum memory using the  $^{31}\text{P}$  nuclear spin". *Nature* **455** (7216): 1085–1088. Bibcode 2008Natur.455.1085M. DOI:10.1038/nature07295.

(50)^ S. Weisner (1983). "Conjugate coding". *Association of Computing Machinery, Special Interest Group in Algorithms and Computation Theory* **15**: 78–88.

(51)^ A. Zeilinger, *Dance of the Photons: From Einstein to Quantum Teleportation*, Farrar, Straus & Giroux, New York, 2010, pp. 189, 192, ISBN 0374239665

(52)^ B. Schumacher (1995). "Quantum coding". *Physical Review A* **51** (4): 2738–2747. Bibcode 1995PhRvA..51.2738S. DOI:10.1103/PhysRevA.51.2738.

(53) Delamotte, Bertrand; *A hint of renormalization*, *American Journal of Physics* **72** (2004) pp. 170–184. Beautiful elementary introduction to the ideas, no prior knowledge of field theory being necessary. Full text available at: [hep-th/0212049](http://hep-th/0212049)

(54) Baez, John; *Renormalization Made Easy*, (2005). A qualitative introduction to the subject.

(55) Blechman, Andrew E. ; *Renormalization: Our Greatly Misunderstood Friend*, (2002). Summary of a lecture; has more information about specific regularization and divergence-subtraction schemes.

(56) Cao, Tian Yu & Schweber, Silvan S. ; *The Conceptual Foundations and the Philosophical Aspects of Renormalization Theory*, *Synthese*, **97**(1) (1993), 33–108.

(57) Shirkov, Dmitry; *Fifty Years of the Renormalization Group*, *C.E.R.N. Courier* **41**(7) (2001). Full

text available at: *I.O.P Magazines*.

(58)E. Elizalde; *Zeta regularization techniques with Applications*.

(59)N. N. Bogoliubov, D. V. Shirkov (1959): *The Theory of Quantized Fields*. New York, Interscience. The first text-book on the renormalization group theory.

(60)Ryder, Lewis H. ; *Quantum Field Theory* (Cambridge University Press, 1985), ISBN 0-521-33859-X Highly readable textbook, certainly the best introduction to relativistic Q.F.T. for particle physics.

(61)Zee, Anthony; *Quantum Field Theory in a Nutshell*, Princeton University Press (2003) ISBN 0-691-01019-6. Another excellent textbook on Q.F.T.

(62)Weinberg, Steven; *The Quantum Theory of Fields* (3 volumes) Cambridge University Press (1995). A monumental treatise on Q.F.T. written by a leading expert, *Nobel laureate 1979*.

(63)Pokorski, Stefan; *Gauge Field Theories*, Cambridge University Press (1987) ISBN 0-521-47816-2.

(64)'t Hooft, Gerard; *The Glorious Days of Physics – Renormalization of Gauge theories*, lecture given at Erice (August/September 1998) by the *Nobel laureate 1999* . Full text available at: *hep-th/9812203*.

(65)Rivasseau, Vincent; *An introduction to renormalization*, Poincaré Seminar (Paris, Oct. 12, 2002), published in: Duplantier, Bertrand; Rivasseau, Vincent (Eds.) ; *Poincaré Seminar 2002*, Progress in Mathematical Physics 30, Birkhäuser (2003) ISBN 3-7643-0579-7. Full text available in *PostScript*.

(66) Rivasseau, Vincent; *From perturbative to constructive renormalization*, Princeton University Press (1991) ISBN 0-691-08530-7. Full text available in *PostScript*.

(67) Iagolnitzer, Daniel & Magnen, J. ; *Renormalization group analysis*, Encyclopaedia of Mathematics, Kluwer Academic Publisher (1996). Full text available in PostScript and pdfhere.

(68) Scharf, Günter; *Finite quantum electrodynamics: The casual approach*, Springer Verlag Berlin Heidelberg New York (1995) ISBN 3-540-60142-2.

(69)A. S. Švarc (Albert Schwarz), Математические основы квантовой теории поля, (Mathematical aspects of quantum field theory), Atomizdat, Moscow, 1975. 368 pp.

(70) A. N. Vasil'ev *The Field Theoretic Renormalization Group in Critical Behavior Theory and Stochastic Dynamics* (Routledge Chapman & Hall 2004); ISBN 978-0-415-31002-4

(71) Nigel Goldenfeld ; *Lectures on Phase Transitions and the Renormalization Group*, Frontiers in Physics 85, West view Press (June, 1992) ISBN 0-201-55409-7. Covering the elementary aspects of the physics of phase's transitions and the renormalization group, this popular book emphasizes understanding and clarity rather than technical manipulations.

(72) Zinn-Justin, Jean; *Quantum Field Theory and Critical Phenomena*, Oxford University Press (4th edition – 2002) ISBN 0-19-850923-5. A masterpiece on applications of renormalization methods to the calculation of critical exponents in statistical mechanics, following Wilson's ideas (Kenneth Wilson was *Nobel laureate 1982*).

(73)Zinn-Justin, Jean; *Phase Transitions & Renormalization Group: from Theory to Numbers*, Poincaré Seminar (Paris, Oct. 12, 2002), published in: Duplantier, Bertrand; Rivasseau, Vincent (Eds.); *Poincaré Seminar 2002*, Progress in Mathematical Physics 30, Birkhäuser (2003) ISBN 3-7643-0579-7. Full text available in *PostScript*.

(74 )Domb, Cyril; *The Critical Point: A Historical Introduction to the Modern Theory of Critical Phenomena*, CRC Press (March, 1996) ISBN 0-7484-0435-X.

(75)Brown, Laurie M. (Ed.) ; *Renormalization: From Lorentz to Landau (and Beyond)*, Springer-Verlag (New York-1993) ISBN 0-387-97933-6.

(76) Cardy, John; *Scaling and Renormalization in Statistical Physics*, Cambridge University Press (1996) ISBN 0-521-49959-3.

(77)Shirkov, Dmitry; *The Bogoliubov Renormalization Group*, JINR Communication E2-96-15 (1996). Full text available at: [hep-th/9602024](http://hep-th/9602024)

(78)Zinn Justin, Jean; *Renormalization and renormalization group: From the discovery of UV divergences to the concept of effective field theories*, in: de Witt-Morette C., Zuber J.-B. (eds), Proceedings of the NATO ASI on *Quantum Field Theory: Perspective and Prospective*, June 15–26, 1998, Les Houches, France, Kluwer Academic Publishers, NATO ASI Series C 530, 375–388 (1999). Full text available in *PostScript*.

(79) Connes, Alain; *Symmetries Galoisiennes & Renormalisation*, Poincaré Seminar (Paris, Oct. 12, 2002), published in: Duplantier, Bertrand; Rivasseau, Vincent (Eds.) ; *Poincaré Seminar 2002*,



---

Progress in Mathematical Physics 30, Birkhäuser (2003) ISBN 3-7643-0579-7. French mathematician *Alain Connes* (Fields medallist 1982) describes the mathematical underlying structure (the Hopf algebra) of renormalization, and its link to the Riemann-Hilbert problem. Full text (in French) available at *math/0211199v1*.

---

(80)^ <http://www.ingentaconnect.com/con/klu/math/1996/00000037/00000004/00091778>

(81)^ F. J. Dyson, *Phys. Rev.* **85** (1952) 631.

(82)^ A. W. Stern, *Science* **116** (1952) 493.

(83)^ P.A.M. Dirac, "The Evolution of the Physicist's Picture of Nature," in *Scientific American*, May 1963, p. 53.

(84)^ Kragh, Helge ; *Dirac: A scientific biography*, CUP 1990, p. 184

(85)^ Feynman, Richard P. ; *QED, The Strange Theory of Light and Matter*, Penguin 1990, p. 128

(86)^ C.J.Isham, A.Salam and J.Strathdee, 'Infinity Suppression Gravity Modified Quantum Electrodynamics,' *Phys. Rev. D5*, 2548 (1972)

(87)^ Ryder, Lewis. *Quantum Field Theory*, page 390 (Cambridge University Press 1996).

---

#### Acknowledgments:

**The introduction is a collection of information from various articles, Books, News Paper reports, Home Pages Of authors, Journal Reviews, Nature 's Letters, Article Abstracts, Research papers, Abstracts Of Research Papers, Stanford Encyclopedia, Web Pages, Ask a Physicist Column, Deliberations with Professors, the internet including Wikipedia. We acknowledge all authors who have contributed to the same. In the eventuality of the fact that there has been any act of omission on the part of the authors, we regret with great deal of compunction, contrition, regret, trepidation and remorse. As Newton said, it is only because erudite and eminent people allowed one to piggy ride on their backs; probably an attempt has been made to look slightly further. Once again, it is stated that the references are only illustrative and not comprehensive**

**First Author:** <sup>1</sup>**Mr. K. N.Prasanna Kumar** has three doctorates one each in Mathematics, Economics, Political Science. Thesis was based on Mathematical Modeling. He was recently awarded D.litt. for his work on 'Mathematical Models in Political Science'--- Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Corresponding [Author:drknpkumar@gmail.com](mailto:drknpkumar@gmail.com)

**Second Author:** <sup>2</sup>**Prof. B.S Kiranagi** is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150 publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent publication history.-- UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

**Third Author:** <sup>3</sup>**Prof. C.S. Bagewadi** is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. Several Books on Differential Geometry, Differential Equations are coauthored by him--- Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India

=====

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### **IISTE Knowledge Sharing Partners**

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

