Advances in Physics Theories and Applications
ISSN 2224-719X (Paper) ISSN 2225-0638 (Online)
Vol 10, 2012

Attenuation of Mesoscopic Echoes In A Spin-Chain Channel, Anti-Zeno Polarization Control of Spin Ensembles, Quantum Information, Quantum Wells, Hadrons, Spatial Profiles of Fusion Product Flux, Proton Spin-Lattice Relaxation In A Liquid Crystal , Quantum Confinement And Lattice Gauge Field Theories Et Al: Deo Ac Veritati-Deus Ex Machina Model

[1]Dr K N Prasanna Kumar, [2]Prof B S Kiranagi [3]Prof C S Bagewadi

[1]Dr K N Prasanna Kumar, Post doctoral researcher, Dr KNP Kumar has three PhD's, one each in Mathematics, Economics and Political science and a D.Litt. in Political Science, Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Correspondence Mail id : drknpkumar@gmail.com

[2]Prof B S Kiranagi, UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

[3]Prof C S Bagewadi, Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India

Keywords: Decoherence ,Mesoscopic Echoes , Spin Ensembles , Zeno effect, Quantum measurements, quantum information, Localization effects, many-spin quantum states, Quantum Zeno Effect , TOCSY, Intersubband Transitions, particles , torque , colloidal particle, liquid crystal host , topological defects, fusion product flux, dynamic trap , Nonperturbative Effects , Quantum Confinement, Lattice Gauge Field Theories, Quark-Antiquark Potential , Radiative corrections, Transport Theory , Quantum Tunneling, Heisenberg Model, Ground States , Quantum Monte Carlo Method , Black Holes , Stability Of Geons, Boson Stars, Neutrons, Higher Derivative Gravity, Superconductivity, Holographic Anomalies

This paper is organized as follows

Section 1: Decoherence Vis-à-vis Attenuation Of Mesoscopic Echoes In A Spin-Chain Channel, Spin Ladder Vis-à-vis Mesoscopic Echoes Et Al, A Thunderstorm Thermistor-Verkrampte Reactionary Model

Section 2: Zeno and Anti-Zeno Polarization Control of Spin Ensembles Vis a Vis Induced Dephasing, Zeno effect Vis a Vis anti-Zeno effect, Quantum measurements Vis a Vis quantum information: A Testamentary Dictyoptera-Synecdochal Syncretism Model

Section 3 : Localization effects Vis-à-vis decoherence in superpositions of many-spin quantum states, Anderson localization Vis-à-vis thermalization, quantum information Vis-à-vis state transfer: A Sustenational sussurationer-Poindexter poinsettia Model

Section 4: Quantum Zeno Effect Vis A Vis TOCSY, Quantum Information Vis-à-vis Spin-Spin Couplings Shift-Driven Modulations Vis-à-vis Spin-Echo Signals: A Burlesque Moccasin – Harmonious Key Note Model

Section 5: Polaritons Vis-à-vis Intersubband, Quantum Wells Vis-à-vis Hadrons, Signatures Of The Ultra Strong Light-Mater Coupling Regime Vis-à-vis Intersubband Transitions: A Apriori Determination And Aposteori Differentiation Model

Section 6: Drag on particles in a nematic suspension Vis-à-vis moving nematic-isotropic interface, Defect

structures and torque on an elongated colloidal particle immersed Vis-à-vis a liquid crystal host with topological defects, Entropic torque. Vis-à-vis depletion effects: A Herpestidae Mongoose and Ineluctable Aquiline Model

Section 7: Spatial profiles of fusion product flux Vis-à-vis the gas dynamic trap with deuterium neutral beam injection, Fusion regimes of axisymmetric open system Vis-à-vis powerful injection of fast particles, Enhancement of order fluctuations Vis-à-vis nematic liquid crystal by sonication (nematic, acoustic): A Filibustered Conventicler- Serene canopies and Trellises Model

Section 8: Proton field-cycling nuclear magnetic resonance relaxometry Vis-à-vis the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication, Proton spin-lattice relaxation in a liquid crystalvis Vis-à-vis Aerosol complex above the bulk isotropization temperature:

Section 9: Nonperturbative Effects Vis-à-vis Quantum Confinement And Lattice Gauge Field Theories, 3D SU(N) Georgi-Glashow Model Vis-à-vis Magnetic Monopoles Confining Strings, Quark Confinement Vis-à-vis String And Field Theories At Finite Temperature: A Reinforcing Template-Dissipation Denosjour Model

Section 10: Dual Abelian-Higgs-Type Theories Vis-à-vis Aharonov-Bohm Effect, Luescher Term In The Quark-Antiquark Potential Vis-à-vis Nonperturbative Effects, Mixed Heavy-Quark-Gluon(Nonperturbative Effects) Bose Einstein Condensate Vis-à-vis In The Stochastic Vacuum Model , Et Al, :A Exogenous Endomorph-Arrondissement Arrogate Model

Section 11: Radiative corrections to scalar fermion pair production Vis-à-vis High-energy e+ e- collisions, Radiative corrections Vis-à-vis Supersymmetry, Higgs decays in the two Higgs doublet model: Large quantum effects Vis-à-vis the decoupling regime: A Dorcas gazelle-Weeny Weaver Model

Section 12: Transport Theory Vis-à-vis Quantum Tunneling, Heisenberg Model Vis-à-vis Exchange Interactions (Electron); Ground States Vis-à-vis Spin Fluctuations, Mesoscopic Systems Vis-à-vis Green's Function Methods And Band Structure Et Al,: Coextensive Representations And Apriori Determination Thereof:

Section 13: Quantum Monte Carlo Method Vis-à-vis Molecular Nanodevices, Anderson-Holstein Model Vis-à-vis Molecular Transistor, Green-Function Approach Vis-à-vis Transport Phenomena In Quantum Pumps: A Crotchet-Conundrum Model

Section 14: Stability Of Geons, Boson Stars And Black Holes (Asymptotically Nonlinear Solutions Which Are Nonlinearly Stable Vis-à-vis Black Hole Binaries, Wilhelm Weber And The Surface Charges Vis-à-vis Resistive Conductors Carrying Steady Currents Surface: Model With primary Focus On The Homologous Receptiveness Of Differentially Interactional And Instrumental Activity

Section 15: Holographic Anomalies And Higher Derivative Gravity, Two-Hadron Semi-Inclusive Production Including Subleading Twist Et Al,: A Rational Leibneizism And Socratic Subjectivity Model

Section 16: Superconductivity and Neutrons, magnetic ordering on dynamical conductivity, Transport Properties and Magnetic properties Et Al: Predicational Interiority and Primordial Exactitude thereof

Acknowledgements:

It is to be stated in unmistakable, unambiguous, unequivocal terms that_the contential constitutions, presciential dynamics, predicational anteriority, character constitution, ontological consonance, primordial exactitude, Acolytish representation, propositional subsistence, substantive corporeality, apocryphal aneurism, atrophied asseverations, functional topology, manifestation of written histories, perceptual field, form-background, theme-potentialities, predicational pronouncements, profiles and unity of perspectives. principal frontier of diurnal dynamics, corporeal depth and sonorous continuum, locus and focus of essence, sense and expression, hypostatized and hyper dynamic confederation, of the Introduction, has been taken from various sources, such as: Google Search, Encyclopedia, Stanford Encyclopedia, Deliberations with the Professors, Authors Home Page, various Columns like Ask a Physicist, Reports of Scientific American, and Newspapers reports. Follow up notes of ATLAS, TOTEM, ALICE, and various

photographs from NASA or Wikipedia. This is purely done spending my own terminal benefits and without any sense of profit motivation.

I have put all concerted efforts, sustained struggle, and protracted endeavor, to put on recordable evidence the sources to the widest commonalty spread. In the eventuality of the fact that there had been any act of omission is attributable to the fact that it has happened inadvertently, in deliberately, and despite eminent and erudite people helping me to piggy ride on their backs, I probably might not have come to their expectations and produced an average work. Singlehandedly, I have browsed millions of pages, and there are possibilities of repetition, ascribable to the forgetfulness of what has been done before, for I never looked back, with diligence and enterprise. Emphasis also shifted sometimes from attention to the problem solving to style. Despite all these kinks in my amour, if I have been able to put forth any worthy piece of work, credit goes to all the people and not to me at all. I beg pardon on that count. References are only illustrative and not exhaustive. I once again thank all the people who have contributed for this work, and frankly, there is no question of sense of belonging what with a feeling of whole wide ocean waiting to be explored, one tends only impatient, and astutely truculent, and serenadingly pushy to consummate, consolidate, concretize, corporate, consubstantiate whatever one has done . Introduction so prepared has helped me in the continuation of the work and I humbly hope it does serve the purpose of the others, in the similar realm of progression as it has served mine. Predicational anteriorities, ontological consonance, primordial exactitude, analytical treatment of concatenated equations, Studies of alternative methodologies, possibility of wave motion, Lee paradigm and optimization problems and stochastic treatment shall be done in the subsequent papers.

Introduction

Dreams are but interludes which Fancy makes
When monarch Reason sleeps, this mimic wakes
Compounds a medley of disjointed things
A mob of cobblers, and a court of kings
Light fumes are merry, grosser fumes are sad
Both are the reasonable soul run mad
And many monstrous forms in sleep we see
That neither were, nor are, nor e'er can be
Sometimes forgotten things long cast behind
Rush forward in the brain, and come to mind.
The nurse's legends are for truths received
And the man dreams but what the boy believed
Sometimes we but rehearse a former play
The night restores our actions done by day
As hounds in sleep will open for their prey
In short, the farce of dreams is of a piece
Chimeras all; and more absurd, or less

John Dryden

“Life is but a dream for the dead.”

— Gerard Way

“Live as if you were to die tomorrow. Learn as if you were to live forever.”

Mahatma Gandhi
(Indian Philosopher, internationally esteemed for his doctrine of nonviolent protest, 1869-1948)

"[T]he atoms or elementary particles themselves are not real; they form a world of potentialities or possibilities rather than one of things or facts."

Werner Heisenberg

Anyone not shocked by quantum mechanics has not yet understood it."

Niels Bohr

In the beginning there were only probabilities. The universe could only come into existence if someone observed it. It does not matter that the observers turned up several billion years later. The universe exists because we are aware of it."

Martin Rees

We give a consolidated and consummated model for the following systems:

1. Decoherence Vis-à-vis attenuation of mesoscopic echoes in a spin-chain channel.
2. Spin ladder Vis-à-vis mesoscopic echoes
3. spin channels Vis-à-vis , quantum channels
4. quantum information Vis-à-vis quantum computation
5. Keldysh formalism Vis-à-vis Spin Dynamics
6. dynamical decoupling sequences for a qubit Vis-à-vis rapidly fluctuating spin bath
7. Dynamical decoupling Vis-à-vis quantum information
8. quantum computation Vis-à-vis time reversion
9. Spin echoes Vis-à-vis spin baths
10. Zeno and Anti-Zeno Polarization Control of Spin Ensembles Vis-à-vis Induced Dephasing
11. Zeno effect Vis-à-vis anti-Zeno effect
12. Quantum measurements Vis-à-vis quantum information
13. quantum computation Vis-à-vis spin dynamics
14. Decoherence Vis-à-vis Thermalization
15. Optimal pulse spacing for dynamical decoupling Vis-à-vis presence of a purely-dephasing spin-bath
16. Open systems Vis-à-vis quantum interference
17. Robust Dynamical Decoupling Vis-à-vis Quantum Computing and Quantum Memory
18. Quantum Relaxation **Vis-à-vis** quantum memories and decoherence
19. Localization effects Vis-à-vis decoherence in superpositions of many-spin quantum states
20. Anderson localization Vis-à-vis thermalization
21. quantum information Vis-à-vis state transfer
22. Robustness of spin-coupling distributions Vis-à-vis for perfect quantum state transfer
23. Perfect state transfer Vis-à-vis spin chains and mesoscopic echo
24. Measuring the spectrum of colored noise Vis-à-vis by dynamical decoupling
25. dynamical decoupling Vis-à-vis spin systems
26. spin echoes Vis-à-vis noise spectroscopy
27. Controlling Spin-Spin Network Dynamics Vis-à-vis Projective Measurements
28. quantum Zeno effect Vis-à-vis TOCSY
29. quantum information Vis-à-vis spin-spin couplings
30. Shift-driven modulations Vis-à-vis spin-echo signals
31. dynamical decoupling Vis-à-vis NMR
32. Hahn echo Vis-à-vis spin echo
33. Time reversion Vis-à-vis decoherence
34. Chemical shift determination Vis-à-vis field inhomogeneities
35. in-medium properties of hadrons Vis-à-vis Field Inhomogeneities of nuclear physics
36. low-energy pi pi Photoproduction Vis-à-vis Hadronic OFF NUCLEIL
37. Sub-cycle switch-on Vis-à-vis ultra strong light-matter interaction
38. light-matter interaction Vis-à-vis Microcavity
39. polaritons Vis-à-vis intersubband
40. quantum wells Vis-à-vis Hadrons
41. Signatures of the ultra strong light-mater coupling regime Vis-à-vis intersubband transitions
42. Towards Intersubband Polari tonics Vis-à-vis Light and Electrons Mate
43. Intersubband transitions Vis-à-vis ultra strong light-matter coupling
44. non-adiabatic switch-on Vis-à-vis Ultrafast phenomenon in Semiconductors
45. Extreme THz nonlinearities Vis-à-vis in bulk and nanostructured semiconductors
46. electrophonic sounds Vis-à-vis from Leonid fireballs
47. Time resolved diagnostics of plasmas Vis-à-vis ablative capillary discharges
48. Meteorites Vis-à-vis electrophonic sounds
49. Numerical modeling of laser beam-target interaction Vis-à-vis the plasma formation threshold

50. Lasing in an ablative capillary discharge Vis-à-vis structured return conductor.
51. Rogowski coil Vis-à-vis fast pulsed-current measurements of capillary discharge
52. Drag on particles in a nematic suspension Vis-à-vis moving nematic-isotropic interface
53. Defect structures and torque on an elongated colloidal particle immersed Vis-à-vis a liquid crystal host with topological defects
54. Entropic torque. Vis aVis depletion effects
55. confined liquid Vis-à-vis density functional,
56. Boundary slip Vis-à-vis of a prewetting transition
57. Capillary bridging Vis-à-vis long-range attractive forces in a mean-field approach
58. Vanishing Perturbative Vacuum Energy Vis-à-vis non-Supersymmetric Orientifolds
59. Axially Symmetric Gas Dynamic Trap and Magnetic confinement
60. Spatial profiles of fusion product flux Vis-à-vis the gas dynamic trap with deuterium neutral beam injection
61. Fusion regimes of axisymmetric open system Vis-à-vis powerful injection of fast particles
62. Enhancement of order fluctuations Vis-à-vis nematic liquid crystal by sonication (nematic, acoustic)
63. Low-frequency NMR relaxometry Vis-à-vis spatially constrained liquid crystals
64. Apparent low-field dispersion in the smectic Vis-à-vis A mesophase of thermotropic cyanobiphenyls
65. Proton field-cycling nuclear magnetic resonance relaxometry Vis-à-vis the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication.
66. Proton spin-lattice relaxation in a liquid crystalVis-à-vis Aerosol complex above the bulk isotropization temperature
67. Field-cycling NMR relaxometry Vis-à-vis rotating-frame spin-lattice relaxation
68. Aerosol Vis-à-vis order director fluctuations, Monte Carlo
69. Field-cycling NMR relaxometry Vis-à-vis spin-lattice relaxation
70. Rouse model Vis-à-vis molecular dynamics.
71. The "Corset Effect" of Spin-Lattice Relaxation in Polymer Melts Vis-à-vis Nanoporous Media
72. Acoustically stimulated NMR relaxometry Vis-à-vis ultrasound order fluctuations
73. magnetoacoustically manipulated nematic ordered states Vis-à-vis Quantum memory effects
74. Ebert: k-string tensions Vis-à-vis the 4-d SU(N)-inspired dual Abelian-Higgs-type theory
75. . Nonperturbative effects Vis-à-vis Quantum confinement and lattice gauge field theories
76. 3D SU(N) Georgi-Glashow model Vis-à-vis magnetic monopoles confining strings
77. quark confinement Vis-à-vis string and field theories at finite temperature
78. Wilson and Polyakov loops Vis-à-vis string effective action,
79. String breaking in QCD(dual superconductor)**Vis-à-vis** stochastic vacuum model
80. Confinement in the Abelian-Higgs-type theories **Vis-à-vis** String picture and field correlators
81. Field correlators **Vis-à-vis** string representation
82. Quark confinement **Vis-à-vis** stochastic vacuum model of QCD
83. dual Abelian-Higgs-type theories **Vis-à-vis** Aharonov-Bohm effect
84. Luescher term in the quark-antiquark potential Vis-à-vis Nonperturbative effects
85. Mixed heavy-quark-gluon(Nonperturbative effects) Bose Einstein condensate **Vis-à-vis** in the stochastic vacuum model
86. gauge-invariant two-point correlation function Vis-à-vis of gluonic field strengths
87. Heavy-quark condensate at zero and finite temperatures Vis-à-vis Quantum chromodynamics
88. heavy-quark condensate Vis-à-vis world-line formalism
89. Abelian-type theories with confinement Vis-à-vis finite-temperature effects in gauge theories
90. Photostimulated luminescence Vis-à-vis Thermal neutron imaging
91. Determination of the density of defect states Vis-à-vis thermally stimulated conductivity studied from numerical simulations
92. CP-odd A0 production at e+e-colliders Vis-à-vis the minimal supersymmetric standard model with SP violating phases.
93. Fourth generation CP phase Vis-à-vis b s transitions.
94. Radiative corrections to scalar fermion pair production Vis-à-vis high energy e+ e- collisions
95. Radiative corrections Vis-à-vis Supersymmetry
96. Higgs decays in the two Higgs doublet model: Large quantum effects Vis-à-vis the decoupling regime
97. d-wave to p-wave pairing Vis-à-vis the t'-t Hubbard model at zero temperature
98. Dynamical Cluster Approximation Projective Quantum Monte Carlo method Vis-à-vis Hubbard model triplet superconductivity

99. Orbital-selective Mott-Hubbard transition Vis-à-vis the two-band Hubbard model.
100. Orbital selective Mott transition Dynamical means field theory Vis-à-vis Projective Quantum Monte Carlo method Hubbard model
101. Phonon extinction rules **Vis-à-vis** thermal diffuse scattering experiments.
102. Transport theory **Vis-à-vis** Quantum Tunneling.
103. Heisenberg model **Vis-à-vis** exchange interactions (electron);
104. Ground states **Vis-à-vis** spin fluctuations.
105. Mesoscopic systems **Vis-à-vis** Green's function methods and band structure.
106. U-J synergy effect **Vis-à-vis** high Tc superconductors.
107. High Tc superconductivity Vis-à-vis - Extended Hubbard model
108. Fluctuation-dissipation relation of a dissipative driven mesoscopic system Vis-à-vis Time dependent Transport
109. Quantum Monte Carlo method Vis-à-vis molecular nanodevices
110. Anderson-Holstein model Vis-à-vis molecular transistor.
111. Green-function approach Vis-à-vis transport phenomena in quantum pumps.
112. Time-dependent quantum transport Vis-à-vis Keldysh formalism
113. Short optical pulses, Vis-à-vis Autocorrelation
114. FEL Efficiency Increase Vis-à-vis Magnetic Field Tapering of the Undulators
115. Cerenkov Free Electron Lasers and tapering and treatment of saturation
116. Free electron laser oscillators with tapered undulators Vis-à-vis Inclusion of harmonic generation and pulse propagation
117. On the nonlinear stability Vis-à-vis asymptotically anti-de Sitter solutions
118. Stability of geons, boson stars and black holes (asymptotically nonlinear solutions which are nonlinearly stable Vis-à-vis Black hole binaries.
119. Wilhelm Weber and the Surface Charges Vis-à-vis Resistive Conductors Carrying Steady Currents surface
120. Andre K.T. Assis: The Experimental and Historical Foundations of Electricity Montreal: Apeiron, 2010, 268 pp. [Physics, electricity, Stephen Gray, electrostatics
121. Electron-beam propagation in a two-dimensional electron gas.
122. Quantum point contact, ballistic transport, potential fluctuations, and two-dimensional electron gas
123. Quasiballistic transport in HgTe quantum-well nanostructures
124. Isotopic dilution, self-association, and Raman non-coincidence in the binary system
125. The black hole and Global topological k-defects
126. Chiral Symmetry Breaking and Pions in Nonsupersymmetric Gauge / Gravity Duals
127. Space-time dependent couplings **in** n=1 SUSY gauge theories: anomalies and central functions.
128. Holographic anomalies and higher derivative gravity
129. Two-hadron semi-inclusive production including subleading twist
130. New observables in longitudinal single-spin asymmetries in semi-inclusive DIS
131. Single-spin asymmetries in semi-inclusive deep-inelastic scattering
132. Single-spin asymmetries: and Collins fragmentation function
133. Di hadron interference fragmentation functions in proton-proton collisions.
134. non-centrosymmetric and penetration depth,
135. Non-centrosymmetric superconductor and penetration depth,
136. Super neutronic, Superheavy of Anti Matter
137. Bose-Einstein condensation and neutron star core.
138. Distribution of the Relaxation Times from Dielectric Spectra
139. Phase transition and Ionic conductivity
140. Acoustoelectric effects in Sn2P2S6-type ferroelectric semiconductors
141. Superconductivity and Neutrons
142. magnetic ordering on dynamical conductivity
143. Transport Properties and Magnetic properties
144. Discriminating between superconducting fluctuations and pseudogap effects
145. Pseudogap and Superconductivity
146. Vortex dynamics and Phase diagram
147. Mechanical and electronic properties of vanadium oxide nanotubes
148. Crystal-to-amorphous state transitions at normal and high pressure in the systems with semiconductor phases
149. Scaling in a temperature quench in systems with a Lifshitz point with nonconserved and conserved order parameters.
150. Dynamo mechanism: Effects of correlations and viscosities

151. Magnetohydrodynamic turbulence in three dimensions and classes of coupled driven diffusive systems.
152. Angle resolved photoemission and Surface electronic phenomena
153. work function and surface potential
154. Fermi Gap Stabilization of an Incommensurate Two-Dimensional (Surface electronic structure) Superstructure
155. Shockley state and Surface reconstruction
156. Electronic structure of organic thin films on metal surfaces
157. The surface reconstruction induced band-gap in the Shockley state on monolayer systems on noble metals.
158. surface modifications by ARUPS on Shockley-type surface states
159. Evolution of a symmetry gap and synergetic quantum well states in ultrathin Ag films
160. growth and self organization of nanostructures
161. Angle resolved photoemission and Surface electronic phenomena
162. The Shockley-type surface state on Solid-gas interfaces
163. Effect of rare-gas adsorption on the spin-orbit split bands of a surface alloy
164. Role of intermolecular interactions **on** the electronic and geometric structure of a large pi-conjugated molecule adsorbed on a metal surface
165. Adsorption energy and geometry of physisorbed organic molecules
166. the atomic contributionVis-a-Vis the Rashba spin-orbit splitting in surface alloys
167. spin-orbit interactions and two-dimensional electron gas
168. White Noise Analysis and the Feynman Path Integral
169. Functional Integrals in Stochastic and Quantum Dynamics
170. White Noise Path Integrals with Dirichlet, Neumann, and Periodic Boundary Conditions
171. Reactivity of water-electron complexes on crystalline ice surfaces
172. spin-1/2 XXZ chain and the Bose gas
173. Supramolecular Assemblies and Nano dots
174. Highly ordered nanoporous thin films from block copolymer supramolecular assembly
175. Arrays of Inorganic Nanodots and Nanowires Using Nanotemplates Based on Switchable Block Copolymer Supramolecular Assemblies.
176. Highly ordered palladium Nanodots and nanowires from switchable block copolymer thin films
177. Hexagonally ordered arrays of metallic Nanodots from thin films of functional block copolymers
178. Supramolecular Assembly of Poly (styrene)-b-poly (4-vinylpyridine) and 1-Pyrenebutyric Acid in Thin Film and Their Use for Nanofabrication
179. Solvent Vapour Induced Orientation Switching in PS-b-P4VP Block Copolymer Thin Films.
180. Nanostructures based on self-assembly of block copolymers
181. Effects of post-growth annealing on physical properties of SrRuO3 thin film grown by MOCVD
182. Optical constants, and Surface morphology
183. X-ray diffraction and Metal-organic chemical vapor deposition.
184. role of built-in strain and extra phases on Electrical resistivity
185. Quantum order in the chiral magnet MnSi
186. Topological Hall effect in the A-phase of MnSi
187. Metastable superfluidity of repulsive fermionic atoms in optical lattices
188. Skyrmion lattice in a chiral magnet
189. Chirality induced anomalous-Hall effect in helical spin crystals.
190. Polymers in crowded environment on a random walk under stretching force: globule-coil transitions
191. Walking on fractals: diffusion and self-avoiding walks on percolation clusters
192. Multifractality of self-avoiding walks on percolation clusters
193. Scaling behavior of self-avoiding walks on percolation clusters
194. Effect of chromatic mechanisms **on the** detection of mesopic incremental targets at different eccentricities
195. Determination of alpha_s **from** tau decays.
196. Kondo screening cloud **in a** one dimensional wire: Numerical renormalization group study
197. Friedel oscillations and the Kondo screening cloud: A Quantum impurity problem
198. The applicability of bosonization and the Anderson-Yuval **methods at the** strong coupling limit of quantum impurity problems.
199. Toward a microscopic description of Kondo dimers **on** metallic surfaces
200. Kondo cloud and spin-spin correlations **around a** partially screened magnetic impurity
201. Feynman path integrals **for** quantum particles propagators interacting with environments
202. Some studies on functional integrals representations **for** fluid motion with random conditions.

203. The Static Confining Potential for QCD **in the** Mandelstam Model
204. Steady state and time-resolved photoluminescence characterization **of c**opper gallium disulphide
205. BCS-BEC crossover in dense relativistic matter: Collective excitations
206. Nambu-Jona-Lasinio model and Spontaneous symmetry breaking
207. Type-II Goldstone bosons and Bose-Einstein condensation
208. Strongly interacting Fermi systems in 1/N expansion:
209. Cold atoms to color superconductivity
210. Helical ordering in the ground state of spin-one color superconductors **as a consequence** of parity violation.
211. Producing high fidelity single photons with optimal **brightness via** wave guided parametric down-conversion
212. Entanglement **and** optical parametric oscillator
213. A Quantum Walk with **Adjustable** Coin Operations
214. Robustness of bipartite Gaussian entangled beams and **propagating** in lossy channels
215. Hong-Ou-Mandel interference **and** parametric down-conversion
216. Negative Wigner Function of a Pulsed Single Photon Point by Point **and** parametric down conversion
217. Multimode squeezing **with** correlation functions.
218. Multimode quantum state **and** Parametric down-conversion
219. Schmidt modes **and** Correlation functions
220. spectral filters **as(=)** Gaussian quantum optical channels
221. Spectral Filtering **AND ,** Schmidt Number

Section 1:

Decoherence Vis-à-vis Attenuation Of Mesoscopic Echoes In A Spin-Chain Channel, Spin Ladder Vis-à-vis Mesoscopic Echoes Et Al, A Thunderstorm Thermistor-Verkrampte Reactionary Model

1. Decoherence Vis-à-vis attenuation of mesoscopic echoes in a spin-chain channel.
2. Spin ladder Vis-à-vis mesoscopic echoes
3. spin channels Vis-à-vis, quantum channels
4. quantum information Vis-à-vis quantum computation
5. Keldysh formalism Vis-à-vis Spin Dynamics
6. dynamical decoupling sequences for a qubit Vis-à-vis rapidly fluctuating spin bath
7. Dynamical decoupling Vis-à-vis quantum information
8. quantum computation Vis-à-vis time reversion
9. Spin echoes Vis-à-vis spin baths

NOTATION

Module One

Decoherence **Vis a Vis** attenuation of mesoscopic echoes in a spin-chain channel

G_{13} : Category one of Decoherence

G_{14} : Category two of Decoherence

G_{15} : Category three of Decoherence

T_{13} : Category one of attenuation of mesoscopic echoes in a spin-chain channel

T_{14} : Category two of attenuation of mesoscopic echoes in a spin-chain channel

T_{15} : Category three of attenuation of mesoscopic echoes in a spin-chain channel

Spin ladder Vis-à-vis mesoscopic echoes

Module Two

G_{16} : Category one of Spin ladder

G_{17} : Category two of Spin ladder

G_{18} : Category three of Spin ladder

T_{16} : Category one of mesoscopic echoes

T_{17} : Category two of mesoscopic echoes

T_{18} : Category three of mesoscopic echoes

Module three

spin channels Vis-à-vis, quantum channels

G_{20} : Category one of spin channels

G_{21} : Category two of spin channels

G_{22} : Category three of spin channels

T_{20} : Category one of quantum channels

T_{21} : Category two of quantum channels

T_{22} : Category three of quantum channels

Module four

quantum information Vis-à-vis quantum computation

G_{24} : Category one of quantum information

G_{25} : Category two of quantum information

G_{26} : Category three of quantum information

T_{24} : Category one of quantum computation

T_{25} : Category two of quantum computation

T_{26} : Category three of quantum computation

Module five

Keldysh formalism Vis-à-vis Spin Dynamics

G_{28} : Category one of Keldysh formalism

G_{29} : Category two of Keldysh formalism

G_{30} : Category three of Keldysh formalism

T_{28} : Category one of Spin Dynamics

T_{29} : Category two of Spin Dynamics

T_{30} : Category three of Spin Dynamics

Module six

dynamical decoupling sequences for a qubit Vis-à-vis rapidly fluctuating spin bath

G_{32} : Category one of dynamical decoupling sequences for a qubit

G_{33} : Category two of dynamical decoupling sequences for a qubit

G_{34} : Category three of dynamical decoupling sequences for a qubit

T_{32} : Category one of rapidly fluctuating spin bath

T_{33} : Category two of rapidly fluctuating spin bath

T_{34} : Category three of rapidly fluctuating spin bath

Module seven

Dynamical decoupling Vis-à-vis quantum information

G_{36} : Category one of Dynamical decoupling

G_{37} : Category two of Dynamical decoupling

G_{38} : Category three of Dynamical decoupling

T_{36} : Category one of quantum information

T_{37} : Category two of quantum information

T_{38} : Category three of quantum information

Module eight

quantum computation Vis-à-vis time reversion

G_{40} : Category one of quantum computation

G_{41} : Category two of quantum computation

G_{42} : Category three of quantum computation

T_{40} : Category one of time reversion

T_{41} : Category two of time reversion

T_{42} : Category three of time reversion

Module Nine

Spin echoes Vis-à-vis spin baths

G_{44} : Category one of Spin echoes

G_{45} : Category two of Spin echoes

G_{46} : Category three of Spin echoes

T_{44} : Category one of spin baths

T_{45} : Category two of spin baths

T_{46} : Category three of spin baths

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}$ $(a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$: $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}$,
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$,
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$, $(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$, $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$,
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$,
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$,

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right] & \left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right] & \left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right] & \left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for

category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1,2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1,2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1,2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1,2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1,2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1,2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1,2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1,2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1,2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* $1,2$ *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation

coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3
$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3
$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3
$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3
$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3
$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3
$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3
$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \quad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37}$$

$$- \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36}$$

$$- \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)} G_{37}$$

$$- \left[\begin{array}{ccc} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second

detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3
$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth

augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth

detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$

$$= (a_{45})^{(9)}G_{44}$$

$$-\begin{bmatrix} (a'_{45})^{(9)}+(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt}$$

$$= (a_{46})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{44}}{dt} =$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)}\ \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\left[\begin{array}{c} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)}\ \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \quad (98)$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: (100)

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: (101)

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \le (p_i)^{(2)} \le (\hat{A}_{16})^{(2)} \quad (102)$$

$$(b_i'')^{(2)}(G_{19},t) \le (r_i)^{(2)} \le (b_i')^{(2)} \le (\hat{B}_{16})^{(2)} \quad (103)$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad (104)$$

$$\lim_{G\to\infty}(b_i'')^{(2)}((G_{19}),t) = (r_i)^{(2)} \quad (105)$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\,(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{\mathrm{G}\to\infty}(b_i'')^{(3)}\,(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t)-(a_i'')^{(3)}(T_{21},t)|\le(\hat{k}_{20})^{(3)}|T_{21}-T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}{}',t)-(b_i'')^{(3)}(G_{23},t)|<(\hat{k}_{20})^{(3)}||G_{23}-G_{23}{}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)},(\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)}=1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}},\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)},(\hat{A}_{20})^{(3)}and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)},(a_i')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(p_i)^{(3)},\ (r_i)^{(3)},i=20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)}+(a_i')^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)}+(b_i')^{(3)}+(\hat{B}_{20})^{(3)}+(\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

Where we suppose

$$(a_i)^{(4)},(a_i')^{(4)},(a_i'')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(b_i'')^{(4)}>0,\quad i,j=24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)},(b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t)\le(p_i)^{(4)}\le(\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big)\le\ (r_i)^{(4)}\le(b_i')^{(4)}\le(\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t)=(p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big)=\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i=24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t)-(a_i'')^{(4)}(T_{25},t)|\le(\hat{k}_{24})^{(4)}|T_{25}-T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)}=1$then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the

system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)} |T_{29} - T_{29}'| e^{-(\hat{M}_{28})^{(5)} t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} ||(G_{31}) - (G_{31})'|| e^{-(\hat{M}_{28})^{(5)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)},$ are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}<1$$

Definition of $(\hat{P}_{28})^{(5)},(\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)},(\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)},(a_i')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(p_i)^{(5)},\ (r_i)^{(5)},i=28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\,(a_i)^{(5)}+(a_i')^{(5)}+\ (\hat{A}_{28})^{(5)}+\ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}]<1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)}+(b_i')^{(5)}+\ (\hat{B}_{28})^{(5)}+\ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}]<1$$

Where we suppose

$$(a_i)^{(6)},(a_i')^{(6)},(a_i'')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(b_i'')^{(6)}>0,\quad i,j=32{,}33{,}34$$ 127

The functions $(a_i'')^{(6)},(b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t)\le(p_i)^{(6)}\le(\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t)\le\ (r_i)^{(6)}\le(b_i')^{(6)}\le(\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33},t)=(p_i)^{(6)}$$ 128

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}),t\right)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-\ T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\left((G_{35}),t\right)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\widehat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$: 129

$(\widehat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\ ,\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}<1$$

Definition of $(\hat{P}_{32})^{(6)},(\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38 \qquad (131)$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \qquad (132)$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty}(a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty}(b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (a_i)^{(9)} + (a_i')^{(9)} + \ (\hat{A}_{44})^{(9)} + \ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + \ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{\left(\hat{M}_{20}\right)^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{\left(\hat{M}_{20}\right)^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \;,\; \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \;,\; \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \;,\; \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \;,\; \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^{0} + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^{0} + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^{0} + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:
Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy
$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)} + G_{29}^0) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)} + G_{33}^0) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)}\left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)}\left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t)\leq G_{44}^{0}+\int_{0}^{t}\left[(a_{44})^{(9)}\left(G_{45}^{0}+(\hat{P}_{44})^{(9)}e^{(\hat{M}_{44})^{(9)}s_{(44)}}\right)\right]ds_{(44)}=$$

$$\left(1+(a_{44})^{(9)}t\right)G_{45}^{0}+\frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)}t}-1\right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)},T^{(1)}),(G^{(2)},T^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{13})^{(1)}|G_{14}^{(1)}-G_{14}^{(2)}|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}ds_{(13)}+$$

$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$

$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{20})^{(3)}|G_{21}^{(1)}-G_{21}^{(2)}|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}|G_{20}^{(1)}-G_{20}^{(2)}|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}(T_{21}^{(1)},s_{(20)})|G_{20}^{(1)}-G_{20}^{(2)}|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}|(a''_{20})^{(3)}(T_{21}^{(1)},s_{(20)})-(a''_{20})^{(3)}(T_{21}^{(2)},s_{(20)})|\;e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$|G_{23}{}^{(1)}-G_{23}{}^{(2)}|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 216

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})-(a''_{24})^{(4)}(T_{25}^{(2)},s_{(24)})|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)})d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $((\widehat{M}_{24})^{(4)})_1,((\widehat{M}_{24})^{(4)})_2$ *and* $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 239

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right)$, $t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)})\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32{,}33{,}34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})}$: $\left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$
$G_{36}^{(2)}\left|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a_{36}')^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right) \quad (259)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a_{36}'')^{(7)}$ and $(b_{36}'')^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t} \; and \; (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (261)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: (262)

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37} \; or \; G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36} \; is$ bounded, from below, the same property holds for $G_{37} \; and \; G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. (263)

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. (264)

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t), t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can 274

postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t),t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t\ > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t),t(t),t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose
$(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)}+\left((\hat{P}_{44})^{(9)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\;e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\,(G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(9)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a_{13}')^{(1)} + (a_{14}')^{(1)} - (a_{13}'')^{(1)}(T_{14}, t) + (a_{14}'')^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b_{13}')^{(1)} + (b_{14}')^{(1)} - (b_{13}'')^{(1)}(G, t) - (b_{14}'')^{(1)}(G, t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$

$\left(\frac{(a_{15})^{(1)}G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le\right.$ 286

$\left.\frac{(a_{15})^{(1)}G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$

$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$ 287

$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$ 288

$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le$ 289

$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16}\mathrm{e}^{\left((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)}\right)\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$
$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)},\ \boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)},\ \boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)},$ and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big) - (b''_{25})^{(4)}\big((G_{27}),t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)},, (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, $\boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24}e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G^0_{24}e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)}G^0_{24}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G^0_{26}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T^0_{24}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T^0_{26}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)}=(u_1)^{(5)},(\mu_1)^{(5)}=(\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)}<(u_0)^{(5)}<(\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right) \quad 366$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)},, (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^{0}}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^{0}e^{-(S_2)^{(9)}t}\leq G_{46}(t)\leq \frac{(a_{46})^{(9)}G_{44}^{0}}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}\left[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}\right]+G_{46}^{0}e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)\leq\frac{d\nu^{(2)}}{dt}\leq-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)},(\nu_0)^{(2)}$:-

For $0<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\nu_1)^{(2)}<(\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t)\geq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}\quad,\quad\boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\bar{\nu}_1)^{(2)}$ 392

If $0<(\nu_1)^{(2)}<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)}\leq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}\leq\nu^{(2)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\bar{\nu}_1)^{(2)}$$

If $0<(\nu_1)^{(2)}\leq(\bar{\nu}_1)^{(2)}\leq(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)}\leq\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)}\leq\nu^{(2)}(t)\leq(m_1)^{(2)},\quad\boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)}\leq u^{(2)}(t)\leq(\mu_1)^{(2)},\quad\boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)}=(a_{17}'')^{(2)},then\ (\sigma_1)^{(2)}=(\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)}=(\bar{\nu}_1)^{(2)}$ if in addition

$(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\quad \boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(v_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get 409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{v}_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right) - (a''_{33})^{(6)}(T_{33},t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \le \frac{d\nu^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\quad \boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad (483)$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad (484)$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad (484A)$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: (485)

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: (486)

(a) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: (487)

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: (488)

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: (489)

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T_{25}^{*} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^{*} for which $f(T_{25}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)} G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)} G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

Definition and uniqueness of T_{29}^{*} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^{*} for which $f(T_{29}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)} G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)} G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - \left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \qquad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \tag{545}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \tag{546}$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\big((b_{29}')^{(5)} - (r_{29})^{(5)}\big)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\big(s_{(29)(j)}T_{29}^*\mathbb{G}_j\big) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}\ and\ (b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \ , \ \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\ (G_{35})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}\ and\ (b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left(\ (G_{39})^{**}\ \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\Big(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\Big(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\Big(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\Big(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\Big(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\Big(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 2:

Zeno and Anti-Zeno Polarization Control of Spin Ensembles Vis a Vis Induced Dephasing, Zeno effect Vis a Vis anti-Zeno effect, Quantum measurements Vis a Vis quantum information: A Testamentary Dictyoptera-Synecdochal Syncretism Model

1. Zeno and Anti-Zeno Polarization Control of Spin Ensembles Vis-à-vis Induced Dephasing
2. Zeno effect Vis-à-vis anti-Zeno effect
3. Quantum measurements Vis-à-vis quantum information
4. quantum computation Vis-à-vis spin dynamics
5. Decoherence Vis-à-vis Thermalization
6. Optimal pulse spacing for dynamical decoupling Vis-à-vis presence of a purely-dephasing spin-bath
7. Open systems Vis-à-vis quantum interference
8. Robust Dynamical Decoupling Vis-à-vis Quantum Computing and Quantum Memory
9. Quantum Relaxation Vis-à-vis quantum memories and decoherence

NOTATION

Module One

Zeno and Anti-Zeno Polarization Control of Spin Ensembles Vis-à-vis Induced Dephasing

G_{13} : Category one of Zeno and Anti-Zeno Polarization Control of Spin Ensembles

G_{14} : Category two of Zeno and Anti-Zeno Polarization Control of Spin Ensembles

G_{15} : Category three of Zeno and Anti-Zeno Polarization Control of Spin Ensembles

T_{13} : Category one of Induced Dephasing

T_{14} : Category two of Induced Dephasing

T_{15} : Category three of Induced Dephasing

Module Two

Zeno effect Vis-à-vis anti-Zeno effect

G_{16} : Category one of Zeno effect

G_{17} : Category two of Zeno effect

G_{18} : Category three of Zeno effect

T_{16} : Category one of anti-Zeno effect

T_{17} : Category two of anti-Zeno effect

T_{18} : Category three of anti-Zeno effect

Module three

Quantum measurements Vis-à-vis quantum information

G_{20} : Category one of Quantum measurements

G_{21} : Category two of Quantum measurements

G_{22} : Category three of Quantum measurements

T_{20} : Category one of quantum information

T_{21} : Category two of quantum information

T_{22} : Category three of quantum information

Module four

quantum computation Vis-à-vis spin dynamics

G_{24} : Category one of quantum computation

G_{25} : Category two of quantum computation

G_{26} : Category three of quantum computation

T_{24} : Category one of spin dynamics

T_{25} : Category two of spin dynamics

T_{26} : Category three of spin dynamics

Module five

Decoherence Vis-à-vis Thermalization

G_{28} : Category one of Decoherence

G_{29} : Category two of Decoherence

G_{30} : Category three of Decoherence

T_{28} : Category one of Thermalization

T_{29} : Category two of Thermalization

T_{30} : Category three of Thermalization

Module six

Optimal pulse spacing for dynamical decoupling Vis-à-vis presence of a purely-dephasing spin-bath

G_{32} : Category one of Optimal pulse spacing for dynamical decoupling

G_{33} : Category two of Optimal pulse spacing for dynamical decoupling

G_{34} : Category three of Optimal pulse spacing for dynamical decoupling

T_{32} : Category one of presence of a purely-dephasing spin-bath

T_{33} : Category two of presence of a purely-dephasing spin-bath

T_{34} : Category three of presence of a purely-dephasing spin-bath

Open systems Vis-à-vis quantum interference

Module seven

G_{36} : Category one of Open systems

G_{37} : Category two of Open systems

G_{38} : Category three of Open systems

T_{36} : Category one of quantum interference

T_{37} : Category two of quantum interference

T_{38} : Category three of quantum interference

Module eight

Robust Dynamical Decoupling Vis-à-vis Quantum Computing and Quantum Memory

G_{40} : Category one of Robust Dynamical Decoupling

G_{41} : Category two of Robust Dynamical Decoupling

G_{42} : Category three of Robust Dynamical Decoupling

T_{40} : Category one of Quantum Computing and Quantum Memory

T_{41} : Category two of Quantum Computing and Quantum Memory

T_{42} : Category three of Quantum Computing and Quantum Memory

Module Nine

Quantum Relaxation Vis-à-vis quantum memories and decoherence

G_{44} : Category one of Quantum Relaxation

G_{45} : Category two of Quantum Relaxation

G_{46} : Category three of Quantum Relaxation

T_{44} : Category one of quantum memories and decoherence

T_{45} : Category two of quantum memories and decoherence

T_{46} : Category three of quantum memories and decoherence

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \qquad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients

for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3
$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b_{20}')^{(3)} & -(b_{20}'')^{(3)}(G_{23},t) & -(b_{16}'')^{(2,2,2)}(G_{19},t) & -(b_{13}'')^{(1,1,1,)}(G,t) \\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b_{21}')^{(3)} & -(b_{21}'')^{(3)}(G_{23},t) & -(b_{17}'')^{(2,2,2)}(G_{19},t) & -(b_{14}'')^{(1,1,1,)}(G,t) \\ -(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{37}'')^{(7,7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b_{22}')^{(3)} & -(b_{22}'')^{(3)}(G_{23},t) & -(b_{18}'')^{(2,2,2)}(G_{19},t) & -(b_{15}'')^{(1,1,1,)}(G,t) \\ -(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{38}'')^{(7,7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b_{20}'')^{(3)}(G_{23},t)$, $-(b_{21}'')^{(3)}(G_{23},t)$, $-(b_{22}'')^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b_{16}'')^{(2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b_{13}'')^{(1,1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition

coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition

coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad (89)$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad (90)$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad (91)$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad (92)$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)} G_{37}$$

$$- \left[\begin{array}{ccc} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second

detrition coefficients for category 1, 2 and 3
$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3
$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$
$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$
$$= (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth

augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth

detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$
$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3
$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{44}}{dt} =$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)}\ \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\left[\begin{array}{c} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)}\ \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \quad (98)$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: (100)

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: (101)

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \le (p_i)^{(2)} \le (\hat{A}_{16})^{(2)} \quad (102)$$

$$(b_i'')^{(2)}(G_{19},t) \le (r_i)^{(2)} \le (b_i')^{(2)} \le (\hat{B}_{16})^{(2)} \quad (103)$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad (104)$$

$$\lim_{G\to\infty}(b_i'')^{(2)}((G_{19}),t) = (r_i)^{(2)} \quad (105)$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}\big((G_{19}), t\big)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\,(a_i)^{(2)} + (a_i')^{(2)} + \ (\hat{A}_{16})^{(2)} + \ (\hat{P}_{16})^{(2)}\,(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ \ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\,(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{\mathrm{G}\to\infty}(b_i'')^{(3)}\,(G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t)-(a_i'')^{(3)}(T_{21},t)|\le(\hat{k}_{20})^{(3)}|T_{21}-T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}{}',t)-(b_i'')^{(3)}(G_{23},t)|<(\hat{k}_{20})^{(3)}||G_{23}-G_{23}{}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)},(\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)}=1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}},\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)},(\hat{A}_{20})^{(3)}and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)},(a_i')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(p_i)^{(3)},\ (r_i)^{(3)},i=20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)}+(a_i')^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)}+(b_i')^{(3)}+(\hat{B}_{20})^{(3)}+(\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

Where we suppose

$$(a_i)^{(4)},(a_i')^{(4)},(a_i'')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(b_i'')^{(4)}>0,\quad i,j=24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)},(b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t)\le(p_i)^{(4)}\le(\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big)\le\ (r_i)^{(4)}\le(b_i')^{(4)}\le(\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t)=(p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big)=\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i=24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t)-(a_i'')^{(4)}(T_{25},t)|\le(\hat{k}_{24})^{(4)}|T_{25}-T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)}=1$then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the

system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\widehat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\widehat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\widehat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\widehat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38 \qquad (131)$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)} \qquad (132)$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty}(a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty}(b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \le \ (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \le (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}\big((G_{47}), t\big)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (a_i)^{(9)} + (a_i')^{(9)} + \ (\hat{A}_{44})^{(9)} + \ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + \ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:
Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy
$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)} + G_{33}^0\right) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)}\left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)}\left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t)\leq G_{44}^{0}+\int_{0}^{t}\left[(a_{44})^{(9)}\left(G_{45}^{0}+(\hat{P}_{44})^{(9)}e^{(\hat{M}_{44})^{(9)}s_{(44)}}\right)\right]ds_{(44)}=$$

$$\left(1+(a_{44})^{(9)}t\right)G_{45}^{0}+\frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)}t}-1\right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)},T^{(1)}),(G^{(2)},T^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{13})^{(1)}|G_{14}^{(1)}-G_{14}^{(2)}|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}ds_{(13)}+$$

$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$

$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\bar{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\bar{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\bar{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\hat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}$ *and* $(\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 216

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})-(a''_{24})^{(4)}(T_{25}^{(2)},s_{(24)})|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)})d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $((\widehat{M}_{24})^{(4)})_1,((\widehat{M}_{24})^{(4)})_2$ *and* $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right)$, $t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)})\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32{,}33{,}34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})}$: $\left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$
$G_{36}^{(2)}\left|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad (259)$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a_{36}')^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a_{36}'')^{(7)}$ and $(b_{36}'')^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (261)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: (262)

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. (263)

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. (264)

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t), t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can 274

postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t),t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t\ > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t),t(t),t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose
$(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)}+\left((\hat{P}_{44})^{(9)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\;e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\,(G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(9)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a_{13}')^{(1)} + (a_{14}')^{(1)} - (a_{13}'')^{(1)}(T_{14}, t) + (a_{14}'')^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b_{13}')^{(1)} + (b_{14}')^{(1)} - (b_{13}'')^{(1)}(G, t) - (b_{14}'')^{(1)}(G, t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$

$\left(\frac{(a_{15})^{(1)}G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le\right.$ 286

$\left.\frac{(a_{15})^{(1)}G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$

$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$ 287

$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$ 288

$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le$ 289

$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16}\mathrm{e}^{\left((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)}\right)\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$
$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)},\ \boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)},\ \boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)},$ and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big) - (b''_{25})^{(4)}\big((G_{27}),t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)},, (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, $\boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$ 331

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24}e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G^0_{24}e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G^0_{24}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G^0_{26}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T^0_{24}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T^0_{26}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$

Definition of $(v_1)^{(5)},(v_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},v^{(5)},u^{(5)}$: 339

By $(v_1)^{(5)}>0$, $(v_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2+(\sigma_1)^{(5)}v^{(5)}-(a_{28})^{(5)}=0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{v}_1)^{(5)},,(\bar{v}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)}>0$, $(\bar{v}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2+(\sigma_2)^{(5)}v^{(5)}-(a_{28})^{(5)}=0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(v_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(v_0)^{(5)},(m_1)^{(5)}=(v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)}<(v_1)^{(5)}$

$(m_2)^{(5)}=(v_1)^{(5)},(m_1)^{(5)}=(\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)}<(v_0)^{(5)}<(\bar{v}_1)^{(5)}$,

and $\boxed{(v_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(v_1)^{(5)},(m_1)^{(5)}=(v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)}<(v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)}=(u_1)^{(5)},(\mu_1)^{(5)}=(\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)}<(u_0)^{(5)}<(\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$
and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$
and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right) \quad 366$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)},, (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^{0}}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^{0}e^{-(S_2)^{(9)}t}\leq G_{46}(t)\leq \frac{(a_{46})^{(9)}G_{44}^{0}}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}\left[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}\right]+G_{46}^{0}e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)\leq\frac{d\nu^{(2)}}{dt}\leq-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)},(\nu_0)^{(2)}$:-

For $0<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\nu_1)^{(2)}<(\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t)\geq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}\quad,\quad\boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\bar{\nu}_1)^{(2)}$ 392

If $0<(\nu_1)^{(2)}<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)}\leq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}\leq\nu^{(2)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\bar{\nu}_1)^{(2)}$$

If $0<(\nu_1)^{(2)}\leq(\bar{\nu}_1)^{(2)}\leq(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)}\leq\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)}\leq\nu^{(2)}(t)\leq(m_1)^{(2)},\quad\boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)}\leq u^{(2)}(t)\leq(\mu_1)^{(2)},\quad\boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)}=(a_{17}'')^{(2)},then\ (\sigma_1)^{(2)}=(\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)}=(\bar{\nu}_1)^{(2)}$ if in addition

$(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(v_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get 409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{v}_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)}=(a''_{29})^{(5)}, then\ (\sigma_1)^{(5)}=(\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)}=(\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)}=(v_5)^{(5)}$ then $v^{(5)}(t)=(v_0)^{(5)}$ and as a consequence $G_{28}(t)=(v_0)^{(5)}G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)}=(b''_{29})^{(5)}, then\ (\tau_1)^{(5)}=(\tau_2)^{(5)}$ and then $(u_1)^{(5)}=(\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)}=(u_1)^{(5)}$ then $T_{28}(t)=(u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt}=(a_{32})^{(6)}-\left((a'_{32})^{(6)}-(a'_{33})^{(6)}+(a''_{32})^{(6)}(T_{33},t)\right)-(a''_{33})^{(6)}(T_{33},t)v^{(6)}-(a_{33})^{(6)}v^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \le \frac{d\nu^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $v^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t) v^{(8)} - (a_{41})^{(8)} v^{(8)}$$

Definition of $v^{(8)}$:- $\quad \boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)} v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)} v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq v^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(b) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T_{25}^{*} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^{*} for which $f(T_{25}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)} G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)} G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

Definition and uniqueness of T_{29}^{*} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^{*} for which $f(T_{29}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)} G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)} G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - \left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \qquad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \tag{545}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \tag{546}$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\big((b_{29}')^{(5)} - (r_{29})^{(5)}\big)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\big(s_{(29)(j)}T_{29}^*\mathbb{G}_j\big) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}\ and\ (b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \ , \ \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\ (G_{35})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}\ and\ (b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left(\ (G_{39})^{**}\ \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section3 :

Localization effects Vis-à-vis decoherence in superpositions of many-spin quantum states, Anderson localization Vis-à-vis thermalization, quantum information Vis-à-vis state transfer: A Sustenational sussurationer-Poindexter poinsettia Model

INTRODUCTION—VARIABLES USED

1. Localization effects Vis-à-vis decoherence in superpositions of many-spin quantum states
2. Anderson localization Vis-à-vis thermalization
3. quantum information Vis-à-vis state transfer
4. Robustness of spin-coupling distributions Vis-à-vis for perfect quantum state transfer
5. Perfect state transfer Vis-à-vis spin chains and mesoscopic echo
6. Measuring the spectrum of colored noise Vis-à-vis by dynamical decoupling.
7. dynamical decoupling Vis-à-vis spin systems
8. spin echoes Vis-à-vis noise spectroscopy
9. Controlling Spin-Spin Network Dynamics Vis-à-vis Projective Measurements

NOTATION

Module One

Localization effects Vis-à-vis decoherence in superpositions of many-spin quantum states

G_{13} : Category one of Localization effects

G_{14} : Category two of Localization effects

G_{15} : Category three of Localization effects

T_{13} : Category one of decoherence in superpositions of many-spin quantum states

T_{14} : Category two of decoherence in superpositions of many-spin quantum states

T_{15} : Category three of decoherence in superpositions of many-spin quantum states

Module Two

Anderson localization Vis-à-vis thermalization

G_{16} : Category one of Anderson localization

G_{17} : Category two of Anderson localization

G_{18} : Category three of Anderson localization

T_{16} : Category one of thermalization

T_{17} : Category two of thermalization

T_{18} : Category three of thermalization

Module three

quantum information Vis-à-vis state transfer

G_{20} : Category one of quantum information

G_{21} : Category two of quantum information

G_{22} : Category three of quantum information

T_{20} : Category one of state transfer

T_{21} : Category two of state transfer

T_{22} : Category three of state transfer

Module four

Robustness of spin-coupling distributions Vis-à-vis perfect quantum state transfer

G_{24} : Category one of Robustness of spin-coupling distributions

G_{25} : Category two of Robustness of spin-coupling distributions

G_{26} : Category three of Robustness of spin-coupling distributions

T_{24} : Category one of perfect quantum state transfer

T_{25} : Category two of perfect quantum state transfer

T_{26} : Category three of perfect quantum state transfer

Module five

Perfect state transfer Vis-à-vis spin chains and mesoscopic echo

G_{28} : Category one of Perfect state transfer

G_{29} : Category two of Perfect state transfer

G_{30} : Category three of Perfect state transfer

T_{28} : Category one of spin chains and mesoscopic echo

T_{29} : Category two of spin chains and mesoscopic echo

T_{30} : Category three of spin chains and mesoscopic echo

Module six

Measuring the spectrum of colored noise Vis-à-vis dynamical decoupling

G_{32} : Category one of Measuring the spectrum of colored noise

G_{33} : Category two of Measuring the spectrum of colored noise

G_{34} : Category three of Measuring the spectrum of colored noise

T_{32} : Category one of dynamical decoupling

T_{33} : Category two of dynamical decoupling

T_{34} : Category three of dynamical decoupling

Module seven

dynamical decoupling Vis-à-vis spin systems

G_{36} : Category one of dynamical decoupling

G_{37} : Category two of dynamical decoupling

G_{38} : Category three of dynamical decoupling

T_{36} : Category one of spin systems

T_{37} : Category two of spin systems

T_{38} : Category three of spin systems

Module eight

spin echoes Vis-à-vis noise spectroscopy

G_{40} : Category one of spin echoes

G_{41} : Category two of spin echoes

G_{42} : Category three of spin echoes

T_{40} : Category one of noise spectroscopy

T_{41} : Category two of noise spectroscopy

T_{42} : Category three of noise spectroscopy

Module Nine

Controlling Spin-Spin Network Dynamics Vis-à-vis Projective Measurements

G_{44} : Category one of Controlling Spin-Spin Network Dynamics

G_{45} : Category two of Controlling Spin-Spin Network Dynamics

G_{46} : Category three of Controlling Spin-Spin Network Dynamics

T_{44} : Category one of Projective Measurements

T_{45} : Category two of Projective Measurements

T_{46} : Category three of Projective Measurements

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients

for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3
$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b_{20}')^{(3)} & -(b_{20}'')^{(3)}(G_{23},t) & -(b_{16}'')^{(2,2,2)}(G_{19},t) & -(b_{13}'')^{(1,1,1,)}(G,t) \\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b_{21}')^{(3)} & -(b_{21}'')^{(3)}(G_{23},t) & -(b_{17}'')^{(2,2,2)}(G_{19},t) & -(b_{14}'')^{(1,1,1,)}(G,t) \\ -(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{37}'')^{(7,7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b_{22}')^{(3)} & -(b_{22}'')^{(3)}(G_{23},t) & -(b_{18}'')^{(2,2,2)}(G_{19},t) & -(b_{15}'')^{(1,1,1,)}(G,t) \\ -(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{38}'')^{(7,7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b_{20}'')^{(3)}(G_{23},t)$, $-(b_{21}'')^{(3)}(G_{23},t)$, $-(b_{22}'')^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b_{16}'')^{(2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b_{13}'')^{(1,1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b_{36}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7)}(G_{39},t)$ are seventh detrition

coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & \boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* $1, 2$ *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category $1, 2$ *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category $1, 2$ *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* $1, 2$ *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition

coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37}$$

$$- \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36}$$

$$- \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)} G_{37}$$

$$- \left[\begin{array}{ccc} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{array} \right] G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \left[\begin{array}{ccc} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[\begin{array}{ccc} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second

detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3
$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$
$$= (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a_{40}')^{(8)}\boxed{+(a_{40}'')^{(8)}(T_{41},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt}$$
$$= (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a_{41}')^{(8)}\boxed{+(a_{41}'')^{(8)}(T_{41},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a_{42}')^{(8)}\boxed{+(a_{42}'')^{(8)}(T_{41},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a_{40}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8)}(T_{41},t)}$, $\boxed{+(a_{42}'')^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3
$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3
$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth

augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth

detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{44})^{(9)} & +(a''_{44})^{(9)}(T_{45},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$

$$= (a_{45})^{(9)}G_{44}$$

$$-\begin{bmatrix} (a'_{45})^{(9)} & +(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt}$$

$$= (a_{46})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{46})^{(9)} & +(a''_{46})^{(9)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{44}}{dt} =$

$$(b_{44})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)}\ \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\left[\begin{array}{c} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)}\ \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \quad (98)$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: (100)

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: (101)

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t) \le (p_i)^{(2)} \le (\hat{A}_{16})^{(2)} \quad (102)$$

$$(b_i'')^{(2)}(G_{19},t) \le (r_i)^{(2)} \le (b_i')^{(2)} \le (\hat{B}_{16})^{(2)} \quad (103)$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t) = (p_i)^{(2)} \quad (104)$$

$$\lim_{G\to\infty}(b_i'')^{(2)}((G_{19}),t) = (r_i)^{(2)} \quad (105)$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and \ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, \ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t)-(a_i'')^{(3)}(T_{21},t)|\le(\hat{k}_{20})^{(3)}|T_{21}-T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}{}',t)-(b_i'')^{(3)}(G_{23},t)|<(\hat{k}_{20})^{(3)}||G_{23}-G_{23}{}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)},(\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)}=1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}},\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)},(\hat{A}_{20})^{(3)}and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)},(a_i')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(p_i)^{(3)},\ (r_i)^{(3)},i=20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)}+(a_i')^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)}+(b_i')^{(3)}+(\hat{B}_{20})^{(3)}+(\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

Where we suppose

$$(a_i)^{(4)},(a_i')^{(4)},(a_i'')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(b_i'')^{(4)}>0,\quad i,j=24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)},(b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t)\le(p_i)^{(4)}\le(\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big)\le\ (r_i)^{(4)}\le(b_i')^{(4)}\le(\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t)=(p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big)=\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i=24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t)-(a_i'')^{(4)}(T_{25},t)|\le(\hat{k}_{24})^{(4)}|T_{25}-T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)}=1$then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the

system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\widehat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\widehat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\widehat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ \ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\widehat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)},$ are positive constants

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \ , \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty}(a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty}(b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}\big((G_{47}), t\big)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (a_i)^{(9)} + (a_i')^{(9)} + \ (\hat{A}_{44})^{(9)} + \ (\hat{P}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)} + (b_i')^{(9)} + \ (\hat{B}_{44})^{(9)} + \ (\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{(\hat{M}_{20})^{(3)}t} \;,\quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \;,\quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \;,\quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \;,\; \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \;,\; \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \;,\; \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \;,\; \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^{0} + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^{0} + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^{0} + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:
Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy
$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)} + G_{33}^0\right) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)}\left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)}\left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t)\leq G_{44}^{0}+\int_{0}^{t}\left[(a_{44})^{(9)}\left(G_{45}^{0}+(\hat{P}_{44})^{(9)}e^{(\hat{M}_{44})^{(9)}s_{(44)}}\right)\right]ds_{(44)}=$$

$$\left(1+(a_{44})^{(9)}t\right)G_{45}^{0}+\frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)}t}-1\right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)},T^{(1)}),(G^{(2)},T^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{13})^{(1)}|G_{14}^{(1)}-G_{14}^{(2)}|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}ds_{(13)}+$$

$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$

$T_i\,(t)\geq T_i^0e^{(-(b_i')^{(1)}t)}>0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1, T_{13}\,(t)>(m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\bar{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\bar{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\bar{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\hat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}$ *and* $(\hat{Q}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 216

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})-(a''_{24})^{(4)}(T_{25}^{(2)},s_{(24)})|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)})d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 228

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $((\widehat{M}_{24})^{(4)})_1,((\widehat{M}_{24})^{(4)})_2$ *and* $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right)$, $t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)})\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32{,}33{,}34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})}$: $\left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$
$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right) \quad 259$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t} \; and \; (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If $G_{37} \; or \; G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36} \; is$ bounded, from below, the same property holds for $G_{37} \; and \; G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\,((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t), t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can 274

postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t),t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t),t(t),t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose
$(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)}+\left((\hat{P}_{44})^{(9)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\;e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\;(G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(9)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t),t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a_{13}')^{(1)} + (a_{14}')^{(1)} - (a_{13}'')^{(1)}(T_{14},t) + (a_{14}'')^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b_{13}')^{(1)} + (b_{14}')^{(1)} - (b_{13}'')^{(1)}(G,t) - (b_{14}'')^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$

$\left(\frac{(a_{15})^{(1)}G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le\right.$ 286

$\left.\frac{(a_{15})^{(1)}G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$

$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$ 287

$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$ 288

$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le$ 289

$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16}\mathrm{e}^{\left((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)}\right)\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$
$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)},\ \boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)},\ \boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)},$ and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right) \quad 324$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big) - (b''_{25})^{(4)}\big((G_{27}),t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)},, (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, $\boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$ 331

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24}e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G^0_{24}e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G^0_{24}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G^0_{26}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T^0_{24}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T^0_{26}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)}=(u_1)^{(5)},(\mu_1)^{(5)}=(\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)}<(u_0)^{(5)}<(\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$

where $(p_i)^{(5)}$ is defined by equation

$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t}$ 343

$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le\right.$ 344
(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}}$ 345

$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}$ 346

$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le$ 347

$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$
and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$
and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right) \quad 366$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, , (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^{0}}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^{0}e^{-(S_2)^{(9)}t}\leq G_{46}(t)\leq \frac{(a_{46})^{(9)}G_{44}^{0}}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}\left[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}\right]+G_{46}^{0}e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)\leq\frac{d\nu^{(2)}}{dt}\leq-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)},(\nu_0)^{(2)}$:-

For $0<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\nu_1)^{(2)}<(\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t)\geq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}\quad,\quad\boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\bar{\nu}_1)^{(2)}$ 392

If $0<(\nu_1)^{(2)}<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)}\leq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}\leq\nu^{(2)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\bar{\nu}_1)^{(2)}$$

If $0<(\nu_1)^{(2)}\leq(\bar{\nu}_1)^{(2)}\leq(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)}\leq\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)}\leq\nu^{(2)}(t)\leq(m_1)^{(2)},\quad\boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)}\leq u^{(2)}(t)\leq(\mu_1)^{(2)},\quad\boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)}=(a_{17}'')^{(2)},then\ (\sigma_1)^{(2)}=(\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)}=(\bar{\nu}_1)^{(2)}$ if in addition

$(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\quad \boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(v_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get 409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{v}_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)}=(a''_{29})^{(5)}, then\ (\sigma_1)^{(5)}=(\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)}=(\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)}=(v_5)^{(5)}$ then $v^{(5)}(t)=(v_0)^{(5)}$ and as a consequence $G_{28}(t)=(v_0)^{(5)}G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)}=(b''_{29})^{(5)}, then\ (\tau_1)^{(5)}=(\tau_2)^{(5)}$ and then $(u_1)^{(5)}=(\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)}=(u_1)^{(5)}$ then $T_{28}(t)=(u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt}=(a_{32})^{(6)}-\left((a'_{32})^{(6)}-(a'_{33})^{(6)}+(a''_{32})^{(6)}(T_{33},t)\right)-(a''_{33})^{(6)}(T_{33},t)v^{(6)}-(a_{33})^{(6)}v^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \le \frac{d\nu^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $v^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t) v^{(8)} - (a_{41})^{(8)} v^{(8)}$$

Definition of $v^{(8)}$:- $\quad \boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)} v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)} v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq v^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(c) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T_{25}^{*} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^{*} for which $f(T_{25}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)} G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)} G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

Definition and uniqueness of T_{29}^{*} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^{*} for which $f(T_{29}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)} G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)} G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - \left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \qquad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \tag{545}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \tag{546}$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\big((b_{29}')^{(5)} - (r_{29})^{(5)}\big)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\big(s_{(29)(j)}T_{29}^*\mathbb{G}_j\big) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}\ and\ (b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \ , \ \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\ (G_{35})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}\ and\ (b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left(\ (G_{39})^{**}\ \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\Big(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\Big(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\Big(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\Big(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\Big(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\Big(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 4:

Quantum Zeno Effect Vis A Vis TOCSY, Quantum Information Vis-à-vis Spin-Spin Couplings Shift-Driven Modulations Vis-à-vis Spin-Echo Signals: A Burlesque Moccasin – Harmonious Key Note Model

NOTATION

1. Controlling Spin-Spin Network Dynamics Vis-à-vis Projective Measurements
2. Quantum Zeno effect Vis-à-vis TOCSY
3. Quantum information Vis-à-vis spin-spin couplings
4. Shift-driven modulations Vis-à-vis spin-echo signals
5. Dynamical decoupling Vis-à-vis NMR
6. Hahn echo Vis-à-vis spin echo
7. Time reversion Vis-à-vis decoherence
8. Chemical shift determination Vis-à-vis field inhomogeneities
9. In-medium properties of hadrons Vis-à-vis Field Inhomogeneities of nuclear physics
10. Low-energy pi pi Photoproduction Vis-à-vis Hadronic OFF NUCLEIL

Module One

Dynamics Vis-à-vis Projective Measurements controlling spin-spin network

G_{13} : Category one of Controlling Spin-Spin Network Dynamics

G_{14} : Category two of Controlling Spin-Spin Network Dynamics

G_{15} : Category three of Controlling Spin-Spin Network Dynamics

T_{13} : Category one of Projective Measurements

T_{14} : Category two of Projective Measurements

T_{15} : Category three of Projective Measurements

Module Two

Quantum Zeno effect Vis-à-vis TOCSY

G_{16} : Category one of quantum Zeno effect

G_{17} : Category two of quantum Zeno effect

G_{18} : Category three of quantum Zeno effect

T_{16} : Category one of TOCSY

T_{17} : Category two of TOCSY

T_{18} : Category three of TOCSY

Module three

Quantum information Vis-à-vis spin-spin couplings

G_{20} : Category one of quantum information

G_{21} : Category two of quantum information

G_{22} : Category three of quantum information

T_{20} : Category one of spin-spin couplings

T_{21} : Category two of spin-spin couplings

T_{22} : Category three of spin-spin couplings

Module four

Shift-driven modulations Vis-à-vis spin-echo signals

G_{24} : Category one of Shift-driven modulations

G_{25} : Category two of Shift-driven modulations

G_{26} : Category three of Shift-driven modulations

T_{24} : Category one of spin-echo signals

T_{25} : Category two of spin-echo signals

T_{26} : Category three of spin-echo signals

Module five

Dynamical decoupling Vis-à-vis NMR

G_{28} : Category one of dynamical decoupling

G_{29} : Category two of dynamical decoupling

G_{30} : Category three of dynamical decoupling

T_{28} : Category one of NMR

T_{29} : Category two of NMR

T_{30} : Category three of NMR

Module six

Hahn echo Vis-à-vis spin echo

G_{32} : Category one of Hahn echo

G_{33} : Category two of Hahn echo

G_{34} : Category three of Hahn echo

T_{32} : Category one of spin echo

T_{33} : Category two of spin echo

T_{34} : Category three of spin echo

Module seven

Time reversion Vis-à-vis decoherence

G_{36} : Category one of Time reversion

G_{37} : Category two of Time reversion

G_{38} : Category three of Time reversion

T_{36} : Category one of decoherence

T_{37} : Category two of decoherence

T_{38} : Category three of decoherence

Module eight

Chemical shift determination Vis-à-vis field inhomogeneities

G_{40} : Category one of Chemical shift determination

G_{41} : Category two of Chemical shift determination

G_{42} : Category three of Chemical shift determination

T_{40} : Category one of field inhomogeneities (A disturbance of the field homogeneity, because of magnetic material (inside or outside the patient), technical problems or scanning at the edge of the field. When images were obtained in a progression from the center to the edge of the coil, the homogeneity of the field observed by the imaged volume, changes when the distance from the center of the volume increase. The same problem appears by scanning at a distance from the isocenter in left-right direction or too large field of view. There are different types of bad image quality, the images are noisy, distorted or the fat suppression doesn't work because of badly set shim currents. E.g. by using an IR sequence, changes in the T1 recovery rates of the tissues are involved. The inversion time at the center of the imaged volume is appropriate to suppress fat, but at the edge of the coil the same inversion time is sufficient to suppress water. Since the inversion time is not changed, the T1 recovery rates will increase)

T_{41} : Category two of field inhomogeneities (MR imaging of susceptibility-induced magnetic field inhomogeneities.: Wendt RE 3rd, Wilcott MR 3rd, Nitz W, Murphy PH, Bryan RN.: A 90 degrees-tau 1-90 degrees-tau 2-image acquisition pulse sequence allows spatial mapping of resonant frequency. This sort of sequence has previously been used for magnet shimming, and its use in chemical-shift imaging has been proposed. The authors used this sequence in magnetic resonance imaging of a phantom to demonstrate the magnetic field gradients arising from susceptibility differences within the phantom and allow those gradients to be measured. Gradients may arise near interfaces between substances that cannot support the same magnetic flux density. The pulse sequence was found to work well in lower-field-strength instruments.)

T_{42} : Category three of field inhomogeneities

Module Nine

in-medium properties of hadrons Vis-à-vis Field Inhomogeneities of nuclear physics

G_{44} : Category one of in-medium properties of hadrons

G_{45} : Category two of in-medium properties of hadrons

G_{46} : Category three of in-medium properties of hadrons

T_{44} : Category one of Field Inhomogeneities of nuclear physics

T_{45} : Category two of Field Inhomogeneities of nuclear physics

T_{46} : Category three of Field Inhomogeneities of nuclear physics

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$

$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$

$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation

coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3
$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3
$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3
$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3
$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3
$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3
$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3
$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1, 2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* $1, 2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$

are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a_{20}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3)}(T_{21},t)$

are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a_{36}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7)}(T_{37},t)$

are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a_{40}'')^{(8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8)}(T_{41},t)$

are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a_{46}'')^{(9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b_{24}')^{(4)} & -(b_{24}'')^{(4)}(G_{27},t) & -(b_{28}'')^{(5,5,)}(G_{31},t) & -(b_{32}'')^{(6,6,)}(G_{35},t) \\ & -(b_{13}'')^{(1,1,1,1)}(G,t) & -(b_{16}'')^{(2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3)}(G_{23},t) \\ & -(b_{36}'')^{(7,7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b_{25}')^{(4)} & -(b_{25}'')^{(4)}(G_{27},t) & -(b_{29}'')^{(5,5,)}(G_{31},t) & -(b_{33}'')^{(6,6,)}(G_{35},t) \\ & -(b_{14}'')^{(1,1,1,1)}(G,t) & -(b_{17}'')^{(2,2,2,2)}(G_{19},t) & -(b_{21}'')^{(3,3,3,3)}(G_{23},t) \\ & -(b_{37}'')^{(7,7,7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b_{26}')^{(4)} & -(b_{26}'')^{(4)}(G_{27},t) & -(b_{30}'')^{(5,5,)}(G_{31},t) & -(b_{34}'')^{(6,6,)}(G_{35},t) \\ & -(b_{15}'')^{(1,1,1,1)}(G,t) & -(b_{18}'')^{(2,2,2,2)}(G_{19},t) & -(b_{22}'')^{(3,3,3,3)}(G_{23},t) \\ & -(b_{38}'')^{(7,7,7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b_{24}'')^{(4)}(G_{27},t)$, $-(b_{25}'')^{(4)}(G_{27},t)$, $-(b_{26}'')^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b_{28}'')^{(5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{32}'')^{(6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b_{13}'')^{(1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1)}(G,t)$

are fourth detrition coefficients for category 1, 2 *and* 3

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

are fifth detrition coefficients for category 1, 2 *and* 3

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

are sixth detrition coefficients for category 1, 2 *and* 3

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

are seventh detrition coefficients for category 1, 2 *and* 3

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b_{32}')^{(6)} & -(b_{32}'')^{(6)}(G_{35},t) & -(b_{28}'')^{(5,5,5)}(G_{31},t) & -(b_{24}'')^{(4,4,4,)}(G_{27},t) \\ & -(b_{13}'')^{(1,1,1,1,1,1)}(G,t) & -(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b_{33}')^{(6)} & -(b_{33}'')^{(6)}(G_{35},t) & -(b_{29}'')^{(5,5,5)}(G_{31},t) & -(b_{25}'')^{(4,4,4,)}(G_{27},t) \\ & -(b_{14}'')^{(1,1,1,1,1,1)}(G,t) & -(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t) & -(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b_{34}')^{(6)} & -(b_{34}'')^{(6)}(G_{35},t) & -(b_{30}'')^{(5,5,5)}(G_{31},t) & -(b_{26}'')^{(4,4,4,)}(G_{27},t) \\ & -(b_{15}'')^{(1,1,1,1,1,1)}(G,t) & -(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t) & -(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34}$$

$-(b_{32}'')^{(6)}(G_{35},t)$, $-(b_{33}'')^{(6)}(G_{35},t)$, $-(b_{34}'')^{(6)}(G_{35},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b_{28}'')^{(5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{24}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$
$$= (a_{36})^{(7)}G_{37}$$
$$-\begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$
$$= (a_{37})^{(7)}G_{36}$$
$$-\begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$
$$= (a_{38})^{(7)}G_{37}$$
$$-\begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{44})^{(9)} & +(a''_{44})^{(9)}(T_{45},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\begin{bmatrix} (a'_{45})^{(9)} & +(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} \boxed{-(b''_{45})^{(9)}(G_{47},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} \boxed{-(b''_{46})^{(9)}(G_{47},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first

augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty} (b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

$$\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$$

$$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and $(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}$, $(r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \quad (136)$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}\left(s_{(13)}\right) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}\left(s_{(13)}\right) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} , T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}\left(s_{(16)}\right) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}\left(T_{17}\left(s_{(16)}\right), s_{(16)}\right)\right) G_{16}\left(s_{(16)}\right)\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}\left(s_{(16)}\right) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}\left(s_{(16)}\right), s_{(17)}\right)\right) G_{17}\left(s_{(16)}\right)\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}\left(s_{(16)}\right) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}\left(s_{(16)}\right), s_{(16)}\right)\right) G_{18}\left(s_{(16)}\right)\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}\left(s_{(16)}\right) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}\left(G_{19}\left(s_{(16)}\right), s_{(16)}\right)\right) T_{16}\left(s_{(16)}\right)\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A
which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that
$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$$ 169

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t\in\mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t\in\mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \le \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} |(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1T_{14}$ which leads to

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1t})+T_{14}^0e^{-\varepsilon_1t}$ If we take t such that $e^{-\varepsilon_1t}=\ \frac{1}{2}$ it results

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16{,}17{,}18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right) \quad (237)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. (241)

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})($and\ not\ on\ t$) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, $if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right) \quad 286$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$
$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le-(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le-(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}, \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}, \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(5)} and\ (b''_i)^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a''_i)^{(6)} and\ (b''_i)^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(d) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]}$, $G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^* \mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^*\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^*\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^*\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\big(\,(G_{39})^{**}\,\big) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^*\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^*\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^*\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^* \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^* \right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\Big]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\Big]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 5:

Polaritons Vis-à-vis Intersubband, Quantum Wells Vis-à-vis Hadrons, Signatures Of The Ultra Strong Light-Mater Coupling Regime Vis-à-vis Intersubband Transitions: A Apriori Determination And Aposteori Differentiation Model

1. Sub-cycle switch-on Vis-à-vis ultra strong light-matter interaction
2. light-matter interaction Vis-à-vis Microcavity
3. polaritons Vis-à-vis intersubband
4. quantum wells Vis-à-vis Hadrons
5. Signatures of the ultra strong light-mater coupling regime Vis-à-vis intersubband transitions
6. Towards Intersubband Polari tonics Vis-à-vis Light and Electrons Mate
7. Intersubband transitions Vis-à-vis ultra strong light-matter coupling
8. non-adiabatic switch-on Vis-à-vis Ultrafast phenomenon in Semiconductors
9. Extreme THz nonlinearities Vis-à-vis bulk , nanostructured semiconductors

NOTATION

Module One

Sub-cycle switch-on Vis-à-vis ultra strong light-matter interaction

G_{13} : Category one of Sub-cycle switch-on

G_{14} : Category two of Sub-cycle switch-on

G_{15} : Category three of Sub-cycle switch-on

T_{13} : Category one of ultra strong light-matter interaction

T_{14} : Category two of ultra strong light-matter interaction

T_{15} : Category three of ultra strong light-matter interaction. Two important papers are cited. (Sub-cycle switch-on of ultra strong light-matter interaction: Günter G, Anappara AA, And Hees J, Sell A, Biasiol G, Sorba L, De Liberato S, Ciuti C, Tredicucci A, Leitenstorfer A, And Huber R.: Controlling the way light interacts with material excitations is at the heart of cavity quantum electrodynamics (QED). In the strong-coupling regime, quantum emitters in a microresonator absorb and spontaneously re-emit a photon many times before dissipation becomes effective, giving rise to mixed light-matter eigenmodes. Recent experiments in semiconductor microcavities reached a new limit of ultrastrong coupling, where

photon exchange occurs on timescales comparable to the oscillation period of light. In this limit, ultrafast modulation of the coupling strength has been suggested to lead to unconventional QED phenomena. Although sophisticated light-matter coupling has been achieved in all three spatial dimensions, control in the fourth dimension, time, is little developed. Here we use a quantum-well waveguide structure to optically tune light-matter interaction from weak to ultrastrong and turn on maximum coupling within less than one cycle of light. In this regime, a class of extremely non-adiabatic phenomena becomes observable. In particular, we directly monitor how a coherent photon population converts to cavity polaritons during abrupt switching. This system forms a promising laboratory in which to study novel sub-cycle QED effects and represents an efficient room-temperature switching device operating at unprecedented speed. Ultrastrong Light-Matter Coupling Regime with Polariton Dots: Y. Todorov1,*, A. M. Andrews2, R. Colombelli3, S. De Liberato1,4, C. Ciuti1, P. Klang2, G. Strasser2, and C. Sirtori1 :The regime of ultrastrong light-matter interaction has been investigated theoretically and experimentally, using zero-dimensional electromagnetic resonators coupled with an electronic transition between two confined states of a semiconductor quantum well. We have measured a splitting between the coupled modes that amounts to 48% of the energy transition, the highest ratio ever observed in a light-matter coupled system. Our analysis, based on a microscopic quantum theory, shows that the nonlinear polariton splitting, a signature of this regime, is a dynamical effect arising from the self-interaction of the collective electronic polarization with its own emitted field).

Module two

Light-matter interaction vis-à-vis microcavity

G_{16} : Category one of light-matter interaction (Light-Matter Interaction: Meystre, Pierre: Optical physics is concerned with the dynamical interactions of atoms and molecules with electromagnetic to fields. Semiclassical theories, which study the interaction of atoms with classical fields, are often said to comprise optical physics, while quantum optics treats the interaction of atoms or molecules with quantized electromagnetic fields. A significant part of optical physics and quantum optics is the study of near-resonant atom-field interactions, and concentrates on nonperturbative dynamics, where the effects of the optical fields have to be kept to all orders. The atomic properties themselves are assumed to be known.Dr. Ulrich Rössler's investigation centres on the investigation of condensed matter systems using light as a probe is a versatile and powerful concept, which provides various kinds of information not only about the structure but also of the electronic and lattice excitations and their dynamics. Besides linear optical processes, like absorption and emission (including photoemission), light-scattering, two- (or three-) photon absorption, photo-luminescence and high-excitation spectroscopy also belong to the methods to gain information about the single-particle and collective excitations of a solid. In semiconductor samples with a proper design, coherent light emission is possible and used in solid-state lasers. These methods, which allow studies in the frequency or time domain, are all based on the interaction of light with matter.)

G_{17} : Category two of light-matter interaction

G_{18} : Category three of light-matter interaction

T_{16} : Category one of Microcavity

T_{17} : Category two of Microcavity

T_{18} : Category three of Microcavity

Module three

Polaritons Vis-à-vis intersubband

G_{20} : Category one of polaritons

G_{21} : Category two of polaritons

G_{22} : Category three of polaritons

T_{20} : Category one of intersubband

T_{21} : Category two of intersubband

T_{22} : Category three of intersubband

Module four

Quantum wells Vis-à-vis Hadrons

G_{24} : Category one of quantum wells

G_{25} : Category two of quantum wells

G_{26} : Category three of quantum wells

T_{24} : Category one of Hadrons

T_{25} : Category two of Hadrons

T_{26} : Category three of Hadrons

Module five

Signatures of the ultra strong light-mater coupling regime Vis-à-vis intersubband transitions

G_{28} : Category one of Signatures of the ultra strong light-mater coupling regime

G_{29} : Category two of Signatures of the ultra strong light-mater coupling regime

G_{30} : Category three of Signatures of the ultra strong light-mater coupling regime

T_{28} : Category one of intersubband transitions

T_{29} : Category two of intersubband transitions

T_{30} : Category three of intersubband transitions

Intersubband Polari tonics Vis-à-vis Light and Electrons Mate

Module six

G_{32} : Category one of Intersubband Polari tonics

G_{33} : Category two of Intersubband Polari tonics

G_{34} : Category three of Intersubband Polari tonics

T_{32} : Category one of Electrons Mate

T_{33} : Category two of Electrons Mate

T_{34} : Category three of Electrons Mate

Module seven

Intersubband transitions Vis-à-vis ultra strong light-matter coupling

G_{36} : Category one of Intersubband transitions

G_{37} : Category two of Intersubband transitions

G_{38} : Category three of Intersubband transitions

T_{36} : Category one of ultra strong light-matter coupling

T_{37} : Category two of ultra strong light-matter coupling

T_{38} : Category three of ultra strong light-matter coupling

Module eight

Non-adiabatic switch-on Vis-à-vis Ultrafast phenomenon in Semiconductors

G_{40} : Category one of non-adiabatic switch-on

G_{41} : Category two of non-adiabatic switch-on

G_{42} : Category three of non-adiabatic switch-on

T_{40} : Category one of Ultrafast phenomenon in Semiconductors

T_{41} : Category two of Ultrafast phenomenon in Semiconductors

T_{42} : Category three of Ultrafast phenomenon in Semiconductors

Module Nine

Extreme THz nonlinearities Vis-à-vis bulk , nanostructured semiconductors

G_{44} : Category one of Extreme THz nonlinearities

G_{45} : Category two of Extreme THz nonlinearities

G_{46} : Category three of Extreme THz nonlinearities

T_{44} : Category one of bulk , nanostructured semiconductors

T_{45} : Category two of bulk , nanostructured semiconductors

T_{46} : Category three of bulk , nanostructured semiconductors

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}$ $(a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$: $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$

$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$

$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$

$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$

$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}),t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \tag{67}$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}+(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{22} \tag{69}$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \tag{70}$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \tag{71}$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & \boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \tag{72}$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & \boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \tag{73}$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \tag{74}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \tag{75}$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ are third augmentation

$coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34}$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$
$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$
$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$
$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{37}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} & +(a''_{41})^{(8)}(T_{41},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) \\ & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) \\ & +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) \\ & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) \\ & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & \end{array}\right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) \\ & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) \\ & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\left[\begin{array}{ccc} (b'_{42})^{(8)} & \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} & \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \\ \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} & & \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\left[\begin{array}{ccc} (a'_{44})^{(9)} & \boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \\ \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} & \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \\ \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} & \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & & \end{array}\right] G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\left[\begin{array}{ccc} (a'_{45})^{(9)} & \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \\ \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} & \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \\ \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} & \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & & \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} \boxed{-(b''_{45})^{(9)}(G_{47},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} \boxed{-(b''_{46})^{(9)}(G_{47},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first

augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty} (b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

$$\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$$

$$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and $(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}$, $(r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \quad (136)$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} , T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A
which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that
$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$$ 169

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1T_{14}$ which leads to

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1t})+T_{14}^0e^{-\varepsilon_1t}$ If we take t such that $e^{-\varepsilon_1t}=\frac{1}{2}$ it results

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)} \left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)} \left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{32})^{(6)}|G_{33}^{(1)} - G_{33}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}|G_{32}^{(1)} - G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right)$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 250

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)}t\right)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If $G_{32} \; is$ bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\,((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(7)}-(a''_i)^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b''_i)^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, $if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$
$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le -(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le -(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le -(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le -(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, $\boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, $\boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\Big(\frac{(a_{46})^{(9)} G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\Big] + G^0_{46} e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \frac{(a_{46})^{(9)} G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\big[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\big] + G^0_{46} e^{-(a'_{46})^{(9)} t}\Big)$$

$$\boxed{T^0_{44} e^{(R_1)^{(9)} t} \le T_{44}(t) \le T^0_{44} e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(5)} and\ (b''_i)^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a''_i)^{(6)} and\ (b''_i)^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(e) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]}$, $G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j} (\, G^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b'_{13})^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \textstyle\sum_{j=13}^{15}\big(s_{(13)(j)} T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b'_{14})^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \textstyle\sum_{j=13}^{15}\big(s_{(14)(j)} T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b'_{15})^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \textstyle\sum_{j=13}^{15}\big(s_{(15)(j)} T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}} (T_{17}^{*}) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j} (\, (G_{19})^{*}\,) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a'_{16})^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^{*}\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\big(\,(G_{39})^{**}\,\big) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\Big)\right]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\Big)\right]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\Big)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29} + (b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29} + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29} + (b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}\{\left((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\Big)\right]$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\Big)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}\{\left((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\Big)\right]$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\Big)$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2 + \left((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 6:

Drag on particles in a nematic suspension Vis-à-vis moving nematic-isotropic interface, Defect structures and torque on an elongated colloidal particle immersed Vis-à-vis a liquid crystal host with topological defects, Entropic torque. Vis-à-vis depletion effects: A Herpestidae Mongoose and Ineluctable Aquiline Model

1. Electrophonic sounds Vis-à-vis from Leonid fireballs
2. Time resolved diagnostics of plasmas Vis-à-vis ablative capillary discharges
3. Meteorites Vis-à-vis electrophonic sounds
4. Numerical modeling of laser beam-target interaction Vis-à-vis the plasma formation threshold
5. Lasing in an ablative capillary discharge Vis-à-vis structured return conductor.
6. Rogowski coil Vis-à-vis fast pulsed-current measurements of capillary discharge
7. Drag on particles in a nematic suspension Vis-à-vis moving nematic-isotropic interface
8. Defect structures and torque on an elongated colloidal particle immersed Vis-à-vis a liquid crystal host with topological defects
9. Entropic torque Vis-à-vis depletion effects

NOTATION

Module One

electrophonic sounds Vis-à-vis Leonid fireballs

G_{13} : Category one of electrophonic sounds

G_{14} : Category two of electrophonic sounds

G_{15} : Category three of electrophonic sounds

T_{13} : Category one of Leonid fireballs

T_{14} : Category two of Leonid fireballs

T_{15} : Category three of Leonid fireballs

Module Two

Time resolved diagnostics of plasmas Vis-à-vis ablative capillary discharges

G_{16} : Category one of Time resolved diagnostics of plasmas

G_{17} : Category two of Time resolved diagnostics of plasmas

G_{18} : Category three of Time resolved diagnostics of plasmas

T_{16} : Category one of ablative capillary discharges

T_{17} : Category two of ablative capillary discharges

T_{18} : Category three of ablative capillary discharges

Meteorites Vis-à-vis electrophonic sounds

Module three

G_{20} : Category one of Meteorites

G_{21} : Category two of Meteorites

G_{22} : Category three of Meteorites

T_{20} : Category one of electrophonic sounds

T_{21} : Category two of electrophonic sounds

T_{22} : Category three of electrophonic sounds

Module four

Numerical modeling of laser beam-target interaction Vis-à-vis the plasma formation threshold

G_{24} : Category one of Numerical modeling of laser beam-target interaction

G_{25} : Category two of Numerical modeling of laser beam-target interaction

G_{26} : Category three of Numerical modeling of laser beam-target interaction

T_{24} : Category one of the plasma formation threshold

T_{25} : Category two of the plasma formation threshold

T_{26} : Category three of the plasma formation threshold

Lasing in an ablative capillary discharge Vis-à-vis structured return conductor

Module five

G_{28} : Category one of Lasing in an ablative capillary discharge

G_{29} : Category two of Lasing in an ablative capillary discharge

G_{30} : Category three of Lasing in an ablative capillary discharge

T_{28} : Category one of structured return conductor

T_{29} : Category two of structured return conductor

T_{30} : Category three of structured return conductor

Module six

Rogowski coil Vis-à-vis fast pulsed-current measurements of capillary discharge

G_{32} : Category one of Rogowski coil

G_{33} : Category two of Rogowski coil

G_{34} : Category three of Rogowski coil

T_{32} : Category one of fast pulsed-current measurements of capillary discharge

T_{33} : Category two of fast pulsed-current measurements of capillary discharge

T_{34} : Category three of fast pulsed-current measurements of capillary discharge

Module seven

Drag on particles in a nematic suspension Vis-à-vis moving nematic-isotropic interface

G_{36} : Category one of Drag on particles in a nematic suspension

G_{37} : Category two of Drag on particles in a nematic suspension

G_{38} : Category three of Drag on particles in a nematic suspension

T_{36} : Category one of moving nematic-isotropic interface

T_{37} : Category two of moving nematic-isotropic interface

T_{38} : Category three of moving nematic-isotropic interface

Module eight

Defect structures and torque on an elongated colloidal particle immersed Vis-à-vis a liquid crystal host with topological defects

G_{40} : Category one of Defect structures and torque on an elongated colloidal particle immersed

G_{41} : Category two of Defect structures and torque on an elongated colloidal particle immersed

G_{42} : Category three of Defect structures and torque on an elongated colloidal particle immersed

T_{40} : Category one of liquid crystal host with topological defects

T_{41} : Category two of liquid crystal host with topological defects

T_{42} : Category three of liquid crystal host with topological defects

Module Nine

Entropic torque Vis-à-vis depletion effects

G_{44} : Category one of Entropic torque

G_{45} : Category two of Entropic torque

G_{46} : Category three of Entropic torque

T_{44} : Category one of depletion effects

T_{45} : Category two of depletion effects

T_{46} : Category three of depletion effects

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt}=(b_{13})^{(1)}T_{14}-\begin{bmatrix}(b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt}=(b_{14})^{(1)}T_{13}-\begin{bmatrix}(b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt}=(b_{15})^{(1)}T_{14}-\begin{bmatrix}(b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)}\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients

for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{c} (a_{28}')^{(5)}\ +(a_{28}'')^{(5)}(T_{29},t)\ +(a_{24}'')^{(4,4,)}(T_{25},t)\ +(a_{32}'')^{(6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)\ +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)\ +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)\ +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)\ +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{c} (a_{29}')^{(5)}\ +(a_{29}'')^{(5)}(T_{29},t)\ +(a_{25}'')^{(4,4,)}(T_{25},t)\ +(a_{33}'')^{(6,6,6)}(T_{33},t) \\ +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)\ +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)\ +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)\ +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)\ +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{c} (a_{30}')^{(5)}\ +(a_{30}'')^{(5)}(T_{29},t)\ +(a_{26}'')^{(4,4,)}(T_{25},t)\ +(a_{34}'')^{(6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)\ +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)\ +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)\ +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)\ +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{30} \quad 81$$

$Where\ +(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ $are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ +(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ $are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ $are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
are first detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$
$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$
$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b_{28}'')^{(5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{24}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad (92)$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad (93)$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)} & -(b_{36}'')^{(7)}(G_{39},t) & -(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t) & -(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)} & -(b_{37}'')^{(7)}(G_{39},t) & -(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t) & -(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)} & -(b_{38}'')^{(7)}(G_{39},t) & -(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t) & -(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b_{36}'')^{(7)}(G_{39},t)$, $-(b_{37}'')^{(7)}(G_{39},t)$, $-(b_{38}'')^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

95

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right]G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right]G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right]G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} -$$

$$\begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} -$$

$$\begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} -$$

$$\begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}},\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)}+(a_i')^{(1)}+(\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)}+(b_i')^{(1)}+(\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)},\ (\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le(\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le\ (r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\big((G_{19}),t\big)=\ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)},(\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)},(\hat{B}_{16})^{(2)},(p_i)^{(2)},\ (r_i)^{(2)}}$ are positive constants and $\boxed{i=16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t)-(a_i'')^{(2)}(T_{17},t)|\le(\hat{k}_{16})^{(2)}|T_{17}-T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t)-(b_i'')^{(2)}\big((G_{19}),t\big)|<(\hat{k}_{16})^{(2)}||(G_{19})-(G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)},(\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \ , \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\widehat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\widehat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\widehat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\widehat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\widehat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\ ,\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants

$(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$ 132

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$:

137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$: B

$$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A
which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$$ 169

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t\in\mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t\in\mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1T_{14}$ which leads to

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1t})+T_{14}^0e^{-\varepsilon_1t}$ If we take t such that $e^{-\varepsilon_1t}=\frac{1}{2}$ it results

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad (191)$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad (192)$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{\left(-(b_i')^{(3)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right) \quad (237)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. (241)

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)})\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, $if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$
$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$$

$$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$$

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le -(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le -(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le -(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le -(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2+(\sigma_2)^{(3)}\nu^{(3)}-(a_{20})^{(3)}\right)\leq\frac{d\nu^{(3)}}{dt}\leq-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2+(\sigma_1)^{(3)}\nu^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(\nu_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\nu_1)^{(3)}<(\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t)\geq\frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)}\leq\nu^{(3)}(t)\leq(\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t)\leq\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)}\leq\nu^{(3)}(t)\leq(\bar{\nu}_1)^{(3)}$

If $0<(\nu_1)^{(3)}<(\nu_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)}\leq\frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}\leq\nu^{(3)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}\leq(\bar{\nu}_1)^{(3)}$$

If $0<(\nu_1)^{(3)}\leq(\bar{\nu}_1)^{(3)}\leq(\nu_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)}\leq\nu^{(3)}(t)\leq\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}\leq(\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)}\leq\nu^{(3)}(t)\leq(m_1)^{(3)},\quad\boxed{\nu^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\leq u^{(3)}(t)\leq(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)}=(\bar{\nu}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(5)} and\ (b''_i)^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a''_i)^{(6)} and\ (b''_i)^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(f) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^* \mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}(\,(G_{39})^{**}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29} + (b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29} + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29} + (b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}\{\left((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\Big)\right]$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\Big)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}\{\left((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\Big)\right]$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\Big)$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2 + \left((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 7:

Spatial profiles of fusion product flux Vis-à-vis the gas dynamic trap with deuterium neutral beam injection, Fusion regimes of axisymmetric open system Vis-à-vis powerful injection of fast particles, Enhancement of order fluctuations Vis-à-vis nematic liquid crystal by sonication (nematic, acoustic): A Filibustered Conventicler- Serene canopies and Trellises Model

INTRODUCTION—VARIABLES USED

1. confined liquid Vis-à-vis density functional,
2. Boundary slip Vis-à-vis of a prewetting transition
3. Capillary bridging Vis-à-vis long-range attractive forces in a mean-field approach
4. Vanishing Perturbative Vacuum Energy Vis-à-vis non-Supersymmetric Orientifolds
5. Axially Symmetric Gas Dynamic Trap and Magnetic confinement
6. Spatial profiles of fusion product flux Vis-à-vis the gas dynamic trap with deuterium neutral beam injection
7. Fusion regimes of axisymmetric open system Vis-à-vis powerful injection of fast particles
8. Enhancement of order fluctuations Vis-à-vis nematic liquid crystal by sonication (nematic acoustic)
9. Low-frequency NMR relaxometry Vis-à-vis spatially constrained liquid crystals

NOTATION

Module One

confined liquid Vis-à-vis density functional

G_{13} : Category one of confined liquid

G_{14} : Category two of confined liquid

G_{15} : Category three of confined liquid

T_{13} : Category one of density functional

T_{14} : Category two of density functional

T_{15} : Category three of density functional

Module Two

Boundary slip Vis-à-vis prewetting transitions

G_{16} : Category one of Boundary slip

G_{17} : Category two of Boundary slip

G_{18} : Category three of Boundary slip

T_{16} : Category one of prewetting transitions

T_{17} : Category two of prewetting transitions

T_{18} : Category three of prewetting transitions

Module three

Capillary bridging Vis-à-vis long-range attractive forces in a mean-field approach

G_{20} : Category one of Capillary bridging

G_{21} : Category two of Capillary bridging

G_{22} : Category three of Capillary bridging

T_{20} : Category one of long-range attractive forces in a mean-field approach

T_{21} : Category two of long-range attractive forces in a mean-field approach

T_{22} : Category three of long-range attractive forces in a mean-field approach

Module four

Vanishing Perturbative Vacuum Energy Vis-à-vis non-Supersymmetric Orientifolds

G_{24} : Category one of Vanishing Perturbative Vacuum Energy

G_{25} : Category two of Vanishing Perturbative Vacuum Energy

G_{26} : Category three of Vanishing Perturbative Vacuum Energy

T_{24} : Category one of non-Supersymmetric Orientifolds

T_{25} : Category two of non-Supersymmetric Orientifolds

T_{26} : Category three of non-Supersymmetric Orientifolds

Module five

Spatial profiles of fusion product flux Vis-à-vis the gas dynamic trap with deuterium neutral beam injection

G_{28} : Category one of Spatial profiles of fusion product flux

G_{29} : Category two of Spatial profiles of fusion product flux

G_{30} : Category three of Spatial profiles of fusion product flux

T_{28} : Category one of Vis the gas dynamic trap with deuterium neutral beam injection

T_{29} : Category two of Vis the gas dynamic trap with deuterium neutral beam injection

T_{30} : Category three of Vis the gas dynamic trap with deuterium neutral beam injection

Module six

Fusion regimes of axisymmetric open system Vis-à-vis powerful injection of fast particles

G_{32} : Category one of Fusion regimes of axisymmetric open system

G_{33} : Category two of Fusion regimes of axisymmetric open system

G_{34} : Category three of Fusion regimes of axisymmetric open system

T_{32} : Category one of powerful injection of fast particles

T_{33} : Category two of powerful injection of fast particles

T_{34} : Category three of powerful injection of fast particles

Module seven

Axially Symmetric Gas Dynamic Trap and Magnetic confinement

G_{36} : Category one of Axially Symmetric Gas Dynamic Trap

G_{37} : Category two of Axially Symmetric Gas Dynamic Trap

G_{38} : Category three of Axially Symmetric Gas Dynamic Trap

T_{36} : Category one of Magnetic confinement

T_{37} : Category two of Magnetic confinement

T_{38} : Category three of Magnetic confinement

Module eight

Enhancement of order fluctuations Vis-à-vis nematic liquid crystal by sonication (nematic, acoustic)

G_{40} : Category one of Enhancement of order fluctuations

G_{41} : Category two of Enhancement of order fluctuations

G_{42} : Category three of Enhancement of order fluctuations

T_{40} : Category one of nematic liquid crystal by sonication (nematic, acoustic)

T_{41} : Category two of nematic liquid crystal by sonication (nematic, acoustic

T_{42} : Category three of nematic liquid crystal by sonication (nematic, acoustic)

Module Nine

Low-frequency NMR relaxometry Vis-à-vis spatially constrained liquid crystals

G_{44} : Category one of Low-frequency NMR relaxometry

G_{45} : Category two of Low-frequency NMR relaxometry

G_{46} : Category three of Low-frequency NMR relaxometry

T_{44} : Category one of spatially constrained liquid crystals

T_{45} : Category two of spatially constrained liquid crystals

T_{46} : Category three of spatially constrained liquid crystals

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}$ $(a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$: $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$, $(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$, $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}),t)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}),t)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}),t)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}),t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) \quad +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) \quad +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) \quad +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right]G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20} \quad (67)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients

for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & +(a_{28}'')^{(5)}(T_{29},t) & +(a_{24}'')^{(4,4,)}(T_{25},t) \\ +(a_{32}'')^{(6,6,6)}(T_{33},t) & +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t) \\ +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) & +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t) \\ +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) & & \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & +(a_{29}'')^{(5)}(T_{29},t) & +(a_{25}'')^{(4,4,)}(T_{25},t) \\ +(a_{33}'')^{(6,6,6)}(T_{33},t) & +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t) \\ +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) & +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t) \\ +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) & & \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & +(a_{30}'')^{(5)}(T_{29},t) & +(a_{26}'')^{(4,4,)}(T_{25},t) \\ +(a_{34}'')^{(6,6,6)}(T_{33},t) & +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t) \\ +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) & +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t) \\ +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) & & \end{array}\right] G_{30} \quad 81$$

$Where\ +(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ $are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ +(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ $are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ $are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
are first detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32} \quad (85)$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33} \quad (86)$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{matrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right]G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \left[\begin{matrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{matrix}\right]T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \left[\begin{matrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{matrix}\right]T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \left[\begin{matrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{matrix}\right]T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b_{28}'')^{(5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{24}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad (92)$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad (93)$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $-(b_{36}'')^{(7)}(G_{39},t)$, $-(b_{37}'')^{(7)}(G_{39},t)$, $-(b_{38}'')^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41} - \left[\begin{matrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{matrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{matrix} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} -$$

$$\left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} -$$

$$\left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} -$$

$$\left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{46})^{(9)} & \boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}},\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)}+(a_i')^{(1)}+(\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)}+(b_i')^{(1)}+(\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le(\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le(r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\left((G_{19}),t\right)=(r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)},(\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)},(\hat{B}_{16})^{(2)},(p_i)^{(2)},\ (r_i)^{(2)}}$ are positive constants and $\boxed{i=16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t)-(a_i'')^{(2)}(T_{17},t)|\le(\hat{k}_{16})^{(2)}|T_{17}-T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t)-(b_i'')^{(2)}\left((G_{19}),t\right)|<(\hat{k}_{16})^{(2)}||(G_{19})-(G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)},(\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\widehat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\widehat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\widehat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\widehat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\widehat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}\ (T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\ \big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\ ,\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants

$(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$ 132

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$:

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad (138)$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad (139)$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad (140)$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad (141)$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \quad (142)$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \quad (143)$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad (144)$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad (145)$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad (146)$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \quad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44{,}45{,}46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$: B

$$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^{0} + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^{0} + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{32})^{(6)}, T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A
which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that
$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$$ 169

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t\in\mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t\in\mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0 \quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1, T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1 T_{14}$ which leads to

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1 t})+T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t}=\frac{1}{2}$ it results

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16{,}17{,}18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$

$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. (241)

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(7)}$ and $(b''_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 261

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(7)}-(a''_i)^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b''_i)^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, *if* $(\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$$

$$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$$

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le-(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le-(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)}+(\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)}+(\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)}+(C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)}-(v_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)}-(v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)}-(v_0)^{(2)}}{(v_0)^{(2)}-(v_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \le u^{(2)}(t) \le (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}, \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}, \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484A$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(g) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j} (\, G^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b'_{13})^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \textstyle\sum_{j=13}^{15}\big(s_{(13)(j)} T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b'_{14})^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \textstyle\sum_{j=13}^{15}\big(s_{(14)(j)} T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b'_{15})^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \textstyle\sum_{j=13}^{15}\big(s_{(15)(j)} T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}} (T_{17}^{*}) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j} (\, (G_{19})^{*}\,) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a'_{16})^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^*\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^*\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^*\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\big(\,(G_{39})^{**}\,\big) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^*\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^*\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^*\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29} + (b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29} + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29} + (b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}\{\left((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\Big)\right]$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\Big)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}\{\left((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\Big)\right]$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\Big)$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2 + \left((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 8:

Proton field-cycling nuclear magnetic resonance relaxometry Vis-à-vis the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication, Proton spin-lattice relaxation in a liquid crystalvis Vis-à-vis Aerosol complex above the bulk isotropization temperature:

1. Apparent low-field dispersion in the smectic Vis-à-vis A mesophase of thermotropic cyanobiphenyls
2. Proton field-cycling nuclear magnetic resonance relaxometry Vis-à-vis the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication.
3. Proton spin-lattice relaxation in a liquid crystal Vis-à-vis Aerosol complex above the bulk isotropization temperature
4. Field-cycling NMR relaxometry Vis-à-vis rotating-frame spin-lattice relaxation
5. Aerosol Vis-à-vis order director fluctuations, Monte Carlo
6. Field-cycling NMR relaxometry Vis-à-vis spin-lattice relaxation
7. Rouse model Vis-à-vis molecular dynamics.
8. The "Corset Effect" of Spin-Lattice Relaxation in Polymer Melts Vis-à-vis Nanoporous Media
9. Acoustically stimulated NMR relaxometry Vis-à-vis ultrasound order fluctuations

NOTATION

Module One

Apparent low-field dispersion in the smectic Vis-à-vis mesophase of thermotropic cyanobiphenyls

G_{13} : Category one of Apparent low-field dispersion in the smectic

G_{14} : Category two of Apparent low-field dispersion in the smectic

G_{15} : Category three of Apparent low-field dispersion in the smectic

T_{13} : Category one of mesophase of thermotropic cyanobiphenyls

T_{14} : Category two of mesophase of thermotropic cyanobiphenyls

T_{15} : Category three of mesophase of thermotropic cyanobiphenyls

Module Two

Proton field-cycling nuclear magnetic resonance relaxometry Vis-à-vis the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication

G_{16} : Category one of Proton field-cycling nuclear magnetic resonance relaxometry

G_{17} : Category two of Proton field-cycling nuclear magnetic resonance relaxometry

G_{18} : Category three of Proton field-cycling nuclear magnetic resonance relaxometry

T_{16} : Category one of the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication

T_{17} : Category two of the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication

T_{18} : Category three of the smectic A mesophase of thermotropic cyanobiphenyls: effects of sonication

Module three

Proton spin-lattice relaxation in a liquid crystal Vis-à-vis Aerosol complex above the bulk isotropization temperature

G_{20} : Category one of Proton spin-lattice relaxation in a liquid crystal

G_{21} : Category two of Proton spin-lattice relaxation in a liquid crystal

G_{22} : Category three of Proton spin-lattice relaxation in a liquid crystal

T_{20} : Category one of Aerosol complex above the bulk isotropization temperature

T_{21} : Category two of Aerosol complex above the bulk isotropization temperature

T_{22} : Category three of Aerosol complex above the bulk isotropization temperature

Module four

Aerosol Vis-à-vis order director fluctuations, Monte Carlo

G_{24} : Category one of Aerosol

G_{25} : Category two of Aerosol

G_{26} : Category three of Aerosol

T_{24} : Category one of order director fluctuations, Monte Carlo

T_{25} : Category two of order director fluctuations, Monte Carlo

T_{26} : Category three of order director fluctuations, Monte Carlo

Module five

Field-cycling NMR relaxometry Vis-à-vis rotating-frame spin-lattice relaxation

G_{28} : Category one of Field-cycling NMR relaxometry

G_{29} : Category two of Field-cycling NMR relaxometry

G_{30} : Category three of Field-cycling NMR relaxometry

T_{28} : Category one of rotating-frame spin-lattice relaxation

T_{29} : Category two of rotating-frame spin-lattice relaxation

T_{30} : Category three of rotating-frame spin-lattice relaxation

Module six

Field-cycling NMR relaxometry Vis-à-vis spin-lattice relaxation

G_{32} : Category one of Field-cycling NMR relaxometry

G_{33} : Category two of Field-cycling NMR relaxometry

G_{34} : Category three of Field-cycling NMR relaxometry

T_{32} : Category one of spin-lattice relaxation

T_{33} : Category two of spin-lattice relaxation

T_{34} : Category three of spin-lattice relaxation

Module seven

Rouse model Vis-à-vis molecular dynamics

G_{36} : Category one of Rouse model

G_{37} : Category two of Rouse model

G_{38} : Category three of Rouse model

T_{36} : Category one of molecular dynamics

T_{37} : Category two of molecular dynamics

T_{38} : Category three of molecular dynamics

Module eight

The "Corset Effect" of Spin-Lattice Relaxation in Polymer Melts Vis-à-vis Nanoporous Media

G_{40} : Category one of The "Corset Effect" of Spin-Lattice Relaxation in Polymer Melts

G_{41} : Category two of The "Corset Effect" of Spin-Lattice Relaxation in Polymer Melts

G_{42} : Category three of The "Corset Effect" of Spin-Lattice Relaxation in Polymer Melts

T_{40} : Category one of Nanoporous Media

T_{41} : Category two of Nanoporous Media

T_{42} : Category three of Nanoporous Media

Module Nine

Acoustically stimulated NMR relaxometry Vis-à-vis ultrasound order fluctuations

G_{44} : Category one of Acoustically stimulated NMR relaxometry

G_{45} : Category two of Acoustically stimulated NMR relaxometry

G_{46} : Category three of Acoustically stimulated NMR relaxometry

T_{44} : Category one of ultrasound order fluctuations

T_{45} : Category two of ultrasound order fluctuations

T_{46} : Category three of ultrasound order fluctuations

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt}=(b_{13})^{(1)}T_{14}-\begin{bmatrix}(b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt}=(b_{14})^{(1)}T_{13}-\begin{bmatrix}(b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt}=(b_{15})^{(1)}T_{14}-\begin{bmatrix}(b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}\\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)}\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients

for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* $1, 2\ 3$

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category $1, 2$ *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{16}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2)}(G_{19},t)$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{28}')^{(5)} & +(a_{28}'')^{(5)}(T_{29},t) & +(a_{24}'')^{(4,4,)}(T_{25},t) \quad +(a_{32}'')^{(6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a_{29}')^{(5)} & +(a_{29}'')^{(5)}(T_{29},t) & +(a_{25}'')^{(4,4,)}(T_{25},t) \quad +(a_{33}'')^{(6,6,6)}(T_{33},t) \\ +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a_{30}')^{(5)} & +(a_{30}'')^{(5)}(T_{29},t) & +(a_{26}'')^{(4,4,)}(T_{25},t) \quad +(a_{34}'')^{(6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{30} \quad 81$$

$Where\ +(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ $are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ +(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ $are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ $are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a_{40}'')^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
are first detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$
$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$
$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{matrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right]G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \left[\begin{matrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)}\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{matrix}\right]T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \left[\begin{matrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)}\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{matrix}\right]T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \left[\begin{matrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)}\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{matrix}\right]T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$-(b_{28}'')^{(5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b_{24}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b_{13}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1)}(G,t)$, $-(b_{15}'')^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b_{16}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)} & +(a_{36}'')^{(7)}(T_{37},t) & +(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)} & +(a_{37}'')^{(7)}(T_{37},t) & +(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad (92)$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)} & +(a_{38}'')^{(7)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad (93)$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)} & -(b_{36}'')^{(7)}(G_{39},t) & -(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t) & -(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)} & -(b_{37}'')^{(7)}(G_{39},t) & -(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t) & -(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)} & -(b_{38}'')^{(7)}(G_{39},t) & -(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t) & -(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b_{36}'')^{(7)}(G_{39},t)$, $-(b_{37}'')^{(7)}(G_{39},t)$, $-(b_{38}'')^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} -$$

$$\left[\begin{array}{ccc} (b'_{40})^{(8)} & \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} -$$

$$\left[\begin{array}{ccc} (b'_{41})^{(8)} & \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} -$$

$$\left[\begin{array}{ccc} (b'_{42})^{(8)} & \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\left[\begin{array}{ccc} (a'_{44})^{(9)} & \boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$\frac{dT_{46}}{dt} =$

$(b_{46})^{(9)}T_{45} -$

$$\begin{bmatrix} (b'_{46})^{(9)} & \boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}},\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)}+(a_i')^{(1)}+(\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)}+(b_i')^{(1)}+(\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le(\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le(r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\left((G_{19}),t\right)=(r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)},(\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)},(\hat{B}_{16})^{(2)},(p_i)^{(2)},\ (r_i)^{(2)}}$ are positive constants and $\boxed{i=16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t)-(a_i'')^{(2)}(T_{17},t)|\le(\hat{k}_{16})^{(2)}|T_{17}-T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t)-(b_i'')^{(2)}\left((G_{19}),t\right)|<(\hat{k}_{16})^{(2)}||(G_{19})-(G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)},(\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\widehat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\widehat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\widehat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\widehat{M}_{20})^{(3)}]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\widehat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ ,\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\,(a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\,(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ \ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, \ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \le \ (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\,\big((G_{27}), t\big) = \ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - \ T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\big[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\big]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\ ,\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants

$(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - \ T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$ 132

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42$$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$:

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad (138)$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \quad (139)$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad (140)$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad (141)$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \quad (142)$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \quad (143)$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad (144)$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad (145)$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad (146)$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \quad (146A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$: B

$$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''^{(1)}_{13}\right)\left(T_{14}(s_{(13)}), s_{(13)}\right)\right) G_{13}(s_{(13)})\Big] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}\left(T_{14}(s_{(13)}), s_{(13)}\right)\right) G_{14}(s_{(13)})\Big] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}\left(T_{14}(s_{(13)}), s_{(13)}\right)\right) G_{15}(s_{(13)})\Big] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}\left(G(s_{(13)}), s_{(13)}\right)\right) T_{13}(s_{(13)})\Big] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}\left(G(s_{(13)}), s_{(13)}\right)\right) T_{14}(s_{(13)})\Big] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}\left(G(s_{(13)}), s_{(13)}\right)\right) T_{15}(s_{(13)})\Big] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''^{(2)}_{16}\right)\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{16}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}\left(T_{17}(s_{(16)}), s_{(17)}\right)\right) G_{17}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}\left(T_{17}(s_{(16)}), s_{(16)}\right)\right) G_{18}(s_{(16)})\Big] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^{0} + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^{0} + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^{0} + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{32})^{(6)}, T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167 that

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+$ 169
$\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)} \left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1T_{14}$ which leads to

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1t})+T_{14}^0e^{-\varepsilon_1t}$ If we take t such that $e^{-\varepsilon_1t}=\frac{1}{2}$ it results

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16{,}17{,}18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. (241)

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}\left|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq$ 259

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, *if* $(v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$
$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$ 333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$ 334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$ 335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$ 336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le -(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le -(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le -(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le -(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, $\boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, $\boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\Big(\frac{(a_{46})^{(9)} G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\Big] + G^0_{46} e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \frac{(a_{46})^{(9)} G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\big[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\big] + G^0_{46} e^{-(a'_{46})^{(9)} t}\Big)$$

$$\boxed{T^0_{44} e^{(R_1)^{(9)} t} \le T_{44}(t) \le T^0_{44} e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)}+(\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)}+(\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)}+(C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)}-(v_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)}-(v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)}-(v_0)^{(2)}}{(v_0)^{(2)}-(v_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t) v^{(9)} - (a_{45})^{(9)} v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)} v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)} v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right) t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right) t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right) t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right) t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(h) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]}$, $G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^* \mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^*\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^*\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^*\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}$, $\frac{\partial(b_i'')^{(7)}}{\partial G_j}(\,(G_{39})^{*^*}\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^*\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^*\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^*\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(28)}T^*_{29} + (b_{29})^{(5)}s_{(28),(28)}T^*_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^2 + \left((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^2 + \left((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^*_{29} + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^*_{28}\right)$$

$$\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)}\right)s_{(29),(30)}T^*_{29} + (b_{29})^{(5)}s_{(28),(30)}T^*_{28}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}\{\left((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\Big)\right]$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\Big)$$

$$+\left((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}\right)(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\Big)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^2 + \left((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^2 + \left((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)}\{\left((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\Big)\right]$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\Big)$$

$$\left((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^2 + \left((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 9:

Nonperturbative Effects Vis-à-vis Quantum Confinement And Lattice Gauge Field Theories, 3D SU(N) Georgi-Glashow Model Vis-à-vis Magnetic Monopoles Confining Strings, Quark Confinement Vis-à-vis String And Field Theories At Finite Temperature: A Reinforcing Template-Dissipation Denosjour Model

1. magnetoacoustically manipulated nematic ordered states Vis-à-vis Quantum memory effects
2. Ebert: k-string tensions Vis-à-vis the 4-d SU(N)-inspired dual Abelian-Higgs-type theory
3. . Nonperturbative effects Vis-à-vis Quantum confinement and lattice gauge field theories
4. 3D SU(N) Georgi-Glashow model Vis-à-vis magnetic monopoles confining strings
5. quark confinement Vis-à-vis string and field theories at finite temperature
6. Wilson and Polyakov loops Vis-à-vis string effective action,
7. String breaking in QCD(dual superconductor)**Vis-à-vis** stochastic vacuum model
8. Confinement in the Abelian-Higgs-type theories **Vis a Vis** String picture and field correlators.
9. Field correlators **Vis-à-vis** string representation

NOTATION

Module One

Magnetoacoustically manipulated nematic ordered states Vis-à-vis Quantum memory effects

G_{13} : Category one of magnetoacoustically manipulated nematic ordered states

G_{14} : Category two of magnetoacoustically manipulated nematic ordered states

G_{15} : Category three of magnetoacoustically manipulated nematic ordered states

T_{13} : Category one of Quantum memory effects

T_{14} : Category two of Quantum memory effects

T_{15} : Category three of Quantum memory effects

Module Two

Ebert: k-string tensions Vis-à-vis the 4-d SU(N)-inspired dual Abelian-Higgs-type theory(classification based on parametricization and characterization of the systems to which they are applicable)

G_{16} : Category one of Ebert: k-string tensions

G_{17} : Category two of Ebert: k-string tensions

G_{18} : Category three of Ebert: k-string tensions

T_{16} : Category one of the 4-d SU(N)-inspired dual Abelian-Higgs-type theory(classification based on parametricization and characterization of the systems to which they are applicable)

T_{17} : Category two of the 4-d SU(N)-inspired dual Abelian-Higgs-type theory(classification based on parametricization and characterization of the systems to which they are applicable)

T_{18} : Category three of the 4-d SU(N)-inspired dual Abelian-Higgs-type theory(classification based on parametricization and characterization of the systems to which they are applicable)

Module three

Nonperturbative effects Vis a Vis Quantum confinement and lattice gauge field theories

G_{20} : Category one of **Nonperturbative effects**

G_{21} : Category two of **Nonperturbative effects**

G_{22} : Category three of **Nonperturbative effects**

T_{20} : Category one of **Quantum confinement and lattice gauge field theories**

T_{21} : Category two of **Quantum confinement and lattice gauge field theories**

T_{22} : Category three of **Quantum confinement and lattice gauge field theories**

Module four

3D SU(N) Georgi-Glashow model(systemic classification based on characteristics' of the systems which are studied) Vis-à-vis magnetic monopoles confining strings

G_{24} : Category one of**3D SU(N) Georgi-Glashow model(systemic classification based on characteristics' of the systems which are studied)**

G_{25} : Category two of **3D SU(N) Georgi-Glashow model(systemic classification based on characteristics' of the systems which are studied)**

G_{26} : Category three of **3D SU(N) Georgi-Glashow model(systemic classification based on characteristics' of the systems which are studied)**

T_{24} : Category one of **magnetic monopoles confining strings**

T_{25} : Category two of **magnetic monopoles confining strings**

T_{26} : Category three of **magnetic monopoles confining strings**

Module five

Quark confinement Vis-à-vis string and field theories at finite temperature

G_{28} : Category one of **quark confinement**

G_{29} : Category two of **quark confinement**

G_{30} : Category three of **quark confinement**

T_{28} : Category one of **string and field theories at finite temperature**

T_{29} : Category two of **string and field theories at finite temperature**

T_{30} : Category three of **string and field theories at finite temperature**

Module six

Wilson and Polyakov loops Vis-à-vis string effective action

G_{32} : Category one of Wilson and Polyakov loops

G_{33} : Category two of Wilson and Polyakov loops

G_{34} : Category three of Wilson and Polyakov loops

T_{32} : Category one of string effective action

T_{33} : Category two of string effective action

T_{34} : Category three of string effective action

Module seven

String breaking in QCD (dual superconductor)**Vis-à-vis** stochastic vacuum model

G_{36} : Category one of String breaking in QCD dual superconductor

G_{37} : Category two of String breaking in QCD dual superconductor

G_{38} : Category three of String breaking in QCD dual superconductor

T_{36} : Category one of stochastic vacuum model

T_{37} : Category two of stochastic vacuum model

T_{38} : Category three of stochastic vacuum model

Module eight

Confinement in the Abelian-Higgs-type theories Vis-à-vis String picture and field correlators

G_{40} : Category one of Confinement in the Abelian-Higgs-type theories

G_{41} : Category two of Confinement in the Abelian-Higgs-type theories

G_{42} : Category three of Confinement in the Abelian-Higgs-type theories

T_{40} : Category one of String picture and field correlators

T_{41} : Category two of String picture and field correlators

T_{42} : Category three of String picture and field correlators

Module Nine

Field correlators **Vis-à-vis** string representation

G_{44} : Category one of Field correlators

G_{45} : Category two of Field correlators

G_{46} : Category three of Field correlators

T_{44} : Category one of string representation

T_{45} : Category two of string representation

T_{46} : Category three of string representation

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}), t\big)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}), t\big)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\big((G_{47}), t\big)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\big((G_{47}), t\big)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\big((G_{47}), t\big) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14}, t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17}, t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25}, t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29}, t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14}, t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17}, t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25}, t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29}, t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14}, t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17}, t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25}, t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29}, t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14}, t)}$, $\boxed{(a''_{14})^{(1)}(T_{14}, t)}$, $\boxed{(a''_{15})^{(1)}(T_{14}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$,$-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$,$-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$,$-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & \boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & \boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & \boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & \boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{array}\right] T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & \boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{array}\right] T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & \boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{array}\right] T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for

category 1, 2 and 3

$-(b_{13}'')^{(1,1,)}(G,t)$, $-(b_{14}'')^{(1,1,)}(G,t)$, $-(b_{15}'')^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b_{20}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b_{24}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b_{28}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b_{32}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \tag{67}$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a_{20}')^{(3)}+(a_{20}'')^{(3)}(T_{21},t) & +(a_{16}'')^{(2,2,2)}(T_{17},t) & +(a_{13}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a_{21}')^{(3)}+(a_{21}'')^{(3)}(T_{21},t) & +(a_{17}'')^{(2,2,2)}(T_{17},t) & +(a_{14}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a_{22}')^{(3)}+(a_{22}'')^{(3)}(T_{21},t) & +(a_{18}'')^{(2,2,2)}(T_{17},t) & +(a_{15}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{22} \tag{69}$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

$Where\ \boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{matrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{matrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{matrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{30} \quad 81$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1,2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1,2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)} & -(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)} & -(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)} & -(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition

coefficients for category 1, 2 and 3
$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$
are seventh detrition coefficients for category 1, 2 and 3
$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3
$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth

augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\left[\begin{array}{c} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)}\ \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\left[\begin{array}{c} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, \ (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\big((G_{35}), t\big) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} \ (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = \ (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - \ T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{\left(\hat{M}_{20}\right)^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{\left(\hat{M}_{20}\right)^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)} + G_{29}^0) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)} + G_{33}^0) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)}\left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)}\left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t)\leq G_{44}^{0}+\int_{0}^{t}\left[(a_{44})^{(9)}\left(G_{45}^{0}+(\hat{P}_{44})^{(9)}e^{(\hat{M}_{44})^{(9)}s_{(44)}}\right)\right]ds_{(44)}=$$

$$\left(1+(a_{44})^{(9)}t\right)G_{45}^{0}+\frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)}t}-1\right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)},T^{(1)}),(G^{(2)},T^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{13})^{(1)}|G_{14}^{(1)}-G_{14}^{(2)}|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}ds_{(13)}+$$

$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad (186)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$

$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. (189)

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\bar{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\bar{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\bar{M}_{16})^{(2)}s_{(16)}}e^{(\bar{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\bar{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\bar{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 216

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})-(a''_{24})^{(4)}(T_{25}^{(2)},s_{(24)})|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)})d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right)$, $t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)})\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32{,}33{,}34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})}$: $\left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$
$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right) \quad (259)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t} \; and \; (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ (261)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: (262)

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If $G_{37} \; or \; G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36} \; is$ bounded, from below, the same property holds for $G_{37} \; and \; G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. (263)

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. (264)

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t), t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can 274

postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose
$(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)}+\left((\hat{P}_{44})^{(9)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\;e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\,(G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(9)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a_{45}')^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a_{46}')^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t),t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a_{13}')^{(1)} + (a_{14}')^{(1)} - (a_{13}'')^{(1)}(T_{14},t) + (a_{14}'')^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b_{13}')^{(1)} + (b_{14}')^{(1)} - (b_{13}'')^{(1)}(G,t) - (b_{14}'')^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)},\ if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}\ , if\ (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$

and $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)},\ if\ (\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$

$\left(\frac{(a_{15})^{(1)}G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le\right.$ 286

$\left.\frac{(a_{15})^{(1)}G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$

$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$ 287

$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$ 288

$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le$ 289

$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16}\mathrm{e}^{\left((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)}\right)\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$
$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)},\ \boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)},\ \boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)},$ and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right) \quad 324$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big) - (b''_{25})^{(4)}\big((G_{27}),t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)},, (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, $\boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$ 331

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24}e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G^0_{24}e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G^0_{24}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G^0_{26}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T^0_{24}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T^0_{26}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$

Definition of $(\nu_1)^{(5)},(\nu_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},\nu^{(5)},u^{(5)}$: 339

By $(\nu_1)^{(5)}>0$, $(\nu_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_1)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)}>0$, $(\bar{\nu}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2+(\sigma_2)^{(5)}\nu^{(5)}-(a_{28})^{(5)}=0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(\nu_0)^{(5)},(m_1)^{(5)}=(\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)}<(\nu_1)^{(5)}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)}<(\nu_0)^{(5)}<(\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(\nu_1)^{(5)},(m_1)^{(5)}=(\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)}<(\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)}$, ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)}=(u_1)^{(5)},(\mu_1)^{(5)}=(\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)}<(u_0)^{(5)}<(\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$
and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$
and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right) \quad 366$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)},, (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^{0}}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^{0}e^{-(S_2)^{(9)}t}\leq G_{46}(t)\leq \frac{(a_{46})^{(9)}G_{44}^{0}}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}\left[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}\right]+G_{46}^{0}e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)\leq\frac{d\nu^{(2)}}{dt}\leq-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)},(\nu_0)^{(2)}$:-

For $0<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\nu_1)^{(2)}<(\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t)\geq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}\quad,\quad\boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\bar{\nu}_1)^{(2)}$ 392

If $0<(\nu_1)^{(2)}<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)}\leq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}\leq\nu^{(2)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\bar{\nu}_1)^{(2)}$$

If $0<(\nu_1)^{(2)}\leq(\bar{\nu}_1)^{(2)}\leq(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)}\leq\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)}\leq\nu^{(2)}(t)\leq(m_1)^{(2)},\quad\boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)}\leq u^{(2)}(t)\leq(\mu_1)^{(2)},\quad\boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)}=(a_{17}'')^{(2)},then\ (\sigma_1)^{(2)}=(\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)}=(\bar{\nu}_1)^{(2)}$ if in addition

$(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(v_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(v_2)^{(5)}}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get 409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{v}_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (\bar{v}_1)^{(5)}$$

If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33},t)\right) - (a_{33}'')^{(6)}(T_{33},t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \le \frac{d\nu^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\quad \boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(i) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T_{25}^{*} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^{*} for which $f(T_{25}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)} G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)} G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

Definition and uniqueness of T_{29}^{*} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^{*} for which $f(T_{29}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)} G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)} G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - \left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \qquad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \tag{545}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \tag{546}$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\big((b_{29}')^{(5)} - (r_{29})^{(5)}\big)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\big(s_{(29)(j)}T_{29}^*\mathbb{G}_j\big) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}\ and\ (b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \ , \ \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}\ and\ (b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 10:

Dual Abelian-Higgs-Type Theories Vis-à-vis Aharonov-Bohm Effect, Luescher Term In The Quark-Antiquark Potential Vis-à-vis Nonperturbative Effects, Mixed Heavy-Quark-Gluon(Nonperturbative Effects) Bose Einstein Condensate Vis-à-vis In The Stochastic Vacuum Model , Et Al, :A Exogenous Endomorph-Arrondissement Arrogate Model

1. Quark confinement Vis-à-vis stochastic vacuum model of QCD(systemic classification based on characteristics')
2. dual Abelian-Higgs-type theories (systemic classification based on characteristics') Vis-à-vis Aharonov-Bohm effect
3. Luescher term in the quark-antiquark potential Vis-à-vis Nonperturbative effects
4. Mixed heavy-quark-gluon(Nonperturbative effects) Bose Einstein condensate Vis-à-vis in the stochastic vacuum model
5. gauge-invariant two-point correlation function Vis-à-vis of gluonic field strengths
6. Heavy-quark condensate at zero and finite temperatures Vis-à-vis Quantum chromodynamics
7. heavy-quark condensate Vis-à-vis world-line formalism
8. Abelian-type theories with confinement Vis-à-vis finite-temperature effects in gauge theories
9. Photostimulated luminescence Vis-à-vis Thermal neutron imaging

NOTATION

Module One

Quark confinement Vis-à-vis stochastic vacuum model of QCD(systemic classification based on

characteristics

G_{13} : Category one of Quark confinement

G_{14} : Category two of Quark confinement

G_{15} : Category three of Quark confinement

T_{13} : Category one of stochastic vacuum model of QCD(systemic classification based on characteristics')

T_{14} : Category two of stochastic vacuum model of QCD(systemic classification based on characteristics')

T_{15} : Category three of stochastic vacuum model of QCD(systemic classification based on characteristics')

dual Abelian-Higgs-type theories (systemic classification based on characteristics') Vis-à-vis Aharonov-Bohm effect

Module Two

G_{16} : Category one of dual Abelian-Higgs-type theories (systemic classification based on characteristics'

G_{17} : Category two of dual Abelian-Higgs-type theories (systemic classification based on characteristics'

G_{18} : Category three of dual Abelian-Higgs-type theories (systemic classification based on characteristics'

T_{16} : Category one of Aharonov-Bohm effect

T_{17} : Category two of Aharonov-Bohm effect

T_{18} : Category three of Aharonov-Bohm effect

Module three

Luescher term in the quark-antiquark potential Vis-à-vis Nonperturbative effects

G_{20} : Category one of Luescher term in the quark-antiquark potential

G_{21} : Category two of Luescher term in the quark-antiquark potential

G_{22} : Category three of Luescher term in the quark-antiquark potential

T_{20} : Category one of Nonperturbative effects

T_{21} : Category two of Nonperturbative effects

T_{22} : Category three of Nonperturbative effects

Module four

Mixed heavy-quark-gluon(Nonperturbative effects) Bose Einstein condensate (it can have different properties) Vis-à-vis in the stochastic vacuum model

G_{24} : Category one of Mixed heavy-quark-gluon(Nonperturbative effects) Bose Einstein condensate

G_{25} : Category two of Mixed heavy-quark-gluon(Nonperturbative effects) Bose Einstein condensate

G_{26} : Category three of Mixed heavy-quark-gluon(Nonperturbative effects) Bose Einstein condensate

T_{24} : Category one of stochastic vacuum model

T_{25} : Category two of stochastic vacuum model

T_{26} : Category three of stochastic vacuum model

Module five

gauge-invariant two-point correlation function Vis-à-vis gluonic field strengths

G_{28} : Category one of gauge-invariant two-point correlation function

G_{29} : Category two of gauge-invariant two-point correlation function

G_{30} : Category three of gauge-invariant two-point correlation function

T_{28} : Category one of gluonic field strengths

T_{29} : Category two of gluonic field strengths

T_{30} : Category three of gluonic field strengths

Module six

Heavy-quark condensate at zero and finite temperatures Vis-à-vis Quantum chromodynamics

Temperature Dependence of Gluon and Quark Condensates as from Linear Confinement: A. N. Mitra, W-Y. P. Hwang

The gluon and quark condensates and their temperature dependence are investigated within QCD premises. The input for the former is a gauge invariant gg kernel **made up of** the direct (D), exchange (X) and contact(C) **QCD interactions** in the lowest order, but with the perturbative propagator k^{-2} replaced by a `non-perturbative k^{-4} form obtained via two differentiations: $ \mu^2 \partial_m^2 (m^2+k^2)^{-1}$, ($\mu$ a scale parameter), and then setting $m=0$, to simulate linear confinement. Similarly for the input $q{\bar q}$ kernel the gluon propagator is replaced by the above k^{-4} form. With these `linear' simulations, the respective condensates **are obtained by** `looping' up the gluon and quark lines in the standard manner. Using Dimensional regularization (DR), the necessary integrals yield the condensates plus temperature corrections, with a common scale parameter μ for both. For gluons the exact result is $$ = {36\mu^4}\pi^ {-3}\alpha_s (\mu^2)[2-\gamma - 4\pi^2 T^2/(3\mu^2)]$$. Evaluation of the quark condensate is preceded by an approximate solution of the SDE for the mass function $m (p) $, giving a recursive formula, with convergence **achieved at** the third iteration. Setting the scale parameter μ equal to the universal Regge slope $1 GeV^2$, the gluon and quark condensates at T=0 are found to be $0.586 Gev^4$ and $(240-260 MeV)^3$ respectively, in fair accord with QCD sum rule values. Next, the temperature corrections (of order $-T^2$ for both condensates) **is determined via** finite-temperature field theory a la Matsubara. Keywords: Gluon Condensate, mass tensor, gauge invariance, linear confinement, finite-temperature, contour-closing. PACS: 11.15.Tk; 12.38.Lg; 13.20.Cz

G_{32} : Category one of Heavy-quark condensate at zero and finite temperatures

G_{33} : Category two of Heavy-quark condensate at zero and finite temperatures

G_{34} : Category three of Heavy-quark condensate at zero and finite temperatures

T_{32} : Category one of Quantum chromodynamics

T_{33} : Category two of Quantum chromodynamics

T_{34} : Category three of Quantum chromodynamics

heavy-quark condensate Vis-à-vis world-line formalism

Module seven

G_{36} : Category one of heavy-quark condensate

G_{37} : Category two of heavy-quark condensate

G_{38} : Category three of heavy-quark condensate

T_{36} : Category one of world-line formalism

T_{37} : Category two of world-line formalism

T_{38} : Category three of world-line formalism

Module eight

Abelian-type theories with confinement Vis-à-vis finite-temperature effects in gauge theories

G_{40} : Category one of Abelian-type theories with confinement

G_{41} : Category two of Abelian-type theories with confinement

G_{42} : Category three of Abelian-type theories with confinement

T_{40} : Category one of finite-temperature effects in gauge theories

T_{41} : Category two of finite-temperature effects in gauge theories

T_{42} : Category three of finite-temperature effects in gauge theories

Module Nine

Photostimulated luminescence Vis-à-vis Thermal neutron imaging

G_{44} : Category one of Photostimulated luminescence

G_{45} : Category two of Photostimulated luminescence

G_{46} : Category three of Photostimulated luminescence

T_{44} : Category one of Thermal neutron imaging

T_{45} : Category two of Thermal neutron imaging

T_{46} : Category three of Thermal neutron imaging

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \big[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\big]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \big[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\big]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \big[(b'_{32})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big)\big]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \big[(b'_{33})^{(6)} - (b''_{33})^{(6)}\big((G_{35}),t\big)\big]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \big[(b'_{34})^{(6)} - (b''_{34})^{(6)}\big((G_{35}),t\big)\big]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \big[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\big]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \big[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\big]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \big[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\big]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \big[(b'_{36})^{(7)} - (b''_{36})^{(7)}\big((G_{39}),t\big)\big]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \big[(b'_{37})^{(7)} - (b''_{37})^{(7)}\big((G_{39}),t\big)\big]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \big[(b'_{38})^{(7)} - (b''_{38})^{(7)}\big((G_{39}),t\big)\big]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \big[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\big]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \big[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\big]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \big[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\big]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \big[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}),t\big)\big]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \big[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}),t\big)\big]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \big[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}),t\big)\big]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \big[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\big]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{ccc} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a_{20}')^{(3)} & +(a_{20}'')^{(3)}(T_{21},t) & +(a_{16}'')^{(2,2,2)}(T_{17},t) & +(a_{13}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a_{21}')^{(3)} & +(a_{21}'')^{(3)}(T_{21},t) & +(a_{17}'')^{(2,2,2)}(T_{17},t) & +(a_{14}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a_{22}')^{(3)} & +(a_{22}'')^{(3)}(T_{21},t) & +(a_{18}'')^{(2,2,2)}(T_{17},t) & +(a_{15}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b_{20}')^{(3)} & -(b_{20}'')^{(3)}(G_{23},t) & -(b_{16}'')^{(2,2,2)}(G_{19},t) & -(b_{13}'')^{(1,1,1,)}(G,t) \\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & \boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & \boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \qquad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation*

$coefficient\ \ for\ category\ 1,2\ and\ 3$
$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ $are\ third\ \ augmentation$
$coefficient\ for\ category\ 1,2\ and\ 3$
$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where$ $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients$
$for\ category\ 1,2\ and\ 3$
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients$
$for\ category\ 1,2\ and\ 3$
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients$
$for\ category\ 1,2\ and\ 3$
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ are third augmentation

$coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34}$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2, and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b_{46}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)}\ \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)}\ \boxed{+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)}\ \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)}\ \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $(a_{36}'')^{(7)}(T_{37},t)$, $(a_{37}'')^{(7)}(T_{37},t)$, $(a_{38}'')^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{44})^{(9)} & +(a''_{44})^{(9)}(T_{45},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\begin{bmatrix} (a'_{45})^{(9)} & +(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{l} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{l} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)}\ \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[\begin{array}{l} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)}\ \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[\begin{array}{l} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first

augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \tag{127}$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \tag{128}$$

$$\lim_{G \to \infty} (b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \tag{131}$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

$$\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$$

$$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and $(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}$, $(r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \quad (136)$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}\big((G_{43}),t\big) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\big((G_{43}),t\big)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} \text{ , } T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} \text{ , } T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$$ 169

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t\in\mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t\in\mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \le \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} |(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1T_{14}$ which leads to

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1t})+T_{14}^0e^{-\varepsilon_1t}$ If we take t such that $e^{-\varepsilon_1t}=\ \frac{1}{2}$ it results

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad (237)$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (239)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: (240)

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. (241)

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, $if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right. \quad 324$$
$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((s_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le-(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le-(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(j) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]}$, $G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^* \mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^*\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^*\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^*\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}$, $\frac{\partial(b_i'')^{(7)}}{\partial G_j}(\,(G_{39})^{*^*}\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^*\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^*\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^*\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\Big)\right]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\Big)\right]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\Big)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 11:

Radiative corrections to scalar fermion pair production Vis-à-vis high energy e+ e- collisions, Radiative corrections Vis-à-vis Supersymmetry, Higgs decays in the two Higgs doublet model: Large quantum effects Vis-à-vis the decoupling regime: A Dorcas gazelle-Weeny Weaver Model

1. Determination of the density of defect states Vis-à-vis thermally stimulated conductivity studied from numerical simulations
2. CP-odd A0 production at e+e-colliders Vis-à-vis the minimal supersymmetric standard model with SP violating phases.
3. Fourth generation CP phase Vis-à-vis b s transitions.
4. Radiative corrections to scalar fermion pair production Vis-à-vis high energy e+ e- collisions
5. Radiative corrections Vis-à-vis Supersymmetry.
6. Higgs decays in the two Higgs doublet model: Large quantum effects Vis-à-vis the decoupling regime
7. d-wave to p-wave pairing Vis-à-vis the t'-t Hubbard model at zero temperature
8. Dynamical Cluster Approximation Projective Quantum Monte Carlo method Vis-à-vis Hubbard model triplet superconductivity
9. Orbital-selective Mott-Hubbard transition Vis-à-vis the two-band Hubbard model

NOTATION

Module One

Density of defect states Vis-à-vis thermally stimulated conductivity studied from numerical simulations

G_{13} : Category one of density of defect states

G_{14} : Category two of density of defect states

G_{15} : Category three of density of defect states

T_{13} : Category one of thermally stimulated conductivity studied from numerical simulations

T_{14} : Category two of thermally stimulated conductivity studied from numerical simulations

T_{15} : Category three of thermally stimulated conductivity studied from numerical simulations

Module Two

CP-odd A0 production at e+e-colliders Vis-à-vis the minimal supersymmetric standard model with SP

violating phases

G_{16} : Category one of CP-odd A0 production at e+e-colliders

G_{17} : Category two of CP-odd A0 production at e+e-colliders

G_{18} : Category three of CP-odd A0 production at e+e-colliders

T_{16} : Category one of the minimal supersymmetric standard model with SP violating phases

T_{17} : Category two of the minimal supersymmetric standard model with SP violating phases

T_{18} : Category three of the minimal supersymmetric standard model with SP violating phases

Fourth generation CP phase Vis-à-vis b s transitions

Module three

G_{20} : Category one of Fourth generation CP phase

G_{21} : Category two of Fourth generation CP phase

G_{22} : Category three of Fourth generation CP phase

T_{20} : Category one of b s transitions

T_{21} : Category two of b s transitions

T_{22} : Category three of b s transitions

Module four

Radiative corrections to scalar fermion pair production Vis-à-vis high energy e+ e- collisions

G_{24} : Category one of Radiative corrections to scalar fermion pair production

G_{25} : Category two of Radiative corrections to scalar fermion pair production

G_{26} : Category three of Radiative corrections to scalar fermion pair production

T_{24} : Category one of energy e+ e- collisions

T_{25} : Category two of energy e+ e- collisions

T_{26} : Category three of energy e+ e- collisions

Radiative corrections Vis-à-vis Supersymmetry

Module five

G_{28} : Category one of Radiative corrections

G_{29} : Category two of Radiative corrections

G_{30} : Category three of Radiative corrections

T_{28} : Category one of Supersymmetry

T_{29} : Category two of Supersymmetry

T_{30} : Category three of Supersymmetry

Module six

Higgs decays in the two Higgs doublet model: Large quantum effects Vis-à-vis the decoupling regime

G_{32} : Category one of Higgs decays in the two Higgs doublet model: Large quantum effects

G_{33} : Category two of Higgs decays in the two Higgs doublet model: Large quantum effects

G_{34} : Category three of Higgs decays in the two Higgs doublet model: Large quantum effects

T_{32} : Category one of the decoupling regime

T_{33} : Category two of the decoupling regime

T_{34} : Category three of the decoupling regime

Module seven

D-wave to p-wave pairing Vis avis the t'-t Hubbard model at zero temperature(classification based on the characteristics' of the systems to which the model is applicable)

G_{36} : Category one of d-wave to p-wave pairing

G_{37} : Category two of d-wave to p-wave pairing

G_{38} : Category three of d-wave to p-wave pairing

T_{36} : Category one of the t'-t Hubbard model at zero temperature(classification based on the characteristics' of the systems to which the model is applicable)

T_{37} : Category two of the t'-t Hubbard model at zero temperature(classification based on the characteristics' of the systems to which the model is applicable)

T_{38} : Category three of the t'-t Hubbard model at zero temperature(classification based on the characteristics' of the systems to which the model is applicable)

Module eight

Dynamical Cluster Approximation Projective Quantum Monte Carlo method Vis-à-vis Hubbard model triplet superconductivity

G_{40} : Category one of Dynamical Cluster Approximation Projective Quantum Monte Carlo method

G_{41} : Category two of Dynamical Cluster Approximation Projective Quantum Monte Carlo method

G_{42} : Category three of Dynamical Cluster Approximation Projective Quantum Monte Carlo method

T_{40} : Category one of Hubbard model triplet superconductivity

T_{41} : Category two of Hubbard model triplet superconductivity

T_{42} : Category three of Hubbard model triplet superconductivity

Module Nine

Orbital-selective Mott-Hubbard transition Vis-à-vis the two-band Hubbard model

G_{44} : Category one of Orbital-selective Mott-Hubbard transition

G_{45} : Category two of Orbital-selective Mott-Hubbard transition

G_{46} : Category three of Orbital-selective Mott-Hubbard transition

T_{44} : Category one of the two-band Hubbard model

T_{45} : Category two of the two-band Hubbard model

T_{46} : Category three of the two-band Hubbard model

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}), t\big)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}), t\big)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\big((G_{47}), t\big)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\big((G_{47}), t\big)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\big((G_{47}), t\big) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14}, t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17}, t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25}, t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29}, t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14}, t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17}, t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25}, t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29}, t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14}, t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17}, t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25}, t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29}, t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45}, t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14}, t)}$, $\boxed{(a''_{14})^{(1)}(T_{14}, t)}$, $\boxed{(a''_{15})^{(1)}(T_{14}, t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$,$+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$,$-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$,$-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$,$-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & \boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & \boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & \boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} \quad \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & \boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{array}\right] T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & \boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{array}\right] T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & \boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} \quad \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{array}\right] T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for

category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \tag{67}$$

$$(a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \tag{69}$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

$Where\ \boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{matrix}(a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{matrix}(a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{matrix}(a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{30} \quad 81$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1,2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1,2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1,2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition

coefficients for category 1, 2 and 3
$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$
are seventh detrition coefficients for category 1, 2 and 3
$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3
$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3
$+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth

augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)} & \boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)} & \boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, \ (r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(r_i)^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)}$ 113

$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\big((G_{35}), t\big) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} \ (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = \ (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \tag{136}$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\big((G_{43}), t\big) = (r_i)^{(8)} \tag{141}$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \tag{146}$$

A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le \left(\hat{P}_{20}\right)^{(3)} e^{\left(\hat{M}_{20}\right)^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le \left(\hat{Q}_{20}\right)^{(3)} e^{\left(\hat{M}_{20}\right)^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{\left(\hat{M}_{24}\right)^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{\left(\hat{M}_{28}\right)^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{\left(\hat{M}_{32}\right)^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{\left(\hat{M}_{36}\right)^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:
Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy
$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right) \quad 169$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)} + G_{29}^0) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)} + G_{33}^0) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)}\left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)}\left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t)\leq G_{44}^{0}+\int_{0}^{t}\left[(a_{44})^{(9)}\left(G_{45}^{0}+(\hat{P}_{44})^{(9)}e^{(\hat{M}_{44})^{(9)}s_{(44)}}\right)\right]ds_{(44)}=$$

$$\left(1+(a_{44})^{(9)}t\right)G_{45}^{0}+\frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)}t}-1\right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left((G^{(1)},T^{(1)}),(G^{(2)},T^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{13})^{(1)}|G_{14}^{(1)}-G_{14}^{(2)}|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}ds_{(13)}+$$

$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$

$G_{13}^{(2)}|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$

$T_i\,(t)\geq T_i^0e^{\left(-(b_i')^{(1)}t\right)}>0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 216

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_1 / (a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)} \left((\widehat{M}_{20})^{(3)}\right)_2 / (a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}(T_{25}^{(1)},s_{(24)})-(a''_{24})^{(4)}(T_{25}^{(2)},s_{(24)})|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)})d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $((\widehat{M}_{24})^{(4)})_1,((\widehat{M}_{24})^{(4)})_2$ *and* $((\widehat{M}_{24})^{(4)})_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{32})^{(6)} \left|G_{33}^{(1)} - G_{33}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\, ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} \left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} +$$

$$G_{32}^{(2)} \left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)} t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)})\right) \quad (248)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32{,}33{,}34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})}$: $\left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$(a''_{36})^{(7)}\left(T^{(1)}_{37}, s_{(36)}\right)\left|G^{(1)}_{36} - G^{(2)}_{36}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$
$G^{(2)}_{36}\left|(a''_{36})^{(7)}\left(T^{(1)}_{37}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T^{(2)}_{37}, s_{(36)}\right)\right| \; e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})\right) \quad 259$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t} \; and \; (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0 \; and \; T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b'_i)^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2 \; and \; \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37} \; and \; G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G^0_{37} + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G^0_{38} + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If $G_{37} \; or \; G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36} \; is$ bounded, from below, the same property holds for $G_{37} \; and \; G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b''_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t), t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can 274

postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose
$(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)}+\left((\hat{P}_{44})^{(9)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{44})^{(9)}\right]\leq(\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\;e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\;(G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$

From 99 to 44 it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(9)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a_{45}')^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a_{46}')^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t),t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a_{13}')^{(1)} + (a_{14}')^{(1)} - (a_{13}'')^{(1)}(T_{14},t) + (a_{14}'')^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b_{13}')^{(1)} + (b_{14}')^{(1)} - (b_{13}'')^{(1)}(G,t) - (b_{14}'')^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)},,(\bar{v}_2)^{(1)},(\bar{u}_1)^{(1)},(\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)},(\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}$, $(m_2)^{(1)}$, $(\mu_1)^{(1)},(\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}\ , if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(\ m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}\ , if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\ \mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$

$\left(\frac{(a_{15})^{(1)}G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le\right.$ 286

$\left.\frac{(a_{15})^{(1)}G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$

$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$ 287

$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$ 288

$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le$ 289

$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16}\mathrm{e}^{\left((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)}\right)\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$
$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)},\ \boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)},\ \boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)},$ and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)},\ \boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big) - (b''_{25})^{(4)}\big((G_{27}),t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)},, (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, $\boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$ 331

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24}e^{(S_1)^{(4)}t} \quad 332$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G^0_{24}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G^0_{24}e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G^0_{24}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G^0_{26}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T^0_{24}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T^0_{24}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T^0_{24}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T^0_{26}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)}=(a_{24})^{(4)}(m_2)^{(4)}-(a'_{24})^{(4)}$

$(S_2)^{(4)}=(a_{26})^{(4)}-(p_{26})^{(4)}$

$(R_1)^{(4)}=(b_{24})^{(4)}(\mu_2)^{(4)}-(b'_{24})^{(4)}$

$(R_2)^{(4)}=(b'_{26})^{(4)}-(r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$:

$(\sigma_1)^{(5)},(\sigma_2)^{(5)},(\tau_1)^{(5)},(\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)}\le-(a'_{28})^{(5)}+(a'_{29})^{(5)}-(a''_{28})^{(5)}(T_{29},t)+(a''_{29})^{(5)}(T_{29},t)\le-(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)}\le-(b'_{28})^{(5)}+(b'_{29})^{(5)}-(b''_{28})^{(5)}\big((G_{31}),t\big)-(b''_{29})^{(5)}\big((G_{31}),t\big)\le-(\tau_1)^{(5)}$

Definition of $(v_1)^{(5)},(v_2)^{(5)},(u_1)^{(5)},(u_2)^{(5)},v^{(5)},u^{(5)}$: 339

By $(v_1)^{(5)}>0$, $(v_2)^{(5)}<0$ and respectively $(u_1)^{(5)}>0$, $(u_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2+(\sigma_1)^{(5)}v^{(5)}-(a_{28})^{(5)}=0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_1)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$ and

Definition of $(\bar{v}_1)^{(5)},,(\bar{v}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)}>0$, $(\bar{v}_2)^{(5)}<0$ and respectively $(\bar{u}_1)^{(5)}>0$, $(\bar{u}_2)^{(5)}<0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2+(\sigma_2)^{(5)}v^{(5)}-(a_{28})^{(5)}=0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2+(\tau_2)^{(5)}u^{(5)}-(b_{28})^{(5)}=0$

Definition of $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)},(v_0)^{(5)}$:-

If we define $(m_1)^{(5)},(m_2)^{(5)},(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)}=(v_0)^{(5)},(m_1)^{(5)}=(v_1)^{(5)},$ ***if*** $(v_0)^{(5)}<(v_1)^{(5)}$

$(m_2)^{(5)}=(v_1)^{(5)},(m_1)^{(5)}=(\bar{v}_1)^{(5)}$, ***if*** $(v_1)^{(5)}<(v_0)^{(5)}<(\bar{v}_1)^{(5)}$,

and $\boxed{(v_0)^{(5)}=\frac{G_{28}^{0}}{G_{29}^{0}}}$

$(m_2)^{(5)}=(v_1)^{(5)},(m_1)^{(5)}=(v_0)^{(5)},$ ***if*** $(\bar{v}_1)^{(5)}<(v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)}=(u_0)^{(5)},(\mu_1)^{(5)}=(u_1)^{(5)},$ ***if*** $(u_0)^{(5)}<(u_1)^{(5)}$

$(\mu_2)^{(5)}=(u_1)^{(5)},(\mu_1)^{(5)}=(\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)}<(u_0)^{(5)}<(\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$
and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$
and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^{0}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G_{36}^{0}e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G_{36}^{0}e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right) \quad 366$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)},, (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}(\nu^{(8)})^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, ***if*** $(\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, ***if*** $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, ***if*** $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44} e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G^0_{44} e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G^0_{44} e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^{0}}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^{0}e^{-(S_2)^{(9)}t}\leq G_{46}(t)\leq \frac{(a_{46})^{(9)}G_{44}^{0}}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}\left[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}\right]+G_{46}^{0}e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^{0}e^{(R_1)^{(9)}t}\leq T_{44}(t)\leq \frac{1}{(\mu_2)^{(9)}}T_{44}^{0}e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}(\nu^{(1)})^{2}+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \le \nu^{(1)}(t) \le (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \le u^{(1)}(t) \le (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)\leq\frac{d\nu^{(2)}}{dt}\leq-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)},(\nu_0)^{(2)}$:-

For $0<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\nu_1)^{(2)}<(\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t)\geq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}\quad,\quad\boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)}\leq\nu^{(2)}(t)\leq(\bar{\nu}_1)^{(2)}$ 392

If $0<(\nu_1)^{(2)}<(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}<(\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)}\leq\frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}\leq\nu^{(2)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\bar{\nu}_1)^{(2)}$$

If $0<(\nu_1)^{(2)}\leq(\bar{\nu}_1)^{(2)}\leq(\nu_0)^{(2)}=\frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)}\leq\nu^{(2)}(t)\leq\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}\leq(\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)}\leq\nu^{(2)}(t)\leq(m_1)^{(2)},\quad\boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)}\leq u^{(2)}(t)\leq(\mu_1)^{(2)},\quad\boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)}=(a_{17}'')^{(2)},then\ (\sigma_1)^{(2)}=(\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)}=(\bar{\nu}_1)^{(2)}$ if in addition

$(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}$, $\boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}$, $\boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(\nu_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{\nu}_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right) - (a''_{33})^{(6)}(T_{33},t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)} e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \le \frac{d\nu^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $v^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t) v^{(8)} - (a_{41})^{(8)} v^{(8)}$$

Definition of $v^{(8)}$:- $\quad \boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)} v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)} v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq v^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

423

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

424

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following

Theorem : If $(a_i'')^{(1)} and\ (b_i'')^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(2)} and\ (b_i'')^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(k) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T_{25}^{*} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^{*} for which $f(T_{25}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)} G_{25}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)} G_{25}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

Definition and uniqueness of T_{29}^{*} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^{*} for which $f(T_{29}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)} G_{29}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)} G_{29}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

Definition and uniqueness of T_{33}^{*} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^{*} for which $f(T_{33}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

Definition and uniqueness of T_{37}^{*} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^{*} for which $f(T_{37}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

Definition and uniqueness of T_{41}^{*} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^{*} for which $f(T_{41}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

Definition and uniqueness of T_{45}^{*} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^{*} for which $f(T_{45}^{*}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - \left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \qquad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a_{16}')^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a_{17}')^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a_{18}')^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b_{16}')^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T_{16}^*\mathbb{G}_j\right) \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b_{17}')^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T_{17}^*\mathbb{G}_j\right) \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b_{18}')^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T_{18}^*\mathbb{G}_j\right) \tag{539}$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \tag{544}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \tag{545}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \tag{546}$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \ , \ \frac{\partial (b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\big((a_{24}')^{(4)} + (p_{24})^{(4)}\big)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\big((a_{25}')^{(4)} + (p_{25})^{(4)}\big)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\big((a_{26}')^{(4)} + (p_{26})^{(4)}\big)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\big((b_{24}')^{(4)} - (r_{24})^{(4)}\big)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(24)(j)}T_{24}^*\mathbb{G}_j\big) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\big((b_{25}')^{(4)} - (r_{25})^{(4)}\big)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\big(s_{(25)(j)}T_{25}^*\mathbb{G}_j\big) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\big((b_{26}')^{(4)} - (r_{26})^{(4)}\big)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\big(s_{(26)(j)}T_{26}^*\mathbb{G}_j\big) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\big((a_{28}')^{(5)} + (p_{28})^{(5)}\big)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\big((a_{29}')^{(5)} + (p_{29})^{(5)}\big)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\big((a_{30}')^{(5)} + (p_{30})^{(5)}\big)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\big((b_{28}')^{(5)} - (r_{28})^{(5)}\big)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\big(s_{(28)(j)}T_{28}^*\mathbb{G}_j\big) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\big((b_{29}')^{(5)} - (r_{29})^{(5)}\big)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\big(s_{(29)(j)}T_{29}^*\mathbb{G}_j\big) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}\ and\ (b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \ , \ \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\ (G_{35})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}\ and\ (b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left(\ (G_{39})^{**}\ \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{41})^{(8)}}{\partial T_{41}}(T^*_{41}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a'_{40})^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G^*_{40}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a'_{41})^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G^*_{41}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a'_{42})^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G^*_{42}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\ B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad (586C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad (586D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad (586E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad (586F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad (586G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^*_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^*_{21}+(b_{21})^{(3)}s_{(20),(22)}T^*_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^*_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^*_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^*_{25}+(b_{25})^{(4)}s_{(24),(25)}T^*_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^*_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^*_{25}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^*_{25}+(b_{25})^{(4)}s_{(24),(24)}T^*_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^2+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left(\left((\lambda)^{(4)}\right)^2+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\Big(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\Big(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\Big(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\Big(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\Big(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\Big(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 12:

Transport Theory Vis-à-vis Quantum Tunneling, Heisenberg Model Vis-à-vis Exchange Interactions (Electron); Ground States Vis-à-vis Spin Fluctuations, Mesoscopic Systems Vis-à-vis Green's Function Methods And Band Structure Et Al,: Coextensive Representations And Apriori Determination Thereof:

1. Orbital selective Mott transition Dynamical means field theory Vis-à-vis Projective Quantum Monte Carlo method Hubbard model
2. Phonon extinction rules Vis-à-vis thermal diffuse scattering.
3. Transport theory Vis-à-vis Quantum Tunneling.
4. Heisenberg model Vis-à-vis exchange interactions (electron);
5. Ground states Vis-à-vis spin fluctuations.
6. Mesoscopic systems Vis-à-vis Green's function methods and band structure.
7. U-J synergy effect Vis-à-vis high Tc superconductors.
8. High Tc superconductivity Vis-à-vis - Extended Hubbard model
9. Fluctuation-dissipation relation of a dissipative driven mesoscopic system. Vis-à-vis Time dependent Transport

NOTATION

Module One

Orbital selective Mott transition Dynamical means field theory Vis-à-vis Projective Quantum Monte

Carlo method Hubbard model

G_{13} : Category one of Orbital selective Mott transition Dynamical means field theory

G_{14} : Category two of Orbital selective Mott transition Dynamical means field theory

G_{15} : Category three of Orbital selective Mott transition Dynamical means field theory

T_{13} : Category one of Projective Quantum Monte Carlo method Hubbard model

T_{14} : Category two of Projective Quantum Monte Carlo method Hubbard model

T_{15} : Category three of Projective Quantum Monte Carlo method Hubbard model

Module Two

Phonon extinction rules Vis-à-vis thermal diffuse scattering

G_{16} : Category one of Phonon extinction rules

G_{17} : Category two of Phonon extinction rules

G_{18} : Category three of Phonon extinction rules

T_{16} : Category one of thermal diffuse scattering

T_{17} : Category two of thermal diffuse scattering

T_{18} : Category three of thermal diffuse scattering

Module three

Transport theory Vis-à-vis Quantum Tunneling

G_{20} : Category one of Transport theory

G_{21} : Category two of Transport theory

G_{22} : Category three of Transport theory

T_{20} : Category one of Quantum Tunneling

T_{21} : Category two of Quantum Tunneling

T_{22} : Category three of Quantum Tunneling

Module four

Heisenberg model(Classification is done based on the systems to which the theory could be applied and the concomitant and corresponding characterization thereof) Vis-à-vis exchange interactions (electron);

G_{24} : Category one of Heisenberg model(Classification is done based on the systems to which the theory could be applied and the concomitant and corresponding characterization thereof)

G_{25} : Category two of Heisenberg model(Classification is done based on the systems to which the theory could be applied and the concomitant and corresponding characterization thereof)

G_{26} : Category three of Heisenberg model(Classification is done based on the systems to which the theory could be applied and the concomitant and corresponding characterization thereof)

T_{24} : Category one of exchange interactions (electron);

T_{25} : Category two of exchange interactions (electron);

T_{26} : Category three of exchange interactions (electron);

Module five

Ground states Vis-à-vis spin fluctuations

G_{28} : Category one of Ground states

G_{29} : Category two of Ground states

G_{30} : Category three of Ground states

T_{28} : Category one of spin fluctuations

T_{29} : Category two of spin fluctuations

T_{30} : Category three of spin fluctuations

Module six

Mesoscopic systems Vis-à-vis Green's function methods and band structure.

G_{32} : Category one of Mesoscopic systems

G_{33} : Category two of Mesoscopic systems

G_{34} : Category three of Mesoscopic systems

T_{32} : Category one of Green's function methods and band structure

T_{33} : Category two of Green's function methods and band structure

T_{34} : Category three of Green's function methods and band structure

Module seven

U-J synergy effect Vis-à-vis high Tc superconductors

G_{36} : Category one of U-J synergy effect

G_{37} : Category two of U-J synergy effect

G_{38} : Category three of U-J synergy effect

T_{36} : Category one of high Tc superconductors

T_{37} : Category two of high Tc superconductors

T_{38} : Category three of high Tc superconductors

Module eight

High Tc superconductivity Vis-à-vis - Extended Hubbard model

G_{40} : Category one of High Tc superconductivity

G_{41} : Category two of High Tc superconductivity

G_{42} : Category three of High Tc superconductivity

T_{40} : Category one of- Extended Hubbard model

T_{41} : Category two of- Extended Hubbard model

T_{42} : Category three of- Extended Hubbard model

Module Nine

Fluctuation-dissipation relation of a dissipative driven mesoscopic system. Vis-à-vis Time dependent Transport

G_{44} : Category one of Fluctuation-dissipation relation of a dissipative driven mesoscopic system

G_{45} : Category two of Fluctuation-dissipation relation of a dissipative driven mesoscopic system

G_{46} : Category three of Fluctuation-dissipation relation of a dissipative driven mesoscopic system

T_{44} : Category one of Time dependent Transport

T_{45} : Category two of Time dependent Transport

T_{46} : Category three of Time dependent Transport

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad (1)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad (2)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad (3)$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad (4)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad (5)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad (6)$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad (7)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad (8)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad (9)$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad (10)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad (11)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad (12)$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad (13)$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad (14)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad (15)$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad (16)$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad (17)$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad (18)$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}), t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}), t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}), t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,)}(G_{19},t)$, $-(b_{17}'')^{(2,2,)}(G_{19},t)$, $-(b_{18}'')^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)}\,+(a_{16}'')^{(2)}(T_{17},t)\;\;+(a_{13}'')^{(1,1,)}(T_{14},t)\;\;+(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)\;\;+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)\;\;+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t)\;\;+(a_{40}'')^{(8,8,8)}(T_{41},t)\;\;+(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right]G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)}\,+(a_{17}'')^{(2)}(T_{17},t)\;\;+(a_{14}'')^{(1,1,)}(T_{14},t)\;\;+(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)\;\;+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)\;\;+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t)\;\;+(a_{41}'')^{(8,8,8)}(T_{41},t)\;\;+(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right]G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)}\,+(a_{18}'')^{(2)}(T_{17},t)\;\;+(a_{15}'')^{(1,1,)}(T_{14},t)\;\;+(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)\;\;+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)\;\;+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t)\;\;+(a_{42}'')^{(8,8,8)}(T_{41},t)\;\;+(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right]G_{18} \quad (63)$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient

for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients

for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition

coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \left[\begin{array}{c} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)}\ \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \left[\begin{array}{c} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)}\ \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{array}{c} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)}\ \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}\ \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation

coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $-(b_{36}'')^{(7)}(G_{39},t)$, $-(b_{37}'')^{(7)}(G_{39},t)$, $-(b_{38}'')^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)} & +(a_{44}'')^{(9)}(T_{45},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)} & +(a_{45}'')^{(9)}(T_{45},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)} & +(a_{46}'')^{(9)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{15}$$

Where $+(a_{44}'')^{(9)}(T_{45},t)$, $+(a_{45}'')^{(9)}(T_{45},t)$, $+(a_{46}'')^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third

augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth
augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth
augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth
augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh
augmentation coefficient for category 1, 2 and 3
$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth
augmentation coefficient for 1,2,3
$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth
augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$
Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth

detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$lim_{T_2 \to \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)}$ 113

$\lim_{G \to \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,((G_{27}),t) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$

$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}((G_{27}),t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$

$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$

$(b_i'')^{(5)}((G_{31}),t) \le \ (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$

$lim_{T_2\to\infty}(a_i'')^{(5)}\,(T_{29},t) = (p_i)^{(5)}$ 123

$\lim_{G\to\infty}(b_i'')^{(5)}\,(G_{31},t) = \ (r_i)^{(5)}$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\,(a_i)^{(5)} + (a_i')^{(5)} + \ (\hat{A}_{28})^{(5)} + \ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\,(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty}(b_i'')^{(6)}\,\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\big((G_{35}),t\big)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}},\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}<1$$

Definition of $(\hat{P}_{32})^{(6)},(\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)},(\hat{A}_{32})^{(6)}and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)},(a_i')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(p_i)^{(6)},\ (r_i)^{(6)},i=32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)}+(a_i')^{(6)}+(\hat{A}_{32})^{(6)}+(\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}]<1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)}+(b_i')^{(6)}+(\hat{B}_{32})^{(6)}+(\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}]<1$$

Where we suppose

$$(a_i)^{(7)},(a_i')^{(7)},(a_i'')^{(7)},(b_i)^{(7)},(b_i')^{(7)},(b_i'')^{(7)}>0,\qquad i,j=36,37,38 \quad 131$$

The functions $(a_i'')^{(7)},(b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)},\ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t)\le(p_i)^{(7)}\le(\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t)\le\ (r_i)^{(7)}\le(b_i')^{(7)}\le(\hat{B}_{36})^{(7)} \quad 132$$

$$\lim_{T_2\to\infty}\ (a_i'')^{(7)}(T_{37},t)=(p_i)^{(7)}$$

$$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big)=\ (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)},(\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)},(\hat{B}_{36})^{(7)},(p_i)^{(7)},\ (r_i)^{(7)}}$ are positive constants and $\boxed{i=36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t)-(a_i'')^{(7)}(T_{37},t)|\le(\hat{k}_{36})^{(7)}|T_{37}-T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \ , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \le \ (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty}(a_i'')^{(8)}\,(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty}(b_i'')^{(8)}\,\big((G_{43}), t\big) = \ (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \le (\hat{k}_{40})^{(8)}|T_{41} - \ T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}} \ , \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, \ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
$G_i, T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''^{(9)}_{44}\left(T_{45}(s_{(44)}), s_{(44)}\right)\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for G_{14} , $G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for G_{17} , $G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)} + G_{33}^0\right) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)}\left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)}\left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad (184)$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0e^{(-(b_i')^{(1)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{16})^{(2)} \left|G_{17}^{(1)} - G_{17}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}}\, ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} \left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} \left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}}\} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| \mathrm{e}^{-(\widehat{M}_{16})^{(2)} \mathrm{t}} \leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right) \mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} \mathrm{e}^{(\widehat{M}_{16})^{(2)} \mathrm{t}}$ and $(\widehat{Q}_{16})^{(2)} \mathrm{e}^{(\widehat{M}_{16})^{(2)} \mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where G_i (t) $= 0$ and T_i (t) $= 0$ 200

it results

$$G_i \text{ (t)} \geq G_i^0 \mathrm{e}^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} \mathrm{d}s_{(16)}\right]} \geq 0$$

$$T_i \text{ (t)} \geq T_i^0 \mathrm{e}^{(-(b_i')^{(2)} \mathrm{t})} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{\mathrm{d}G_{17}}{\mathrm{dt}} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a'_{17})^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then 203

$T_{17} \to \infty$.

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad (208)$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$:$\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{20})^{(3)}\left|G_{21}^{(1)} - G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}\left|G_{20}^{(1)} - G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)} - G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a_{20}'')^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad (214)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (216)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. (218)

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. (219)

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to (220)

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is

unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b''_{22})^{(3)}\left((G_{23})(t),t\right)=(b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}<1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\widehat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{24})^{(4)}\right]\leq(\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 227

$(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24}\ \ is$ bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}), (\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}), (\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{28})^{(5)}\left|G_{29}^{(1)} - G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and

respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\hat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{32})^{(6)}\right]\leq(\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\hat{A}_{36})^{(7)}+(\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\,(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\hat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36{,}37{,}38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 261

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(7)}t)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38}. indeed if

$G_{36}<(\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt}\leq\left((\widehat{M}_{36})^{(7)}\right)_1-(a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{40})^{(8)} \left|G_{41}^{(1)}-G_{41}^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{(\widehat{M}_{40})^{(8)} s_{(40)}}\, ds_{(40)} +$$
$$\int_0^t \{(a'_{40})^{(8)} \left|G_{40}^{(1)}-G_{40}^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{(\widehat{M}_{40})^{(8)} s_{(40)}} +$$
$$G_{40}^{(2)} \left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{(\widehat{M}_{40})^{(8)} s_{(40)}}\} ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} t} \leq \qquad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right) d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ *and* $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\big((G_{43})(t), t\big) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})}$: $\left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a_{44}')^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a_{44}'')^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a_{44}'')^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a_{44}'')^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a_{44}')^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ and $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(9)}$ and $(b''_i)^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b'_i)^{(9)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ and $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} and G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} or G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} is bounded, from below, the same property holds for G_{45} and G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(9)}\left((G_{47})(t), t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t), t\right) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$\left(\frac{(a_{15})^{(1)} G^0_{13}}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G^0_{15} e^{-(S_2)^{(1)} t} \le G_{15}(t) \le\right.$ 286

$$\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t})$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)}=(\nu_0)^{(2)},(m_1)^{(2)}=(\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)}<(\nu_1)^{(2)}$ 305

$(m_2)^{(2)}=(\nu_1)^{(2)},(m_1)^{(2)}=(\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)}<(\nu_0)^{(2)}<(\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right.$ 312
$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right.$ 324

$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously 331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^{0}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^{0}e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G_{24}^{0}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G_{24}^{0}e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^{0}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t}-e^{-(S_2)^{(4)}t}\right]+G_{26}^{0}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^{0}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^{0}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T_{24}^{0}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T_{24}^{0}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^{0}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t}-e^{-(b'_{26})^{(4)}t}\right]+T_{26}^{0}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \ \boldsymbol{if}\ (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \boldsymbol{if}\ (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and $\boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T^0_{32}}{T^0_{33}}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G^0_{32}e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}}G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}}G^0_{32}e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)}G^0_{32}}{(m_1)^{(6)}((S_1)^{(6)}-(p_{32})^{(6)}-(S_2)^{(6)})}\left[e^{((S_1)^{(6)}-(p_{32})^{(6)})t}-e^{-(S_2)^{(6)}t}\right]+G^0_{34}e^{-(S_2)^{(6)}t}\le G_{34}(t)\le\right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T^0_{32}e^{(R_1)^{(6)}t}\le T_{32}(t)\le T^0_{32}e^{((R_1)^{(6)}+(r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T^0_{32}e^{(R_1)^{(6)}t}\le T_{32}(t)\le\frac{1}{(\mu_2)^{(6)}}T^0_{32}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T^0_{32}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T^0_{34}e^{-(b'_{34})^{(6)}t}\le T_{34}(t)\le \quad 358$$

$$\frac{(a_{34})^{(6)}T^0_{32}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T^0_{34}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$ 360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$-(\sigma_2)^{(7)}\le-(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t)\le-(\sigma_1)^{(7)}$

$-(\tau_2)^{(7)}\le-(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big)\le-(\tau_1)^{(7)}$

Definition of $(\nu_1)^{(7)},(\nu_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},\nu^{(7)},u^{(7)}$: 361

By $(\nu_1)^{(7)}>0$, $(\nu_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0$, $(u_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_1)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{\nu}_1)^{(7)},,(\bar{\nu}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)}>0$, $(\bar{\nu}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0$, $(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_2)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(\nu_0)^{(7)},(m_1)^{(7)}=(\nu_1)^{(7)}$, ***if*** $(\nu_0)^{(7)}<(\nu_1)^{(7)}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\bar{\nu}_1)^{(7)}$, ***if*** $(\nu_1)^{(7)}<(\nu_0)^{(7)}<(\bar{\nu}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$
$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, $\boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40}e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G^0_{40}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G^0_{40}e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G^0_{40}e^{(S_1)^{(8)}t} \quad 376$$

$$\Big(\frac{(a_{42})^{(8)}G^0_{40}}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\Big[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\Big] + G^0_{42}e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \quad 377$$
$$\frac{(a_{42})^{(8)}G^0_{40}}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}] + G^0_{42}e^{-(a'_{42})^{(8)}t}\Big)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^{0}e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^{0}}{G_{45}^{0}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\le\frac{dv^{(1)}}{dt}\le-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\ge\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\le v^{(1)}(t)\le(\nu_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t)\le\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\le v^{(1)}(t)\le(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\le\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\le v^{(1)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\le(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\le(\bar{\nu}_1)^{(1)}\le\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\le v^{(1)}(t)\le\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\le(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\le v^{(1)}(t)\le(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\le u^{(1)}(t)\le(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)}=(a_{14}'')^{(1)},then\ (\sigma_1)^{(1)}=(\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)}=(\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)}=(\nu_1)^{(1)}$ then $v^{(1)}(t)=(\nu_0)^{(1)}$ and as a consequence $G_{13}(t)=(\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \le \frac{dv^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \ge \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}}$$

it follows $(v_0)^{(2)} \le v^{(2)}(t) \le (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \le \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}}$$

From which we deduce $(v_0)^{(2)} \le v^{(2)}(t) \le (\bar{v}_1)^{(2)}$ 392

If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 393

$$(v_1)^{(2)} \le \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)} - (v_2)^{(2)}\right)t\right]}} \le v^{(2)}(t) \le$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \le (\bar{v}_1)^{(2)}$$

If $0 < (v_1)^{(2)} \le (\bar{v}_1)^{(2)} \le (v_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(v_1)^{(2)} \le v^{(2)}(t) \le \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}\right)t\right]}} \le (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le v^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)\right) - (a_{25}'')^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29},t)\right) - (a_{29}'')^{(5)}(T_{29},t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(v^{(5)}\right)^2+(\sigma_2)^{(5)}v^{(5)}-(a_{28})^{(5)}\right)\leq\frac{dv^{(5)}}{dt}\leq-\left((a_{29})^{(5)}\left(v^{(5)}\right)^2+(\sigma_1)^{(5)}v^{(5)}-(a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)},(v_0)^{(5)}$:-

For $0<\boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}<(v_1)^{(5)}<(\bar{v}_1)^{(5)}$

$$v^{(5)}(t)\geq\frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}\quad,\quad\boxed{(C)^{(5)}=\frac{(v_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(v_2)^{(5)}}}$$

it follows $(v_0)^{(5)}\leq v^{(5)}(t)\leq(v_1)^{(5)}$

In the same manner , we get 409

$$v^{(5)}(t)\leq\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(5)}=\frac{(\bar{v}_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)}\leq v^{(5)}(t)\leq(\bar{v}_5)^{(5)}$

If $0<(v_1)^{(5)}<(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}<(\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)}\leq\frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}\leq v^{(5)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}\leq(\bar{v}_1)^{(5)}$$

If $0<(v_1)^{(5)}\leq(\bar{v}_1)^{(5)}\leq\boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)}\leq v^{(5)}(t)\leq\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}\leq(v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)}\leq v^{(5)}(t)\leq(m_1)^{(5)},\quad\boxed{v^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)}\leq u^{(5)}(t)\leq(\mu_1)^{(5)},\quad\boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)}=(a_{29}'')^{(5)},then\ (\sigma_1)^{(5)}=(\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)}=(\bar{v}_1)^{(5)}$ if in addition

$(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \le \frac{dv^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

For $0 < \boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \ge \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \le v^{(6)}(t) \le (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \le \nu^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \le u^{(7)}(t) \le (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \le \frac{d\nu^{(8)}}{dt} \le -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right) - (a''_{45})^{(9)}(T_{45},t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \le \frac{dv^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$v^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le v^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le v^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le v^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$, we obtain

$$(\nu_1)^{(9)} \le v^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)} \le v^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0$ 477

$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0$ 478

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(l) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three

first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0\,, T^*_{14}$ given by $f(T^*_{14}) = 0$ and

$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]}$, $G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$

$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}$, $T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \tag{513}$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \tag{521}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}\left((G_{19})^*\right) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \ , \ \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^{*}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^{*}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)} G_{44}^* \mathbb{T}_{45} \qquad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)} G_{45}^* \mathbb{T}_{45} \qquad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)} G_{46}^* \mathbb{T}_{45} \qquad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)} T_{44}^* \mathbb{G}_j\right) \qquad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)} T_{45}^* \mathbb{G}_j\right) \qquad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)} T_{46}^* \mathbb{G}_j\right) \qquad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)} G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)} G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)} G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)} G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^*\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b_{18}')^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a_{18}')^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\Big]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 13:

Quantum Monte Carlo Method Vis-à-vis Molecular Nanodevices, Anderson-Holstein Model Vis-à-vis Molecular Transistor, Green-Function Approach Vis-à-vis Transport Phenomena In Quantum Pumps: A Crotchet-Conundrum Model

1. Quantum Monte Carlo method Vis-à-vis molecular nanodevices
2. Anderson-Holstein model Vis-à-vis molecular transistor.
3. Green-function approach Vis-à-vis transport phenomena in quantum pumps.
4. Time-dependent quantum transport Vis-à-vis Keldysh formalism
5. Short optical pulses, Vis-à-vis Autocorrelation
6. FEL Efficiency Increase Vis-à-vis Magnetic Field Tapering of the Undulators
7. Cerenkov Free Electron Lasers and tapering and treatment of saturation

8. Free electron laser oscillators with tapered undulators Vis-à-vis Inclusion of harmonic generation and pulse propagation
9. The nonlinear stability Vis-à-vis asymptotically anti-de Sitter solutions

NOTATION

Module One

Quantum Monte Carlo method Vis-à-vis molecular nanodevices

G_{13} : Category one of Quantum Monte Carlo method

G_{14} : Category two of Quantum Monte Carlo method

G_{15} : Category three of Quantum Monte Carlo method

T_{13} : Category one of molecular nanodevices

T_{14} : Category two of molecular nanodevices

T_{15} : Category three of molecular nanodevices

Module Two

Anderson-Holstein model Vis-à-vis molecular transistor

G_{16} : Category one of Anderson-Holstein model

G_{17} : Category two of Anderson-Holstein model

G_{18} : Category three of Anderson-Holstein model

T_{16} : Category one of molecular transistor

T_{17} : Category two of molecular transistor

T_{18} : Category three of molecular transistor

Module three

Green-function approach Vis-à-vis transport phenomena in quantum pumps

G_{20} : Category one of Green-function approach

G_{21} : Category two of Green-function approach

G_{22} : Category three of Green-function approach

T_{20} : Category one of transport phenomena in quantum pumps

T_{21} : Category two of transport phenomena in quantum pumps

T_{22} : Category three of transport phenomena in quantum pumps

Module four

Time-dependent quantum transport Vis-à-vis Keldysh formalism

G_{24} : Category one of Time-dependent quantum transport

G_{25} : Category two of Time-dependent quantum transport

G_{26} : Category three of Time-dependent quantum transport

T_{24} : Category one of Keldysh formalism

T_{25} : Category two of Keldysh formalism

T_{26} : Category three of Keldysh formalism

Module five

Short optical pulses, Vis-à-vis Autocorrelation

G_{28} : Category one of Short optical pulses

G_{29} : Category two of Short optical pulses

G_{30} : Category three of Short optical pulses

T_{28} : Category one of Autocorrelation

T_{29} : Category two of Autocorrelation

T_{30} : Category three of Autocorrelation

FEL Efficiency Increase Vis-à-vis Magnetic Fields Tapering of the Undulators

Module six

G_{32} : Category one of FEL Efficiency Increase

G_{33} : Category two of FEL Efficiency Increase

G_{34} : Category three of FEL Efficiency Increase

T_{32} : Category one of Magnetic Fields Tapering of the Undulators

T_{33} : Category two of Magnetic Fields Tapering of the Undulators

T_{34} : Category three of Magnetic Fields Tapering of the Undulators

Cerenkov Free Electron Lasers and tapering and treatment of saturation

Module seven

G_{36} : Category one of Cerenkov Free Electron Lasers

G_{37} : Category two of Cerenkov Free Electron Lasers

G_{38} : Category three of Cerenkov Free Electron Lasers

T_{36} : Category one of tapering and treatment of saturation

T_{37} : Category two of tapering and treatment of saturation

T_{38} : Category three of tapering and treatment of saturation

Module eight

Free electron laser oscillators with tapered undulators Vis-à-vis Inclusion of harmonic generation and pulse propagation

G_{40} : Category one of Free electron laser oscillators with tapered undulators

G_{41} : Category two of Free electron laser oscillators with tapered undulators

G_{42} : Category three of Free electron laser oscillators with tapered undulators

T_{40} : Category one of Inclusion of harmonic generation and pulse propagation

T_{41} : Category two of Inclusion of harmonic generation and pulse propagation

T_{42} : Category three of Inclusion of harmonic generation and pulse propagation

Module Nine

Nonlinear stability Vis-à-vis asymptotically anti-de Sitter solutions

G_{44} : Category one of nonlinear stability

G_{45} : Category two of nonlinear stability

G_{46} : Category three of nonlinear stability

T_{44} : Category one of asymptotically anti-de Sitter solutions

T_{45} : Category two of asymptotically anti-de Sitter solutions

T_{46} : Category three of asymptotically anti-de Sitter solutions

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}), t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}), t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}), t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}), t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,)}(G_{19},t)$, $-(b_{17}'')^{(2,2,)}(G_{19},t)$, $-(b_{18}'')^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,)}(G_{23},t)$, $-(b_{21}'')^{(3,3,)}(G_{23},t)$, $-(b_{22}'')^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,)}(G_{39},t)$, $-(b_{36}'')^{(7,7,)}(G_{39},t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8)}(G_{43},t)$ $-(b_{41}'')^{(8,8)}(G_{43},t)$ $-(b_{42}'')^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{c} (a_{16}')^{(2)}\,+(a_{16}'')^{(2)}(T_{17},t)\;\;+(a_{13}'')^{(1,1,)}(T_{14},t)\;\;+(a_{20}'')^{(3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)\;\;+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)\;\;+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{36}'')^{(7,7,7)}(T_{37},t)\;\;+(a_{40}'')^{(8,8,8)}(T_{41},t)\;\;+(a_{44}'')^{(9,9)}(T_{45},t) \end{array}\right]G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{c} (a_{17}')^{(2)}\,+(a_{17}'')^{(2)}(T_{17},t)\;\;+(a_{14}'')^{(1,1,)}(T_{14},t)\;\;+(a_{21}'')^{(3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)\;\;+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)\;\;+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{37}'')^{(7,7,7)}(T_{37},t)\;\;+(a_{41}'')^{(8,8,8)}(T_{41},t)\;\;+(a_{45}'')^{(9,9)}(T_{45},t) \end{array}\right]G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{c} (a_{18}')^{(2)}\,+(a_{18}'')^{(2)}(T_{17},t)\;\;+(a_{15}'')^{(1,1,)}(T_{14},t)\;\;+(a_{22}'')^{(3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)\;\;+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)\;\;+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t) \\ +(a_{38}'')^{(7,7,7)}(T_{37},t)\;\;+(a_{42}'')^{(8,8,8)}(T_{41},t)\;\;+(a_{46}'')^{(9,9)}(T_{45},t) \end{array}\right]G_{18} \quad (63)$$

Where $+(a_{16}'')^{(2)}(T_{17},t)$, $+(a_{17}'')^{(2)}(T_{17},t)$, $+(a_{18}'')^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient

for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients

for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* $1, 2$ *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category $1, 2$ *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition

coefficients for category 1,2, and 3

$-(b_{20}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b_{42}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b_{46}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a_{32}')^{(6)} & +(a_{32}'')^{(6)}(T_{33},t) & +(a_{28}'')^{(5,5,5)}(T_{29},t) & +(a_{24}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a_{33}')^{(6)} & +(a_{33}'')^{(6)}(T_{33},t) & +(a_{29}'')^{(5,5,5)}(T_{29},t) & +(a_{25}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a_{34}')^{(6)} & +(a_{34}'')^{(6)}(T_{33},t) & +(a_{30}'')^{(5,5,5)}(T_{29},t) & +(a_{26}'')^{(4,4,4,)}(T_{25},t) \\ +(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a_{32}'')^{(6)}(T_{33},t)$, $+(a_{33}'')^{(6)}(T_{33},t)$, $+(a_{34}'')^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a_{28}'')^{(5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a_{24}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a_{13}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a_{16}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a_{20}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a_{36}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a_{40}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a_{44}'')^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation

coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37}$$

$$- \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36}$$

$$- \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37}$$

$$- \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40}$$

$$- \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $-(b_{36}'')^{(7)}(G_{39},t)$, $-(b_{37}'')^{(7)}(G_{39},t)$, $-(b_{38}'')^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)$, $-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{38}'')^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)} & +(a_{44}'')^{(9)}(T_{45},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)} & +(a_{45}'')^{(9)}(T_{45},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)} & +(a_{46}'')^{(9)}(T_{37},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{15}$$

Where $+(a_{44}'')^{(9)}(T_{45},t)$, $+(a_{45}'')^{(9)}(T_{45},t)$, $+(a_{46}'')^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third

augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth
augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth
augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth
augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh
augmentation coefficient for category 1, 2 and 3
$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth
augmentation coefficient for 1,2,3
$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth
augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$
Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3
$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth

detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: (100)

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: (101)

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22$ 112

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$lim_{T_2 \to \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)}$ 113

$\lim_{G \to \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)} t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and $(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$lim_{T_2\to\infty}(a_i'')^{(4)}\,(T_{25},t) = (p_i)^{(4)}$ 118

$\lim_{G\to\infty}(b_i'')^{(4)}\,((G_{27}),t) = \ (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24{,}25{,}26}$

They satisfy Lipschitz condition: 119

$|(a_i'')^{(4)}(T_{25}',t) - (a_i'')^{(4)}(T_{25},t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$

$|(b_i'')^{(4)}((G_{27})',t) - (b_i'')^{(4)}((G_{27}),t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)},$ are positive constants

$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, \ (r_i)^{(4)}, i = 24{,}25{,}26$, satisfy the inequalities

$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)} + (a_i')^{(4)} + \ (\hat{A}_{24})^{(4)} + \ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}] < 1$

$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)} + (b_i')^{(4)} + \ (\hat{B}_{24})^{(4)} + \ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}] < 1$

Where we suppose

$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28{,}29{,}30$ 122

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, \ (r_i)^{(5)}$:

$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$

$(b_i'')^{(5)}((G_{31}),t) \le \ (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$

$lim_{T_2\to\infty}(a_i'')^{(5)}\,(T_{29},t) = (p_i)^{(5)}$ 123

$\lim_{G\to\infty}(b_i'')^{(5)}\,(G_{31},t) = \ (r_i)^{(5)}$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \ , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and \ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, \ (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, \ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq \ (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\big((G_{35}),t\big)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}},\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}<1$$

Definition of $(\hat{P}_{32})^{(6)},(\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)},(\hat{A}_{32})^{(6)}and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)},(a_i')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(p_i)^{(6)},\ (r_i)^{(6)},i=32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)}+(a_i')^{(6)}+(\hat{A}_{32})^{(6)}+(\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}]<1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)}+(b_i')^{(6)}+(\hat{B}_{32})^{(6)}+(\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}]<1$$

Where we suppose

$$(a_i)^{(7)},(a_i')^{(7)},(a_i'')^{(7)},(b_i)^{(7)},(b_i')^{(7)},(b_i'')^{(7)}>0,\qquad i,j=36,37,38 \quad 131$$

The functions $(a_i'')^{(7)},(b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)},\ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t)\le(p_i)^{(7)}\le(\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t)\le\ (r_i)^{(7)}\le(b_i')^{(7)}\le(\hat{B}_{36})^{(7)} \quad 132$$

$$\lim_{T_2\to\infty}(a_i'')^{(7)}(T_{37},t)=(p_i)^{(7)}$$

$$\lim_{G\to\infty}(b_i'')^{(7)}\big((G_{39}),t\big)=\ (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)},(\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)},(\hat{B}_{36})^{(7)},(p_i)^{(7)},\ (r_i)^{(7)}}$ are positive constants and $\boxed{i=36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t)-(a_i'')^{(7)}(T_{37},t)|\le(\hat{k}_{36})^{(7)}|T_{37}-T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\ , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \le \ (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty}(a_i'')^{(8)}\,(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty}(b_i'')^{(8)}\,\big((G_{43}), t\big) = \ (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \le (\hat{k}_{40})^{(8)}|T_{41} - \ T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}} \ , \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, \ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} t} , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{13})^{(1)}$, $T_i^0 \leq (\hat{Q}_{13})^{(1)}$, 155

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 156

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$ 157

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{20})^{(3)}, T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{24})^{(4)}, T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{28})^{(5)}, T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166 A

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''^{(9)}_{44}\left(T_{45}(s_{(44)}), s_{(44)}\right)\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for G_{14} , $G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for G_{17} , $G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)}\left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}}\right)\right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)} t} - 1\right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)} + G_{33}^0\right) e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[((\hat{P}_{40})^{(8)} + G_{41}^0) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[((\hat{P}_{44})^{(9)} + G_{45}^0) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad (184)$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0e^{(-(b_i')^{(1)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\ (G(t),t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\ (G(t),t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}\ and\ (\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{16})^{(2)} \left|G_{17}^{(1)} - G_{17}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}}\, ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} \left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}} +$$

$$G_{16}^{(2)} \left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)} s_{(16)}} e^{(\widehat{M}_{16})^{(2)} s_{(16)}}\} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| \mathrm{e}^{-(\widehat{M}_{16})^{(2)} \mathrm{t}} \leq$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right) \mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} \mathrm{e}^{(\widehat{M}_{16})^{(2)} \mathrm{t}}$ and $(\widehat{Q}_{16})^{(2)} \mathrm{e}^{(\widehat{M}_{16})^{(2)} \mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where G_i (t) $= 0$ and T_i (t) $= 0$ 200

it results

$$G_i \text{ (t)} \geq G_i^0 \mathrm{e}^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} \mathrm{d}s_{(16)}\right]} \geq 0$$

$$T_i \text{ (t)} \geq T_i^0 \mathrm{e}^{(-(b_i')^{(2)} \mathrm{t})} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{\mathrm{d}G_{17}}{\mathrm{dt}} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a'_{17})^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then 203

$T_{17} \to \infty$.

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad (208)$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$:$\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{20})^{(3)}\left|G_{21}^{(1)} - G_{21}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}\left|G_{20}^{(1)} - G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)} - G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}\left|(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a_{20}'')^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad (214)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ (216)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If $G_{20}\ \ is$ bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. (218)

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. (219)

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to (220)

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$$ By taking now ε_3 sufficiently small one sees that T_{21} is

unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b''_{22})^{(3)}\left((G_{23})(t),t\right)=(b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}<1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\widehat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{24})^{(4)}\right]\leq(\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 227

$(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a_{26}')^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24}\ \ is$ bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{28})^{(5)}\left|G_{29}^{(1)} - G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$\left|(G_{31})^{(1)} - (G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and

respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\hat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{32})^{(6)}\right]\leq(\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\hat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\;e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\hat{A}_{36})^{(7)}+(\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\;(G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\hat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36{,}37{,}38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 261

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(7)}t)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38}. indeed if

$G_{36}<(\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt}\leq\left((\widehat{M}_{36})^{(7)}\right)_1-(a'_{37})^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\frac{2}{\varepsilon_7}$$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{40})^{(8)} \left|G_{41}^{(1)}-G_{41}^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{(\widehat{M}_{40})^{(8)} s_{(40)}}\, ds_{(40)} +$$
$$\int_0^t \{(a'_{40})^{(8)} \left|G_{40}^{(1)}-G_{40}^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{(\widehat{M}_{40})^{(8)} s_{(40)}} +$$
$$G_{40}^{(2)} \left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)} s_{(40)}} e^{(\widehat{M}_{40})^{(8)} s_{(40)}}\} ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)} t} \leq \qquad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right) d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ *and* $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)} t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\big((G_{43})(t), t\big) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})}$: $\left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a_{44}')^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a_{44}'')^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a_{44}'')^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a_{44}'')^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a_{44}')^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(9)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0, (\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0, (\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$\left(\frac{(a_{15})^{(1)} G^0_{13}}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G^0_{15} e^{-(S_2)^{(1)} t} \le G_{15}(t) \le\right.$ 286

$$\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}\right]+G_{15}^{0}e^{-(a'_{15})^{(1)}t})$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le \frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le -(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le -(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)}=(\nu_0)^{(2)},(m_1)^{(2)}=(\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)}<(\nu_1)^{(2)}$ 305

$(m_2)^{(2)}=(\nu_1)^{(2)},(m_1)^{(2)}=(\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)}<(\nu_0)^{(2)}<(\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right.$ 312
$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}((G_{23}), t) \le -(\tau_1)^{(3)}$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad (323)$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right) \quad (324)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad (325)$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad (326)$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad (327)$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously 331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^{0}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^{0}e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}}G_{24}^{0}e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}}G_{24}^{0}e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)}G_{24}^{0}}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t}-e^{-(S_2)^{(4)}t}\right]+G_{26}^{0}e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^{0}e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^{0}e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}}T_{24}^{0}e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}}T_{24}^{0}e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)}T_{24}^{0}}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t}-e^{-(b'_{26})^{(4)}t}\right]+T_{26}^{0}e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)}T_{24}^{0}}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t}-e^{-(R_2)^{(4)}t}\right]+T_{26}^{0}e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \ \boldsymbol{if}\ (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \boldsymbol{if}\ (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and $\boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T^0_{32}}{T^0_{33}}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G^0_{32}e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}}G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}}G^0_{32}e^{(S_1)^{(6)}t}$$ 354

$$\left(\frac{(a_{34})^{(6)}G^0_{32}}{(m_1)^{(6)}((S_1)^{(6)}-(p_{32})^{(6)}-(S_2)^{(6)})}\left[e^{((S_1)^{(6)}-(p_{32})^{(6)})t}-e^{-(S_2)^{(6)}t}\right]+G^0_{34}e^{-(S_2)^{(6)}t}\le G_{34}(t)\le\right. \quad 355$$

(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T^0_{32}e^{(R_1)^{(6)}t}\le T_{32}(t)\le T^0_{32}e^{((R_1)^{(6)}+(r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T^0_{32}e^{(R_1)^{(6)}t}\le T_{32}(t)\le\frac{1}{(\mu_2)^{(6)}}T^0_{32}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T^0_{32}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T^0_{34}e^{-(b'_{34})^{(6)}t}\le T_{34}(t)\le \quad 358$$

$$\frac{(a_{34})^{(6)}T^0_{32}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T^0_{34}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$ 360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$-(\sigma_2)^{(7)}\le-(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t)\le-(\sigma_1)^{(7)}$

$-(\tau_2)^{(7)}\le-(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big)\le-(\tau_1)^{(7)}$

Definition of $(\nu_1)^{(7)},(\nu_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},\nu^{(7)},u^{(7)}$: 361

By $(\nu_1)^{(7)}>0$, $(\nu_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0$, $(u_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_1)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{\nu}_1)^{(7)},,(\bar{\nu}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)}>0$, $(\bar{\nu}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0$, $(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_2)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(\nu_0)^{(7)},(m_1)^{(7)}=(\nu_1)^{(7)}$, ***if*** $(\nu_0)^{(7)}<(\nu_1)^{(7)}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\bar{\nu}_1)^{(7)}$, ***if*** $(\nu_1)^{(7)}<(\nu_0)^{(7)}<(\bar{\nu}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$
$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, $\boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,
and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40}e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G^0_{40}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G^0_{40}e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G^0_{40}e^{(S_1)^{(8)}t} \quad 376$$

$$\Big(\frac{(a_{42})^{(8)}G^0_{40}}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\Big[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\Big] + G^0_{42}e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \quad 377$$
$$\frac{(a_{42})^{(8)}G^0_{40}}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}] + G^0_{42}e^{-(a'_{42})^{(8)}t}\Big)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^{0}e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\left(u^{(9)}\right)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G_{44}^{0}}{G_{45}^{0}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)},\ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\le\frac{dv^{(1)}}{dt}\le-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\ge\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\le v^{(1)}(t)\le(\nu_1)^{(1)}$

In the same manner , we get 384

$$v^{(1)}(t)\le\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\le v^{(1)}(t)\le(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\le\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\le v^{(1)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\le(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\le(\bar{\nu}_1)^{(1)}\le\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\le v^{(1)}(t)\le\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\le(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\le v^{(1)}(t)\le(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\le u^{(1)}(t)\le(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)}=(a_{14}'')^{(1)},then\ (\sigma_1)^{(1)}=(\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)}=(\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)}=(\nu_1)^{(1)}$ then $v^{(1)}(t)=(\nu_0)^{(1)}$ and as a consequence $G_{13}(t)=(\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \le \nu^{(2)}(t) \le (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)\right) - (a_{25}'')^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)}-(\nu_0)^{(4)}}{(\nu_0)^{(4)}-(\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29},t)\right) - (a_{29}'')^{(5)}(T_{29},t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(v^{(5)}\right)^2+(\sigma_2)^{(5)}v^{(5)}-(a_{28})^{(5)}\right)\le\frac{dv^{(5)}}{dt}\le-\left((a_{29})^{(5)}\left(v^{(5)}\right)^2+(\sigma_1)^{(5)}v^{(5)}-(a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)},(v_0)^{(5)}$:-

For $0<\boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}<(v_1)^{(5)}<(\bar{v}_1)^{(5)}$

$$v^{(5)}(t)\ge\frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_0)^{(5)}\right)t\right]}}\quad,\quad\boxed{(C)^{(5)}=\frac{(v_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(v_2)^{(5)}}}$$

it follows $(v_0)^{(5)}\le v^{(5)}(t)\le(v_1)^{(5)}$

In the same manner , we get 409

$$v^{(5)}(t)\le\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(5)}=\frac{(\bar{v}_1)^{(5)}-(v_0)^{(5)}}{(v_0)^{(5)}-(\bar{v}_2)^{(5)}}}$$

From which we deduce $(v_0)^{(5)}\le v^{(5)}(t)\le(\bar{v}_5)^{(5)}$

If $0<(v_1)^{(5)}<(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}<(\bar{v}_1)^{(5)}$ we find like in the previous case, 410

$$(v_1)^{(5)}\le\frac{(v_1)^{(5)}+(C)^{(5)}(v_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((v_1)^{(5)}-(v_2)^{(5)}\right)t\right]}}\le v^{(5)}(t)\le$$

$$\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}\le(\bar{v}_1)^{(5)}$$

If $0<(v_1)^{(5)}\le(\bar{v}_1)^{(5)}\le\boxed{(v_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(v_1)^{(5)}\le v^{(5)}(t)\le\frac{(\bar{v}_1)^{(5)}+(\bar{C})^{(5)}(\bar{v}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{v}_1)^{(5)}-(\bar{v}_2)^{(5)}\right)t\right]}}\le(v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)}\le v^{(5)}(t)\le(m_1)^{(5)},\quad\boxed{v^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)}\le u^{(5)}(t)\le(\mu_1)^{(5)},\quad\boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)}=(a_{29}'')^{(5)},then\ (\sigma_1)^{(5)}=(\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)}=(\bar{v}_1)^{(5)}$ if in addition

$(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$ **this also defines** $(v_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \le \frac{dv^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

For $0 < \boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \ge \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \le v^{(6)}(t) \le (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \le v^{(6)}(t) \le (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \le \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)} - (v_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}} \le (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \le (\bar{v}_1)^{(6)} \le \boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(v_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}\right)t\right]}} \le (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \le \nu^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \le u^{(7)}(t) \le (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \le \frac{d\nu^{(8)}}{dt} \le -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(8)}(t)$:-

$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}$, $\boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$
In a completely analogous way, we obtain
Definition of $u^{(8)}(t)$:-

$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}$, $\boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$ if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right) - (a''_{45})^{(9)}(T_{45},t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \le \frac{dv^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$v^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le v^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le v^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le v^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$, we obtain

$$(\nu_1)^{(9)} \le v^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)} \le v^{(9)}(t) \le (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \le u^{(9)}(t) \le (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0$ 477

$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0$ 478

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(m) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three

first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right]+(b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right]+(b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0\,, T^*_{14}$ given by $f(T^*_{14}) = 0$ and

$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]}$, $G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$

$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]}$, $T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and (512)

$$G_{16}^{*} = \frac{(a_{16})^{(2)} G_{17}^{*}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)} G_{17}^{*}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^{*})]} \tag{513}$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)} T_{17}^{*}}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)} T_{17}^{*}}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^{*})]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution (515)

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)} G_{21}^{*}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)} G_{21}^{*}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)} T_{21}^{*}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)} T_{21}^{*}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution (516)

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution (518)

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution (520)

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \tag{521}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}\left((G_{19})^*\right) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \ , \ \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^{*}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^{*}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)} G_{44}^* \mathbb{T}_{45} \qquad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)} G_{45}^* \mathbb{T}_{45} \qquad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)} G_{46}^* \mathbb{T}_{45} \qquad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)} T_{44}^* \mathbb{G}_j\right) \qquad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)} T_{45}^* \mathbb{G}_j\right) \qquad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)} T_{46}^* \mathbb{G}_j\right) \qquad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)} G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)} G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)} G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)} G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^*\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(2)} + (b_{18}')^{(2)} - (r_{18})^{(2)}\right)\{\left((\lambda)^{(2)} + (a_{18}')^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^2+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^2+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^*_{33}+(b_{33})^{(6)}s_{(32),(34)}T^*_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\Big]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^*_{37}+(b_{37})^{(7)}s_{(36),(37)}T^*_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^*_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^*_{37}+(b_{37})^{(7)}s_{(36),(36)}T^*_{36}\Big)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\Big]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 14:

Stability Of Geons, Boson Stars And Black Holes (Asymptotically Nonlinear Solutions Which Are Nonlinearly Stable Vis-à-vis Black Hole Binaries, Wilhelm Weber And The Surface Charges Vis-à-vis Resistive Conductors Carrying Steady Currents Surface: Model With primary Focus On The Homologous Receptiveness Of Differentially Interactional And Instrumental Activity

1. Stability of geons, boson stars and black holes (asymptotically nonlinear solutions which are nonlinearly stable Vis-à-vis Black hole binaries.
2. Wilhelm Weber and the Surface Charges Vis-à-vis Resistive Conductors Carrying Steady Currents surface
3. Electron-beam propagation Vis-à-vis a two-dimensional electron gas.

4. Quantum point contact, ballistic transport, potential fluctuations, and two-dimensional electron gas
5. Quasiballistic transport Vis-à-vis HgTe quantum-well nanostructures
6. Isotopic dilution, self-association Vis-à-Vis Raman non-coincidence in the binary system
7. The black hole Vis-à-vis Global topological k-defects
8. Chiral Symmetry Breaking and Pions Vis-à-vis Nonsupersymmetric Gauge / Gravity Duals
9. Space-time dependent couplings Vis-à-vis n=1 SUSY gauge theories: anomalies and central functions

NOTATION

Module One

Stability of geons, boson stars and black holes (asymptotically nonlinear solutions which are nonlinearly stable) Vis-à-vis Black hole binaries

G_{13} : Category one of Stability of geons, boson stars and black holes (asymptotically nonlinear solutions which are nonlinearly stable)

G_{14} : Category two of Stability of geons, boson stars and black holes (asymptotically nonlinear solutions which are nonlinearly stable)

G_{15} : Category three of Stability of geons, boson stars and black holes (asymptotically nonlinear solutions which are nonlinearly stable)

T_{13} : Category one of Black hole binaries

T_{14} : Category two of Black hole binaries

T_{15} : Category three of Black hole binaries

Module Two

Weber Surface Charges Vis-à-vis Resistive Conductors Carrying Steady Currents surface

G_{16} : Category one of Weber Surface Charges

G_{17} : Category two of Weber Surface Charges

G_{18} : Category three of Weber Surface Charges

T_{16} : Category one of Resistive Conductors Carrying Steady Currents surface

T_{17} : Category two of Resistive Conductors Carrying Steady Currents surface

T_{18} : Category three of Resistive Conductors Carrying Steady Currents surface

Module three

Electron-beam propagation Vis-à-vis two-dimensional electron gases

G_{20} : Category one of Electron-beam propagation

G_{21} : Category two of Electron-beam propagation

G_{22} : Category three of Electron-beam propagation

T_{20} : Category one of two-dimensional electron gases

T_{21} : Category two of two-dimensional electron gases

T_{22} : Category three of two-dimensional electron gases

Module four

Quantum point contact, ballistic transport, potential fluctuations Vis-à-vis two-dimensional electron gas

G_{24} : Category one of Quantum point contact, ballistic transport, potential fluctuations

G_{25} : Category two of Quantum point contact, ballistic transport, potential fluctuations

G_{26} : Category three of Quantum point contact, ballistic transport, potential fluctuations

T_{24} : Category one of two-dimensional electron gas

T_{25} : Category two of two-dimensional electron gas

T_{26} : Category three of two-dimensional electron gas

Quasiballistic transport Vis-à-vis HgTe quantum-well nanostructures

Module five

G_{28} : Category one of Quasiballistic transport

G_{29} : Category two of Quasiballistic transport

G_{30} : Category three of Quasiballistic transport

T_{28} : Category one of HgTe quantum-well nanostructures

T_{29} : Category two of HgTe quantum-well nanostructures

T_{30} : Category three of HgTe quantum-well nanostructures

Module six

Isotopic dilution, self-association Vis-à-Vis Raman non-coincidence in the binary system

G_{32} : Category one of Isotopic dilution, self-association

G_{33} : Category two of Isotopic dilution, self-association

G_{34} : Category three of Isotopic dilution, self-association

T_{32} : Category one of Raman non-coincidence in the binary system

T_{33} : Category two of Raman non-coincidence in the binary system

T_{34} : Category three of Raman non-coincidence in the binary system

Module seven

The black hole Vis-à-vis Global topological k-defects

G_{36} : Category one of The black hole

G_{37} : Category two of The black hole

G_{38} : Category three of The black hole

T_{36} : Category one of Global topological k-defects

T_{37} : Category two of Global topological k-defects

T_{38} : Category three of Global topological k-defects

Module eight

Chiral Symmetry Breaking and Pions Vis-à-vis Nonsupersymmetric Gauge / Gravity Duals

G_{40} : Category one of Chiral Symmetry Breaking and Pions

G_{41} : Category two of Chiral Symmetry Breaking and Pions

G_{42} : Category three of Chiral Symmetry Breaking and Pions

T_{40} : Category one of Nonsupersymmetric Gauge / Gravity Duals

T_{41} : Category two of Nonsupersymmetric Gauge / Gravity Duals

T_{42} : Category three of Nonsupersymmetric Gauge / Gravity Duals

Module Nine

Space-time dependent couplings Vis-à-vis n=1 SUSY gauge theories: anomalies and central functions

G_{44} : Category one of Space-time dependent couplings

G_{45} : Category two of Space-time dependent couplings

G_{46} : Category three of Space-time dependent couplings

T_{44} : Category one of n=1 SUSY gauge theories: anomalies and central functions

T_{45} : Category two of n=1 SUSY gauge theories: anomalies and central functions

T_{46} : Category three of n=1 SUSY gauge theories: anomalies and central functions

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$

$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$

$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{ccc} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation

coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3
$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3
$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3
$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3
$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3
$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3
$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3
$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1, 2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category $1, 2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$

$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$

$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$

$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$

$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ -(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$
$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$
$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$
$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{44})^{(9)} & +(a''_{44})^{(9)}(T_{45},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\begin{bmatrix} (a'_{45})^{(9)} & +(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{c} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \left[\begin{array}{c} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)} \boxed{-(b''_{45})^{(9)}(G_{47},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)} \boxed{-(b''_{46})^{(9)}(G_{47},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{15}$$

Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first

augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \tag{127}$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \tag{128}$$

$$\lim_{G \to \infty} (b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \tag{131}$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

$$\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$$

$$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and $(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \quad (136)$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad (138)$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad (139)$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad (140)$$

$$\lim_{G\to\infty}(b_i'')^{(8)}\big((G_{43}),t\big) = (r_i)^{(8)} \quad (141)$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad (142)$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\big((G_{43}),t\big)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad (143)$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad (144)$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad (145)$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad (146)$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad (146\,A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} \text{ , } T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} \text{ , } T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$$ 169

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right)\left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1, T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1 T_{14}$ which leads to

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1 t})+T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t}=\frac{1}{2}$ it results

$T_{14}\geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad (226)$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ (228)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: (229)

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right) - (a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right| \; e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\} ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}; \; (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i \; and \; T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \; and \; (\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and \; not \; on \; t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ (250)

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(6)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If $G_{32} \; is$ bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)},, (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le -(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le -(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le -(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le -(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2+(\sigma_2)^{(3)}\nu^{(3)}-(a_{20})^{(3)}\right)\leq\frac{d\nu^{(3)}}{dt}\leq-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2+(\sigma_1)^{(3)}\nu^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(\nu_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\nu_1)^{(3)}<(\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t)\geq\frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)}\leq\nu^{(3)}(t)\leq(\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t)\leq\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)}\leq\nu^{(3)}(t)\leq(\bar{\nu}_1)^{(3)}$

If $0<(\nu_1)^{(3)}<(\nu_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)}\leq\frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}\leq\nu^{(3)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}\leq(\bar{\nu}_1)^{(3)}$$

If $0<(\nu_1)^{(3)}\leq(\bar{\nu}_1)^{(3)}\leq(\nu_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)}\leq\nu^{(3)}(t)\leq\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}\leq(\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)}\leq\nu^{(3)}(t)\leq(m_1)^{(3)},\quad\boxed{\nu^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\leq u^{(3)}(t)\leq(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)}=(\bar{\nu}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(5)} and\ (b''_i)^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a''_i)^{(6)} and\ (b''_i)^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(n) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^{*}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^{*}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^{*}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^{*}) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^{*}) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}(\,(G_{39})^{**}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 15:

Holographic Anomalies And Higher Derivative Gravity, Two-Hadron Semi-Inclusive Production Including Subleading Twist Et Al,: A Rational Leibneizism And Socratic Subjectivity Model

1. Holographic anomalies and higher derivative gravity
2. Two-hadron semi-inclusive production Vis-aVis subleading twist
3. New observables in longitudinal single-spin asymmetries Vis-a-Vis semi-inclusive DIS.
4. Single-spin asymmetries Vis-aVis semi-inclusive deep-inelastic scattering
5. Single-spin asymmetries: and Collins fragmentation function
6. Di hadron interference fragmentation functions Vis-aVis proton-proton collisions.
7. Non-centrosymmetric superconductor and penetration depth,
8. Super neutronic, Superheavy Vis-à-vis Anti Matter
9. Bose-Einstein condensation Vis-aVis neutron star core

NOTATION

Module One

Holographic anomalies and higher derivative gravity

G_{13} : Category one of Holographic anomalies

G_{14} : Category two of Holographic anomalies

G_{15} : Category three of Holographic anomalies

T_{13} : Category one of higher derivative gravity(gravity varies at different points in the universe)

T_{14} : Category two of higher derivative gravity

T_{15} : Category three of higher derivative gravity

Module Two

Two-hadron semi-inclusive production Vis-à-vis subleading twist

G_{16} : Category one of Two-hadron semi-inclusive production

G_{17} : Category two of Two-hadron semi-inclusive production

G_{18} : Category three of Two-hadron semi-inclusive production

T_{16} : Category one of subleading twist

T_{17} : Category two of subleading twist

T_{18} : Category three of subleading twist

Module three

New observables in longitudinal single-spin asymmetries Vis-à-vis semi-inclusive DIS

G_{20} : Category one of New observables in longitudinal single-spin asymmetries

G_{21} : Category two of New observables in longitudinal single-spin asymmetries

G_{22} : Category three of New observables in longitudinal single-spin asymmetries

T_{20} : Category one of semi-inclusive DIS

T_{21} : Category two of semi-inclusive DIS

T_{22} : Category three of semi-inclusive DIS

Module four

Single-spin asymmetries Vis-à-vis semi-inclusive deep-inelastic scattering

G_{24} : Category one of Single-spin asymmetries

G_{25} : Category two of Single-spin asymmetries

G_{26} : Category three of Single-spin asymmetries

T_{24} : Category one of semi-inclusive deep-inelastic scattering

T_{25} : Category two of semi-inclusive deep-inelastic scattering

T_{26} : Category three of semi-inclusive deep-inelastic scattering

Module five

Single-spin asymmetries Vis-à-vis Collins fragmentation function

G_{28} : Category one of Single-spin asymmetries

G_{29} : Category two of Single-spin asymmetries

G_{30} : Category three of Single-spin asymmetries

T_{28} : Category one of Collins fragmentation function

T_{29} : Category two of Collins fragmentation function

T_{30} : Category three of Collins fragmentation function

Module six

Di hadron interference fragmentation functions Vis-à-vis proton-proton collisions

G_{32} : Category one of Di hadron interference fragmentation functions

G_{33} : Category two of Di hadron interference fragmentation functions

G_{34} : Category three of Di hadron interference fragmentation functions

T_{32} : Category one of proton-proton collisions

T_{33} : Category two of proton-proton collisions

T_{34} : Category three of proton-proton collisions

Module seven

non-centrosymmetric superconductors Vis-à-vis penetration depth

G_{36} : Category one of non-centrosymmetric superconductors

G_{37} : Category two of non-centrosymmetric superconductors

G_{38} : Category three of non-centrosymmetric superconductors

T_{36} : Category one of penetration depth

T_{37} : Category two of penetration depth

T_{38} : Category three of penetration depth

Module eight

Super neutronic, Superheavy matter Vis-à-vis Anti Matter

G_{40} : Category one of Super neutronic, Superheavy matter

G_{41} : Category two of Super neutronic, Superheavy matter

G_{42} : Category three of Super neutronic, Superheavy matter

T_{40} : Category one of Anti Matter

T_{41} : Category two of Anti Matter

T_{42} : Category three of Anti Matter

Module Nine

Bose-Einstein condensation Vis-à-vis neutron star core

G_{44} : Category one of Bose-Einstein condensation

G_{45} : Category two of Bose-Einstein condensation

G_{46} : Category three of Bose-Einstein condensation

T_{44} : Category one of neutron star core

T_{45} : Category two of neutron star core

T_{46} : Category three of neutron star core

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \big[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\big]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \big[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\big]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \big[(b'_{32})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big)\big]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \big[(b'_{33})^{(6)} - (b''_{33})^{(6)}\big((G_{35}),t\big)\big]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \big[(b'_{34})^{(6)} - (b''_{34})^{(6)}\big((G_{35}),t\big)\big]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \big[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\big]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \big[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\big]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \big[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\big]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \big[(b'_{36})^{(7)} - (b''_{36})^{(7)}\big((G_{39}),t\big)\big]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \big[(b'_{37})^{(7)} - (b''_{37})^{(7)}\big((G_{39}),t\big)\big]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \big[(b'_{38})^{(7)} - (b''_{38})^{(7)}\big((G_{39}),t\big)\big]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \big[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\big]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \big[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\big]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \big[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\big]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \big[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}),t\big)\big]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \big[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}),t\big)\big]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \big[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}),t\big)\big]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \big[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\big]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & \boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & \boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \qquad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation*

$coefficient\ \ for\ category\ 1,2\ and\ 3$

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ $are\ third\ \ augmentation$
$coefficient\ for\ category\ 1,2\ and\ 3$

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
$are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
$are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
$are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
$are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
$are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \qquad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \qquad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \qquad 78$$

$Where$ $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients$
$for\ category\ 1,2\ and\ 3$

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients$
$for\ category\ 1,2\ and\ 3$

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients$
$for\ category\ 1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ are third augmentation

$coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34}$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ *are first detrition coefficients for category* $1,2\ and\ 3$

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ *are second detrition coefficients for category* $1,2\ and\ 3$

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ *are third detrition coefficients for category* $1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$
$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$
$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$
$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{37}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$\frac{dG_{40}}{dt}$ *95*

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{44})^{(9)} & +(a''_{44})^{(9)}(T_{45},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\begin{bmatrix} (a'_{45})^{(9)} & +(a''_{45})^{(9)}(T_{45},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \left[\begin{array}{l} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array} \right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \left[\begin{array}{l} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)} T_{44} - \left[\begin{array}{l} (b'_{45})^{(9)} \boxed{-(b''_{45})^{(9)}(G_{47},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)} T_{45} - \left[\begin{array}{l} (b'_{46})^{(9)} \boxed{-(b''_{46})^{(9)}(G_{47},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array} \right] T_{15}$$

Where $-(b_{44}'')^{(9)}(G_{47},t)$, $-(b_{45}'')^{(9)}(G_{47},t)$, $-(b_{46}'')^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)},(a_i')^{(1)},(a_i'')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(b_i'')^{(1)}>0,\quad i,j=13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)},(b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t)\le(p_i)^{(1)}\le(\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t)\le\ (r_i)^{(1)}\le(b_i')^{(1)}\le(\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t)=(p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t)=\ (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)},(p_i)^{(1)},\ (r_i)^{(1)}$ are positive constants and $i=13,14,15$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$.(T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first

augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty} (b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

$$\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$$

$$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and $(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}$, $(r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \quad (136)$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G\to\infty}(b_i'')^{(8)}\big((G_{43}),t\big) = (r_i)^{(8)} \quad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\big((G_{43}),t\big)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} , T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \ , \ T_i(0) = T_i^0 \ , \ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} , T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A
which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that
$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+$ 169
$\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}= \quad 180$$
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \le \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right) \quad 186$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\ (G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1, T_{13}\ (t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1 T_{14}$ which leads to

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1 t})+T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t}=\ \frac{1}{2}$ it results

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\ (G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16{,}17{,}18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(3)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})},\widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})},\widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{28})^{(5)} \left|G_{29}^{(1)}-G_{29}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}}\, ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} \left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} +$$

$$G_{28}^{(2)} \left|(a''_{28})^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)}, s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)} s_{(28)}} e^{(\widehat{M}_{28})^{(5)} s_{(28)}} \} ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right| e^{-(\widehat{M}_{28})^{(5)} t} \leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}\right) d\left(((G_{31})^{(1)}, (T_{31})^{(1)};\ (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ *and* $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 239

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a'_{29})^{(5)} G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)} \left((\widehat{M}_{28})^{(5)}\right)_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)} \left((\widehat{M}_{28})^{(5)}\right)_2 / (a'_{30})^{(5)}$$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}\left|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq$ 259

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval [0, t] 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, $if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le -(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le -(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le -(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le -(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, ***if*** $(v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, ***if*** $(v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, ***if*** $(\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(5)} and\ (b''_i)^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a''_i)^{(6)} and\ (b''_i)^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484A$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(o) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]}$, $G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^* \mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}$, $\frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^*\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^*\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^*\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\big(\,(G_{39})^{**}\,\big) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^*\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^*\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^*\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Section 16:

Superconductivity and Neutrons, magnetic ordering on dynamical conductivity, Transport Properties and Magnetic properties Et Al, : Predicational Interiority and Primordial Exactitude thereof

1. Distribution of the Relaxation Times Vis-à-vis Dielectric Spectra
2. Phase transition and Ionic conductivity
3. Acoustoelectric effects Vis-à-vis Sn2P2S6-type ferroelectric semiconductors
4. Superconductivity and Neutrons
5. magnetic ordering on dynamical conductivity
6. Transport Properties and Magnetic properties
7. superconducting fluctuations and pseudogap effects
8. Pseudogap and Superconductivity
9. Vortex dynamics and Phase diagram

NOTATION

Module One

Distribution of the Relaxation Times Vis-à-vis Dielectric Spectra

G_{13} : Category one of Dielectric Spectra

G_{14} : Category two of Dielectric Spectra

G_{15} : Category three of Dielectric Spectra

T_{13} : Category one of Distribution of the Relaxation Times

T_{14} : Category two of Distribution of the Relaxation Times

T_{15} : Category three of Distribution of the Relaxation Times

Module Two

Phase transition and Ionic conductivity

G_{16} : Category one of Phase transition

G_{17} : Category two of Phase transition

G_{18} : Category three of Phase transition

T_{16} : Category one of Ionic conductivity

T_{17} : Category two of Ionic conductivity

T_{18} : Category three of Ionic conductivity

Module three

Acoustoelectric effects Vis-à-vis Sn2P2S6-type ferroelectric semiconductors

G_{20} : Category one of Acoustoelectric effects

G_{21} : Category two of Acoustoelectric effects

G_{22} : Category three of Acoustoelectric effects

T_{20} : Category one of Sn2P2S6-type ferroelectric semiconductors

T_{21} : Category two of Sn2P2S6-type ferroelectric semiconductors

T_{22} : Category three of Sn2P2S6-type ferroelectric semiconductors

Module four

Superconductivity Vis-à-vis Neutrons

G_{24} : Category one of Superconductivity

G_{25} : Category two of Superconductivity

G_{26} : Category three of Superconductivity

T_{24} : Category one of Neutrons

T_{25} : Category two of Neutrons

T_{26} : Category three of Neutrons

Module five

magnetic ordering Vis-à-vis dynamical conductivity

G_{28} : Category one of magnetic ordering

G_{29} : Category two of magnetic ordering

G_{30} : Category three of magnetic ordering

T_{28} : Category one of dynamical conductivity

T_{29} : Category two of dynamical conductivity

T_{30} : Category three of dynamical conductivity

Module six

Transport Properties Vis-à-vis Magnetic properties

G_{32} : Category one of Transport Properties

G_{33} : Category two of Transport Properties

G_{34} : Category three of Transport Properties

T_{32} : Category one of Magnetic properties

T_{33} : Category two of Magnetic properties

T_{34} : Category three of Magnetic properties

Module seven

Superconducting fluctuations and pseudogap effects

G_{36} : Category one of superconducting fluctuations

G_{37} : Category two of superconducting fluctuations

G_{38} : Category three of superconducting fluctuations

T_{36} : Category one of pseudogap effects

T_{37} : Category two of pseudogap effects

T_{38} : Category three of pseudogap effects

Module eight

Pseudogap and Superconductivity

G_{40} : Category one of Pseudogap

G_{41} : Category two of Pseudogap

G_{42} : Category three of Pseudogap

T_{40} : Category one of Superconductivity

T_{41} : Category two of Superconductivity

T_{42} : Category three of Superconductivity

Module Nine

Vortex dynamics and Phase diagram

G_{44} : Category one of Vortex dynamics

G_{45} : Category two of Vortex dynamics

G_{46} : Category three of Vortex dynamics

T_{44} : Category one of Phase diagram

T_{45} : Category two of Phase diagram

T_{46} : Category three of Phase diagram

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$

are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\big((G_{31}),t\big) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \big[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\big]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \big[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\big]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \big[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\big]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \big[(b'_{32})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big)\big]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \big[(b'_{33})^{(6)} - (b''_{33})^{(6)}\big((G_{35}),t\big)\big]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \big[(b'_{34})^{(6)} - (b''_{34})^{(6)}\big((G_{35}),t\big)\big]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \big[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\big]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \big[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\big]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \big[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\big]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \big[(b'_{36})^{(7)} - (b''_{36})^{(7)}\big((G_{39}),t\big)\big]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \big[(b'_{37})^{(7)} - (b''_{37})^{(7)}\big((G_{39}),t\big)\big]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \big[(b'_{38})^{(7)} - (b''_{38})^{(7)}\big((G_{39}),t\big)\big]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \big[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\big]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \big[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\big]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \big[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\big]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \big[(b'_{40})^{(8)} - (b''_{40})^{(8)}\big((G_{43}),t\big)\big]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \big[(b'_{41})^{(8)} - (b''_{41})^{(8)}\big((G_{43}),t\big)\big]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \big[(b'_{42})^{(8)} - (b''_{42})^{(8)}\big((G_{43}),t\big)\big]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \big[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\big]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b_{36}'')^{(7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b_{40}'')^{(8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b_{44}'')^{(9,9)}(G_{47},t)$, $-(b_{46}'')^{(9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a_{20}')^{(3)} & +(a_{20}'')^{(3)}(T_{21},t) & +(a_{16}'')^{(2,2,2)}(T_{17},t) & +(a_{13}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a_{36}'')^{(7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a_{21}')^{(3)} & +(a_{21}'')^{(3)}(T_{21},t) & +(a_{17}'')^{(2,2,2)}(T_{17},t) & +(a_{14}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a_{37}'')^{(7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a_{22}')^{(3)} & +(a_{22}'')^{(3)}(T_{21},t) & +(a_{18}'')^{(2,2,2)}(T_{17},t) & +(a_{15}'')^{(1,1,1,)}(T_{14},t) \\ +(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a_{38}'')^{(7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad (69)$$

$+(a_{20}'')^{(3)}(T_{21},t)$, $+(a_{21}'')^{(3)}(T_{21},t)$, $+(a_{22}'')^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a_{36}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a_{40}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a_{44}'')^{(9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9)}(T_{45},t)$, $+(a_{46}'')^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b_{20}')^{(3)} & -(b_{20}'')^{(3)}(G_{23},t) & -(b_{16}'')^{(2,2,2)}(G_{19},t) & -(b_{13}'')^{(1,1,1,)}(G,t) \\ -(b_{24}'')^{(4,4,4,4,4,4)}(G_{27},t) & -(b_{28}'')^{(5,5,5,5,5,5)}(G_{31},t) & -(b_{32}'')^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b_{36}'')^{(7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \qquad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation*

$coefficient\ \ for\ category\ 1,2\ and\ 3$

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ $are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ $are\ fourth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ $are\ fifth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ $are\ sixth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ $are\ seventh\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ $are\ eighth\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where$ $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$ $are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$ $are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$ $are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$ $are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3
$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3
$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3
$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3
$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ are third augmentation

$coefficients\ for\ category\ 1,2\ and\ 3$

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$ Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35},t) & -(b''_{28})^{(5,5,5)}(G_{31},t) & -(b''_{24})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{13})^{(1,1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35},t) & -(b''_{29})^{(5,5,5)}(G_{31},t) & -(b''_{25})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{14})^{(1,1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35},t) & -(b''_{30})^{(5,5,5)}(G_{31},t) & -(b''_{26})^{(4,4,4,)}(G_{27},t) \\ & -(b''_{15})^{(1,1,1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{34}$$

$-(b''_{32})^{(6)}(G_{35},t)$, $-(b''_{33})^{(6)}(G_{35},t)$, $-(b''_{34})^{(6)}(G_{35},t)$ $are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{28})^{(5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5)}(G_{31},t)$ $are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{24})^{(4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,)}(G_{27},t)$ $are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$-(b''_{13})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,1)}(G,t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{37}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)} & +(a''_{41})^{(8)}(T_{41},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & \\ +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\left[\begin{array}{ccc} (b'_{42})^{(8)} & \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} \\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} & \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} \\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} & \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} \\ \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} & & \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\left[\begin{array}{ccc} (a'_{44})^{(9)} & \boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \\ \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} & \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \\ \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} & \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \\ \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & & \end{array}\right] G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\left[\begin{array}{ccc} (a'_{45})^{(9)} & \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} \\ \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} & \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} \\ \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} & \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} \\ \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & & \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{c} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)}\ \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\ \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[\begin{array}{c} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)}\ \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[\begin{array}{c} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)}\ \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[\begin{array}{c} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)}\ \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}\ \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b_{44}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first

augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\hat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (110)$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty} (b_i'')^{(6)}\left((G_{35}), t\right) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\left((G_{35}), t\right)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\hat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

M) $$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

)The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \quad (132)$$

$$\lim_{T_2\to\infty} (a_i'')^{(7)}(T_{37},t) = (p_i)^{(7)}$$

$$\lim_{G\to\infty} (b_i'')^{(7)}\big((G_{39}),t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: (133)

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\big((G_{39}),t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and $(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: (134)

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: (135)

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)}(\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \quad (136)$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: (137)

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \quad (138)$$

$$(b_i'')^{(8)}((G_{43}),t) \le (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \quad (139)$$

$$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)} \quad (140)$$

$$\lim_{G\to\infty}(b_i'')^{(8)}((G_{43}),t) = (r_i)^{(8)} \quad (141)$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t} \quad (142)$$

$$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}((G_{43}),t)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t} \quad (143)$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad (144)$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad (145)$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1 \quad (146)$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad (146\,A)$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}\big((G_{47}),t\big)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $G_i(0) = G_i^0 > 0$

$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$, $T_i(0) = T_i^0 > 0$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0), T_i(0)$:

$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{G_i(0) = G_i^0 > 0}$
$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{\left(\hat{M}_{44}\right)^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} \text{ , } T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{\left(\hat{M}_{13}\right)^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} \text{ , } T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{\left(\hat{M}_{16}\right)^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^{0} + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{44})^{(9)} \text{ , } T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^{0} + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^{0} + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^{0} + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^{0} + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^{0} + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167
that

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1+(a_{13})^{(1)}t\right)G_{14}^{0}+\frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)}t}-1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+$ 169
$\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right)$

From which it follows that 170

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

179

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)}\left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$G_{13}^{(2)} \left|(a''_{13})^{(1)}\left(T_{14}^{(1)}, s_{(13)}\right) - (a''_{13})^{(1)}\left(T_{14}^{(2)}, s_{(13)}\right)\right| \; e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\} ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}\,(G(t),t))=(b'_{14})^{(1)}$ then $T_{14}\ \to\infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b''_i)^{(1)}(G(t),t)<\varepsilon_1,T_{13}\,(t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt}\geq(a_{14})^{(1)}(m)^{(1)}-\varepsilon_1T_{14}$ which leads to

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1-e^{-\varepsilon_1t})+T_{14}^0e^{-\varepsilon_1t}$ If we take t such that $e^{-\varepsilon_1t}=\ \frac{1}{2}$ it results

$T_{14}\ \geq\left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right),\quad t=log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}\,(G(t),t)=(b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|\mathrm{e}^{-(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}} \leq$$
$$\frac{1}{(\widehat{\mathrm{M}}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{\mathrm{A}}_{16})^{(2)} + (\widehat{\mathrm{P}}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{\mathrm{P}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ and $(\widehat{\mathrm{Q}}_{16})^{(2)}\mathrm{e}^{(\widehat{\mathrm{M}}_{16})^{(2)}\mathrm{t}}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16{,}17{,}18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

206

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:\left(\widetilde{(G_{23})},\widetilde{(T_{23})}\right)=\mathcal{A}^{(3)}\left((G_{23}),(T_{23})\right)$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};\ (G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ and\ (\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{\left(-(b_i')^{(3)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1,\left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{24})^{(4)} \left|G_{25}^{(1)} - G_{25}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\, ds_{(24)} +$$

$$\int_0^t \{(a'_{24})^{(4)} \left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}} +$$

$$G_{24}^{(2)} \left|(a''_{24})^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a''_{24})^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)} s_{(24)}} e^{(\widehat{M}_{24})^{(4)} s_{(24)}}\} ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right| e^{-(\widehat{M}_{24})^{(4)} t} \leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right) d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ *and* $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2 / (a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right)$, $t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_{0}^{t}\{(a_{28}')^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a_{28}'')^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a_{28}'')^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a_{28}'')^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a_{28}')^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a_{28}'')^{(5)}$ and $(b_{28}'')^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 239

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{(-(b_i')^{(5)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\leq\left((\widehat{M}_{28})^{(5)}\right)_1-(a_{29}')^{(5)}G_{29}$ and by integrating

$$G_{29}\leq\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30}\leq\left((\widehat{M}_{28})^{(5)}\right)_3=G_{30}^0+2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$

$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t} \leq \frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)};\ (G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \quad (248)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (250)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\}ds_{(32)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(6)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: (251)

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. (252)

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\ ((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. (253)

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t), t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a_{36}'')^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)} - (G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$

$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 261

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)}G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right),\ t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose $(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}), (\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}), (\widetilde{T_{43}})\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{40})^{(8)}\left|G_{41}^{(1)} - G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)} +$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)\left|G_{40}^{(1)} - G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a''_{40})^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t} \leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)} \left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$ and to choose
$(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[(\hat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44,45,46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(9)}t)}>0\quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44}<(\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt}\le\left((\widehat{M}_{44})^{(9)}\right)_1-(a'_{45})^{(9)}G_{45}$ and by integrating

$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$, $if\ (v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$
$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T^0_{16}}{T^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, ***if*** $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G^0_{16}e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}G^0_{16}e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}G^0_{16}e^{(S_1)^{(2)}t} \quad 311$$

$$\Big(\frac{(a_{18})^{(2)}G^0_{16}}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\Big[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\Big] + G^0_{18}e^{-(S_2)^{(2)}t} \le G_{18}(t) \le \frac{(a_{18})^{(2)}G^0_{16}}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}] + G^0_{18}e^{-(a'_{18})^{(2)}t}\Big) \quad 312$$

$$\boxed{T^0_{16}e^{(R_1)^{(2)}t} \le T_{16}(t) \le T^0_{16}e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} - \mathrm{e}^{-(b'_{18})^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} - \mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right] + \mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying
$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$

$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by
$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^{0}}{G_{21}^{0}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)} t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \le G_{22}(t) \le \right.$ 324
$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$

$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \le -(\tau_1)^{(4)}$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_1)^{(4)} v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_1)^{(4)} u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)} > 0$, $(\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)},(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)},$ ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)},$ ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)},$ ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)},$ ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le\right.$$

334

(a26)4G240(m2)4(S1)4−(a26′)4e(S1)4t−e−(a26′)4t+ G260e−(a26′)4t

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)},(S_2)^{(4)},(R_1)^{(4)},(R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)}, , (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}, \ \boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}, \boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$$

and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \ \boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G^0_{28} e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G^0_{28} e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

(a30)5G280(m2)5(S1)5−(a30′)5e(S1)5t−e−(a30′)5t+ G300e−(a30′)5t

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},\ \boldsymbol{if}\ (\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}, \boldsymbol{if}\ (\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$
and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$
(a34)6G320(m2)6(S1)6−(a34′)6e(S1)6t−e−(a34′)6t+ G340e−(a34′)6t

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

360

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36}e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}}G^0_{36}e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}}G^0_{36}e^{(S_1)^{(7)}t} \quad 365$$

$$\Big(\frac{(a_{38})^{(7)}G^0_{36}}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\Big[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\Big] + G^0_{38}e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \quad 366$$

$\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}]+\ G_{38}^{0}e^{-(a'_{38})^{(7)}t})$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\le T_{36}(t)\le \frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\le T_{38}(t)\le \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$-(\sigma_2)^{(8)}\le -(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\le -(\sigma_1)^{(8)}$

$-(\tau_2)^{(8)}\le -(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\le -(\tau_1)^{(8)}$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0,(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0,(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0,(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0,(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)},(v_0)^{(8)}$:-

If we define $(m_1)^{(8)},(m_2)^{(8)},(\mu_1)^{(8)},(\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(v_0)^{(8)},(m_1)^{(8)}=(v_1)^{(8)},$ ***if*** $(v_0)^{(8)}<(v_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}$ $(\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}$ $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}$ $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}$ $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)} t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)} t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)} t}\right] + G_{42}^0 e^{-(S_2)^{(8)} t} \leq G_{42}(t) \leq \right. \quad 377$$
$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)} t} - e^{-(a'_{42})^{(8)} t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)} t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)} t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)} t} - e^{-(b'_{42})^{(8)} t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)} t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)} t}\right] + T_{42}^0 e^{-(R_2)^{(8)} t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)},, (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}, \ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$$

$$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}\ , \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$$

and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}\ , \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G^0_{44}e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G^0_{44}e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}}G^0_{44}e^{(S_1)^{(9)}t}$$

$$\Big(\frac{(a_{46})^{(9)}G^0_{44}}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\Big[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t}\Big] + G^0_{46}e^{-(S_2)^{(9)}t} \le G_{46}(t) \le \frac{(a_{46})^{(9)}G^0_{44}}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}[e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t}] + G^0_{46}e^{-(a'_{46})^{(9)}t}\Big)$$

$$\boxed{T^0_{44}e^{(R_1)^{(9)}t} \le T_{44}(t) \le T^0_{44}e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0 e^{(R_1)^{(9)}t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}}T_{44}^0 e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)}T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t}\right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right) - (a''_{14})^{(1)}(T_{14},t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \le \frac{dv^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{v}_1)^{(1)}+(\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}} \leq (\bar{v}_1)^{(1)}$$

If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)}+(\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{v}_1)^{(1)}-(\bar{v}_2)^{(1)}\right)t\right]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)}+(C)^{(2)}(v_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)}-(v_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((v_1)^{(2)}-(v_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)}-(v_0)^{(2)}}{(v_0)^{(2)}-(v_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\le\frac{dv^{(3)}}{dt}\le-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\ge\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\le v^{(3)}(t)\le(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\le v^{(3)}(t)\le(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\le\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\le v^{(3)}(t)\le$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\le(\bar{v}_1)^{(3)}\le(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\le v^{(3)}(t)\le\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\le(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\le v^{(3)}(t)\le(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\le u^{(3)}(t)\le(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)}=(a_{21}'')^{(3)},then\ (\sigma_1)^{(3)}=(\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)}=(\bar{v}_1)^{(3)}$ if in addition

$(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \le \frac{d\nu^{(4)}}{dt} \le -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \ge \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \le \nu^{(4)}(t) \le (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$, we obtain 407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(v_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(v_2)^{(6)}}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{v}_1)^{(6)}-(v_0)^{(6)}}{(v_0)^{(6)}-(\bar{v}_2)^{(6)}}}$$

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case, 414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)}+(C)^{(6)}(v_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((v_1)^{(6)}-(v_2)^{(6)}\right)t\right]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (\bar{v}_1)^{(6)}$$

If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)}+(\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{v}_1)^{(6)}-(\bar{v}_2)^{(6)}\right)t\right]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $\nu^{(7)}$:- $\boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$ if in addition $(v_0)^{(8)} = (v_1)^{(8)}$ then $v^{(8)}(t) = (v_0)^{(8)}$ and as a consequence $G_{40}(t) = (v_0)^{(8)} G_{41}(t)$ **this also defines** $(v_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(v_1)^{(8)}$ and $(\bar{v}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

Definition of $v^{(9)}$:- $\quad \boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)}\right) \leq \frac{dv^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(v^{(9)}\right)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$:-

For $0 < \boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)} - (v_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}}$$

it follows $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.

We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(p) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]}$, $G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]}$, $G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]}$, $G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]}$, $G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three

first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right]+(b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right]+(b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$
$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi(G^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - \left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi(G^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(\mathrm{T}^*_{17}) = 0$ and 512

$$\mathrm{G}^*_{16} = \frac{(\mathrm{a}_{16})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{16})^{(2)} + (\mathrm{a}''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad \mathrm{G}^*_{18} = \frac{(\mathrm{a}_{18})^{(2)}\mathrm{G}^*_{17}}{[(\mathrm{a}'_{18})^{(2)} + (\mathrm{a}''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 513$$

$$\mathrm{T}^*_{16} = \frac{(\mathrm{b}_{16})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{16})^{(2)} - (\mathrm{b}''_{16})^{(2)}((G_{19})^*)]} \quad , \quad \mathrm{T}^*_{18} = \frac{(\mathrm{b}_{18})^{(2)}\mathrm{T}^*_{17}}{[(\mathrm{b}'_{18})^{(2)} - (\mathrm{b}''_{18})^{(2)}((G_{19})^*)]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)} G_{25}^{*}}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)} G_{25}^{*}}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)} T_{25}^{*}}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)} T_{25}^{*}}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)} T_{13}^* \mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)} T_{14}^* \mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)} T_{15}^* \mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)} G_{16}^* \mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad (535)$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad (536)$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad (537)$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad (538)$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad (539)$$

ASYMPTOTIC STABILITY ANALYSIS (540)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad (541)$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad (542)$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad (543)$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad (544)$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad (545)$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad (546)$$

ASYMPTOTIC STABILITY ANALYSIS (547)

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- (548)

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}\left((G_{27})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^{*}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^{*}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\big((a_{32}')^{(6)} + (p_{32})^{(6)}\big)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^{*}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\big((a_{33}')^{(6)} + (p_{33})^{(6)}\big)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^{*}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\big((a_{34}')^{(6)} + (p_{34})^{(6)}\big)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^{*}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\big((b_{32}')^{(6)} - (r_{32})^{(6)}\big)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(32)(j)}T_{32}^{*}\mathbb{G}_j\big) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\big((b_{33}')^{(6)} - (r_{33})^{(6)}\big)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\big(s_{(33)(j)}T_{33}^{*}\mathbb{G}_j\big) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\big((b_{34}')^{(6)} - (r_{34})^{(6)}\big)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\big(s_{(34)(j)}T_{34}^{*}\mathbb{G}_j\big) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}(\,(G_{39})^{**}\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\big((a_{36}')^{(7)} + (p_{36})^{(7)}\big)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\big((a_{37}')^{(7)} + (p_{37})^{(7)}\big)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\big((a_{38}')^{(7)} + (p_{38})^{(7)}\big)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\big((b_{36}')^{(7)} - (r_{36})^{(7)}\big)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\big) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\big((b_{37}')^{(7)} - (r_{37})^{(7)}\big)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\big(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\big) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\big((b_{38}')^{(7)} - (r_{38})^{(7)}\big)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\big(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\big) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

ASYMPTOTIC STABILITY ANALYSIS (586 A)

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is

$$(\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\Big]$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\Big)$$

$$+\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^{*}_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\Big)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2} + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left((\lambda)^{(1)}\right)^{2} + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14} + (b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\Big]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\Big)$$

$$+\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\Big)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2} + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left((\lambda)^{(2)}\right)^{2} + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\Big)\} = 0$$

$$+$$

$$(\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21} + (a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\Big)\right]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\Big)\right]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\Big)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)\right.$$

$$\left.\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)\right.$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\right)\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)\right.$$

$$\left.\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)\right.$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

REFERENCES

[1] Van Hove, Leon (1958). "Von Neumann's contributions to quantum mechanics"(PDF). Bulletin of the American Mathematical Society 64: Part2:95–99.
[2] Max Born & Emil Wolf, Principles of Optics, 1999, Cambridge University Press
[3] Mehra, J.; Rechenberg, H. (1982). The historical development of quantum theory. New York: Springer-Verlag. ISBN 0387906428.
[4] http://www.ias.ac.in/resonance/December2010/p1056-1059.pdf
[5] Kuhn, T. S. (1978). Black-body theory and the quantum discontinuity 1894-1912. Oxford: Clarendon Press. ISBN 0195023838.
[6] Einstein, A. (1905). "Über einen die Erzeugung und Verwandlung des Lichtes betreffenden heuristischen Gesichtspunkt [On a heuristic point of view concerning the production and transformation of light]". Annalen der Physik 17 (6): 132–48. Bibcode1905AnP...322...132E. doi:10.1002/andp .19053220607. Reprinted in The collected papers of Albert Einstein, John Stachel, editor, Princeton University Press, 1989, Vol. 2, pp. 149-166, in German; see also Einstein's early work on the quantum hypothesis, ibid. pp. 134-148.
[7] "Quantum - Definition and More from the Free Merriam-Webster Dictionary". Merriam-webster.com. Retrieved 2012-08-18.
[8] http://mooni.fccj.org/~ethall/quantum/quant.htm
[9] Compare the list of conferences presented here
[10] Oocities.com at the Wayback Machine (archived October 26, 2009)
[11] P.A.M. Dirac, The Principles of Quantum Mechanics, Clarendon Press, Oxford, 1930.
[12] D. Hilbert Lectures on Quantum Theory, 1915-1927
[13] Greiner, Walter; Müller, Berndt (1994). Quantum Mechanics Symmetries, Second edition. Springer-Verlag. p. 52. ISBN 3-540-58080-8., Chapter 1, p. 52
[14] "Heisenberg - Quantum Mechanics, 1925-1927: The Uncertainty Relations". Aip.org. Retrieved 2012-08-18.
[15] Greenstein, George; Zajonc, Arthur (2006). The Quantum Challenge: Modern Research on the Foundations of Quantum Mechanics, Second edition. Jones and Bartlett Publishers, Inc. p. 215. ISBN 0-7637-2470-X., Chapter 8, p. 215
[16] "[Abstract] Visualization of Uncertain Particle Movement". Actapress.com. Retrieved 2012-08-18.
[17] Hirshleifer, Jack (2001). The Dark Side of the Force: Economic Foundations of Conflict Theory. Campbridge University Press. p. 265. ISBN 0-521-80412-4., Chapter , p. Dict.cc De .pons .eu
[18] "Topics: Wave-Function Collapse". Phy.olemiss.edu. 2012-07-27. Retrieved 2012-08-18.
[19] "Collapse of the wave-function". Farside.ph.utexas.edu. Retrieved 2012-08-18.
[20] "Determinism and Naive Realism: philosophy". Reddit.com. 2009-06-01. Retrieved 2012-08-18.
[21] Michael Trott. "Time-Evolution of a Wavepacket in a Square Well — Wolfram Demonstrations Project". Demonstrations.wolfram.com. Retrieved 2010-10-15.
[22] Michael Trott. "Time Evolution of a Wavepacket In a Square Well". Demonstrations.wolfram.com. Retrieved 2010-10-15.
[23] Mathews, Piravonu Mathews; Venkatesan, K. (1976). A Textbook of Quantum Mechanics. Tata McGraw-Hill. p. 36. ISBN 0-07-096510-2., Chapter 2, p. 36
[24] "Wave Functions and the Schrödinger Equation" (PDF). Retrieved 2010-10-15.
[25] "Quantum Physics: Werner Heisenberg Uncertainty Principle of Quantum Mechanics. Werner

Heisenberg Biography". Spaceandmotion.com. 1976-02-01. Retrieved 2012-08-18.
[26] http://th-www.if.uj.edu.pl/acta/vol19/pdf/v19p0683.pdf
[27] Nancy Thorndike Greenspan, "The End of the Certain World: The Life and Science of Max Born" (Basic Books, 2005), pp. 124-8 and 285-6.
[28] http://ocw.usu.edu/physics/classical-mechanics/pdf_lectures/06.pdf
[29] "The Nobel Prize in Physics 1979". Nobel Foundation. Retrieved 2010-02-16.
[30] Carl M. Bender, Daniel W. Hook, Karta Kooner (2009-12-31). "Complex Elliptic Pendulum". ArXiv: 1001.0131 [hep-th].
[31] "Quantum mechanics course iwhatisquantummechanics". Scribd.com. 2008-09-14. Retrieved 2012-08-18.
[32] "Between classical and quantum " (PDF). Retrieved 2012-08-19.
[33] "Atomic Properties". Academic.brooklyn.cuny.edu. Retrieved 2012-08-18.
[34] http://assets.cambridge.org/97805218/29526/excerpt/9780521829526_excerpt.pdf
[35] "There is as yet no logically consistent and complete relativistic quantum field theory.", p. 4. — V. B. Berestetskii, E. M. Lifshitz, L P Pitaevskii (1971). J. B. Sykes, J. S. Bell (translators). Relativistic Quantum Theory 4, part I. Course of Theoretical Physics (Landau and Lifshitz) ISBN 0-08-016025-5
[36] http://www.damtp.cam.ac.uk/strings02/dirac/hawking/
[37] "Life on the lattice: The most accurate theory we have". Latticeqcd.blogspot.com. 2005-06-03. Retrieved 2010-10-15.
[38] Parker, B. (1993). Overcoming some of the problems. pp. 259–279.
[39] The Character of Physical Law (1965) Ch. 6; also quoted in The New Quantum Universe (2003), by Tony Hey and Patrick Walters
[40] "Action at a Distance in Quantum Mechanics (Stanford Encyclopedia of Philosophy)". Plato.stanford.edu. 2007-01-26. Retrieved 2012-08-18.
[41] "Everett's Relative-State Formulation of Quantum Mechanics (Stanford Encyclopedia of Philosophy)". Plato.stanford.edu. Retrieved 2012-08-18.
[42] The Transactional Interpretation of Quantum Mechanics by John Cramer. Reviews of Modern Physics 58, 647-688, July (1986)
[43] See, for example, the Feynman Lectures on Physics for some of the technological applications which use quantum mechanics, e.g., transistors (vol III, pp. 14-11 ff),integrated circuits, which are follow-on technology in solid-state physics (vol II, pp. 8-6), and lasers (vol III, pp. 9-13).
[44] Introduction to Quantum Mechanics with Applications to Chemistry - Linus Pauling, E. Bright Wilson. Books.google.com. 1985-03-01. ISBN 9780486648712. Retrieved 2012-08-18.
[45] Anderson, Mark (2009-01-13). "Is Quantum Mechanics Controlling Your Thoughts? | Subatomic Particles". DISCOVER Magazine. Retrieved 2012-08-18.
[46] "Quantum mechanics boosts photosynthesis". physicsworld.com. Retrieved 2010-10-23.
[47] Davies, P. C. W.; Betts, David S. (1984). Quantum Mechanics, Second edition. Chapman and Hall. p. 79. ISBN 0-7487-4446-0., Chapter 6, p. 79
[48] Baofu, Peter (2007-12-31). The Future of Complexity: Conceiving a Better Way to Understand Order and Chaos. Books.google.com. ISBN 9789812708991. Retrieved 2012-08-18.
[49] Derivation of particle in a box, chemistry.tidalswan.com
[50] Malin, Shimon (2012). Nature Loves to Hide: Quantum Physics and the Nature of Reality, a Western Perspective (Revised ed.). World Scientific. ISBN 978-981-4324-57-1.
[51] Chester, Marvin (1987) Primer of Quantum Mechanics. John Wiley. ISBN 0-486-42878-8
[52] Richard Feynman, 1985. QED: The Strange Theory of Light and Matter, Princeton University Press. ISBN 0-691-08388-6. Four elementary lectures on quantum electrodynamics andquantum field theory, yet containing many insights for the expert.
[53] Ghirardi, GianCarlo, 2004. Sneaking a Look at God's Cards, Gerald Malsbary, trans. Princeton Univ. Press. The most technical of the works cited here. Passages using algebra, trigonometry, and bra-ket notation can be passed over on a first reading.
[54] N. David Mermin, 1990, "Spooky actions at a distance: mysteries of the QT" in his Boojums all the way through. Cambridge University Press: 110-76.
[55] Victor Stenger, 2000. Timeless Reality: Symmetry, Simplicity, and Multiple Universes. Buffalo NY: Prometheus Books. Chpts. 5-8. Includes cosmological and philosophical considerations.
[56] Bryce DeWitt, R. Neill Graham, eds., 1973. The Many-Worlds Interpretation of Quantum Mechanics, Princeton Series in Physics, Princeton University Press. ISBN 0-691-08131-X
[57] Dirac, P. A. M. (1930). The Principles of Quantum Mechanics. ISBN 0-19-852011-5. The beginning chapters make up a very clear and comprehensible introduction.
[58] Hugh Everett, 1957, "Relative State Formulation of Quantum Mechanics," Reviews of Modern Physics 29: 454-62.

[59] Feynman, Richard P.; Leighton, Robert B.; Sands, Matthew (1965). The Feynman Lectures on Physics. 1-3. Addison-Wesley. ISBN 0-7382-0008-5.
[60] Griffiths, David J. (2004). Introduction to Quantum Mechanics (2nd ed.). Prentice Hall. ISBN 0-13-111892-7. OCLC 40251748. A standard undergraduate text.
[61] Max Jammer, 1966. The Conceptual Development of Quantum Mechanics. McGraw Hill.
[62] Hagen Kleinert, 2004. Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets, 3rd ed. Singapore: World Scientific. Draft of 4th edition.
[63] Gunther Ludwig, 1968. Wave Mechanics. London: Pergamon Press. ISBN 0-08-203204-1
[64] George Mackey (2004). The mathematical foundations of quantum mechanics. Dover Publications. ISBN 0-486-43517-2.
[65] Albert Messiah, 1966. Quantum Mechanics (Vol. I), English translation from French by G. M. Temmer. North Holland, John Wiley & Sons. Cf. chpt. IV, section III.
[66] Omnès, Roland (1999). Understanding Quantum Mechanics. Princeton University Press. ISBN 0-691-00435-8. OCLC 39849482.
[67] Scerri, Eric R., 2006. The Periodic Table: Its Story and Its Significance. Oxford University Press. Considers the extent to which chemistry and the periodic system have been reduced to quantum mechanics. ISBN 0-19-530573-6
[68] Transnational College of Lex (1996). What is Quantum Mechanics? A Physics Adventure. Language Research Foundation, Boston. ISBN 0-9643504-1-6. OCLC 34661512.
[69] Von Neumann, John (1955). Mathematical Foundations of Quantum Mechanics. Princeton University Press. ISBN 0-691-02893-1.
[70] Hermann Weyl, 1950. The Theory of Groups and Quantum Mechanics, Dover Publications.
[71] D. Greenberger, K. Hentschel, F. Weinert, eds., 2009. Compendium of quantum physics, Concepts, experiments, history and philosophy, Springer-Verlag, Berlin, Heidelberg.
[72] Torben Smith Sørensen (1999). Surface chemistry and electrochemistry of membranes. CRC Press. p. 134. ISBN 0-8247-1922-0.
[73] a b Atkins, Peter and Friedman, Ronald (2005). Molecular Quantum Mechanics, p. 249. Oxford University Press, New York. ISBN 0-19-927498-3.
[74] Atkins, Peter and Friedman, Ronald (2005). Molecular Quantum Mechanics, p. 342. Oxford University Press, New York. ISBN 0-19-927498-3.
[75] Landau, L. D. and Lifshitz, E. M. (1980). Statistical Physics, 3rd Ed. p. 52. Elsevier Butterworth Heinemann, New York. ISBN 0-7506-3372-7.
[76] a b Hill, Terrell L. (1986). Introduction to Statistical Thermodynamics, p. 1. Dover Publications, New York. ISBN 0-486-65242-4.
[77] Schmidt, Lanny D. (2005). The Engineering of Chemical Reactions, 2nd Ed. p. 30. Oxford University Press, New York. ISBN 0-19-516925-5.
[78] Schmidt, Lanny D. (2005). The Engineering of Chemical Reactions, 2nd Ed. p. 25, 32. Oxford University Press, New York. ISBN 0-19-516925-5.
[79] Chandler, David (1987). Introduction to Modern Statistical Mechanics, p. 54. Oxford University Press, New York. ISBN 978-0-19-504277-1.
[80] Alexander Vucinich (1963). Science in Russian culture. Stanford University Press. p. 388. ISBN 0-8047-0738-3.
[81] Josiah Willard Gibbs, 1876, "On the Equilibrium of Heterogeneous Substances", Transactions of the Connecticut Academy of Sciences
[82] Laidler, Keith (1993). The World of Physical Chemistry. Oxford: Oxford University Press. pp. 48. ISBN 0-19-855919-4.
[83] Herbst, Eric (May 12, 2005). "Chemistry of Star-Forming Regions". Journal of Physical Chemistry A 109 (18): 4017–4029. doi:10.1021/jp050461c. PMID 16833724.

Authors

Dr. K. N. Prasanna Kumar has three doctorates one each in Mathematics, Economics and Political Science. Thesis was based on Mathematical Modeling. He was awarded D.litt. For his work on 'Mathematical Models in Political Science' Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India **drknpkumar@gmail.com**

Prof. B. S. Kiranagi is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150 publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent publication history.--UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

Prof. C. S. Bagewadi is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and he has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. He coauthors several Books on Differential Geometry, Differential Equations. Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India