

Assessing the Yield Performance of Improved and Local Potato (*Solanum tuberosum* L.) Cultivars Grown in Eastern Ethiopia

Helen Teshome^{1*} Nigussie Dechassa² Tekalign Tsegaw²

1. Department of Horticulture, College of Agricultural Sciences, Wolaita Sodo University, Ethiopia

2. Department of Plant Sciences, College of Agriculture, Haramaya University, Ethiopia

*E-mail: helenteshome@yahoo.com, P.O.Box 138

Abstract

Potato is an important cash and food security crop in Eastern Ethiopia. However, the yield of the crop is very low due to a number of constraints. Cultivation of low yielding local cultivars is one of the constraints. Therefore, field experiments were conducted at Haramaya and Hirna with the objective of assessing the yield performances of local (farmers') and improved potato cultivars during the main cropping season of 2011. The treatments consisted of five released cultivars (Badhasa, Chala, Chiro, Gabbisa, Zemen) and four local (farmers') potato cultivars (Batte, Daddafa, Jarso, Mashenadima). The experiment was laid out as a Randomized Complete Block Design with three replications. The highest total tuber yield was produced by cultivar Badhasa (37.39 t/ha) while the lowest by Mashenadima (19.46 t/ha) at Hirna. In case of tuber number, Jarso produced (24.37) highest tubers per hill and Mashenadima (7.04) least tubers per hill. Minimum average tuber weight was produced by cultivar Jarso (31.37 g) and maximum by Mashenadima (88.93 g). Zemen at Haramaya had higher (86.60%) harvest index and least was obtained from Daddafa (70.10%) at Hirna.

Keywords: potato (*Solanum tuberosum* L.), yield Performance, Improved, local, Ethiopia

Introduction

Potato (*Solanum tuberosum* L.) is one of the most important food crops in many countries of the world. In volume of production, it is the fourth most important crop after wheat, maize and rice with annual production of 314.1 million tonnes cultivated on about 18.1 million hectares of land (FAOSTAT, 2010). It is a major part of the diet of half a billion consumers in the developing countries (Mondal, 2003). Potato is an important food and cash crop in Eastern and Central Africa, playing a major role in national food security and nutrition, poverty alleviation and income generation, and; provides employment in the production, processing and marketing sub-sectors (Lung'aho *et al.*, 2007).

The low acreage and productivity of potato in Ethiopia are attributed to many factors. The major ones are lack of well adapted and high-yielding cultivars, unavailability and high cost of seed tubers, inappropriate agronomic practices, and lack of marketing and suitable post-harvest management facilities, pests and disease (Berga *et al.*, 1994; Tekalign, 2005; Endale *et al.*, 2008; Gildemacher *et al.*, 2009).

In Eastern Ethiopia, improving and maintaining potato cultivars with high yield and market values is limited by lack of knowledge on the diversity of the local as well as the improved cultivars. This is because; the local cultivars are not well studied as that of the improved ones, which need studying both cultivars for better documentation. Moreover, considerable variations exist in local naming or identification of the cultivars. There is information that each agro-ecological zone has its own unique set of names for different cultivars and duplicate or different names may have been given for the same cultivar. This diverse system of naming cultivars not only limits the proper identity of the cultivar but also hinders monitoring and follow-up of the released improved cultivars. Therefore, documented information on the cultivated potato cultivars is vital for advancing improvement work on the crops.

However, to date, no systematic studies have been done to investigate and document the similarities and differences yield performances among the local and improved potato cultivars grown in the Eastern Ethiopia. Therefore, this study was initiated with the objective of assessing the yield performances of the major local and improved cultivars of potato grown in Eastern Ethiopia.

Materials and Methods

Description of the study area

The study were carried out at Rare, Horticulture section's research field, Haramaya University and Hirna research site of the University under rainfed condition during the 2011 main growing season. Rare research site is located at 9°26' N latitude, 42°03' E longitudes at an altitude of 1980 m.a.s.l. The mean annual rainfall is 760 mm (Belay *et al.*, 1998). Mean annual temperature 16 °C (Mishra *et al.*, 2004). The mean relative humidity is 50%, varying from 20 to 81%. The soil of the experimental site is alluvial type with organic carbon content of 1.15%, total Nitrogen content of 0.11%, available Phosphorus content of 18.2 mg kg soil⁻¹, exchangeable Potassium content of 0.65 cmol_c kg soil⁻¹, pH of 8.0 and per cent sand, silt and clay content of 62.92, 19.64 and 17.44, respectively (Simret, 2010).

Hirna sub-station is located at 9 °12' N latitude, 41 °4' E longitudes at an altitude of 1870 m. a.s.l. The area receives mean annual rainfall of 990 to 1010 mm with an average temperature of 24 °C (HURC, 1996). The soil of Hirna is vertisol with organic carbon content of 1.75%, total Nitrogen content of 0.18%, available Phosphorus content of 32 mg kg soil⁻¹, exchangeable Potassium content of 0.68 cmol_c kg soil⁻¹, pH of 7.09 and percent sand, silt and clay contents of 27, 28 and 45, respectively (Nebret, 2011).

Description of Experimental Material

Five potato cultivars, which were released by Haramaya University at different times and four locally available potato cultivars were used for the experiment (Table 1).

Table 1. Potato cultivars used in the study, year of release and their sources.

Variety	Year of release	Source of planting material
Chiro	1998	HUPIP
Zemen	2001	HUPIP
Badhasa	2001	HUPIP
Gabbisa	2005	HUPIP
Chala	2005	HUPIP
Source (MoARD, 2010)		
Batte	Local	RHSPC
Mashenadima	Local	RHSPC
Jarso	Local	RHSPC
Daddafa	Local	RHSPC

Key: HUPIP = Haramaya University Potato Improvement Programme

RHSPC = Rare Hora Seed Producers' Cooperative

Treatments and Experimental Design

The treatments are nine consisting of five improved cultivars (Chala, Chiro, Badhasa, Gabbisa and Zemen) and four local cultivars (Batte, Mashenadima, Jarso and Daddafa). The experiment was laid out as a Randomized Complete Block Design (RCBD) and replicated three times. Each plot was 3.60 m x 4.50 m = 16.2 m² wide consisting of six rows, which accommodated twelve plants per row and thus 72 plants per plot. The spacing between plots and adjacent replication were 1 m and 1.5 m, respectively. There was a total of 669.3 m² area for experimental site.

Experimental Procedures

The experimental field was cultivated by a tractor to a depth of 25-30 cm and levelled and ridges were made by hand. Medium sized (39-75 g) Lung'aho *et al.*, (2007) and well sprouted tubers were planted at the sides of ridges at the spacing of 75 cm between ridges and 30 cm between tubers. Planting depth was maintained at 5 cm (Mahmood *et al.*, 2001). Phosphorus fertilizer at the rate of 92 kg P₂O₅ ha⁻¹ in the form of Diammonium Phosphate (200 kg ha⁻¹) was used and the whole rate was applied at planting. 75 kg Nitrogen ha⁻¹ was applied in the form of urea in two splits, half rate after full emergence (two weeks after planting) and half rate at the initiation of tubers (start of flowering). Potato plants were treated with Mancozeb 80% WP at the rate of 1.5 kg ha⁻¹ diluted at the rate of 40 g per 20 litre water once a week to control late blight disease.

Plant Data Collection and Analysis

Post harvest observations and measurement were taken from randomly selected plants from each plot for all characters studied. Data were subjected to analysis of variance (ANOVA) using the General Linear Model of the SAS statistical package (SAS, 2007) version 9.1. All significant pairs of treatment means were compared using the Least Significant Difference Test (LSD) at 5% and 1% level of significance.

Results and Discussion

Total tuber yields

Yield is a prime trait which helps in selection of any new cultivars because farmers are more concerned with this trait more than the quality traits. The main effects of both cultivar and location highly significantly ($P < 0.01$) influenced both marketable and total tuber yields. The main effect of location did not significantly influence unmarketable tuber yield. However, cultivar and location interacted to highly significantly ($P < 0.01$) influence marketable, unmarketable and total tuber yields.

The cultivars significantly differed in the ability to produce tuber yields at both locations. Thus, the highest total tuber yields were obtained for Chiro (32.81 t ha⁻¹) and Daddafa (31.53 t ha⁻¹) at Haramaya, which

were in statistical parity. The lowest total tuber yield was obtained for Mashenadima (19.46 t ha⁻¹) at Haramaya. Thus, at Haramaya, the total tuber yield obtained from Chiro significantly exceeded the total tuber yields obtained from Mashenadima, Jarso, Chala, Batte, Gabbisa, Zemen, Badhasa and Daddafa by additional increments of about 69,51,43,41,15,10,8 and 4%, respectively (Table 2).

At Hirna, the highest total tuber yields were recorded for Badhasa (37.39 t ha⁻¹), Chiro (36.57 t ha⁻¹), Zemen (34.19 t ha⁻¹) and Batte (33.13 t ha⁻¹). The four cultivars produced total tuber yields that were in statistical parity with each other as well as with the total tuber yields obtained for Chiro at Haramaya. At this location, Badhasa produced total tuber yields that significantly exceeded the total tuber yields recorded for Jarso, Daddafa, Chala, Mashenadima and Gabbisa by about 50, 46, 35, 28, and 20%, respectively (Table 2).

The results revealed that most of the cultivars differed in yielding ability across the two locations. However, Chiro produced significantly higher total tuber yields at both locations, indicating its high yielding potential as well as yield stability at both locations. The total tuber yields obtained from Gabbisa, Chala and Zemen, although being significantly lower than those obtained from Chiro, remained stable across the two locations. Mashenadima and Badhasa yielded significantly better at Hirna than at Haramaya, with extra total tuber yield advantages of about 50 and 24%, respectively. However, only one cultivar, namely, Daddafa yielded significantly better at Haramaya than at Hirna, with a relative yield advantage amounting to about 34% (Table 2). The results of this study are in accord with those of Wiersema and Cabello (2002) and Tekalign and Hammes (2005) who reported that different potato cultivars varied with respect to total tuber yields.

Marketable tuber yields

Similar to the differences observed in total tuber yields, the cultivars also differed in marketable tuber yields across the two locations (Table 2). The highest marketable tuber yields were recorded for Chiro (30.95 t ha⁻¹), Zemen (30.49 t ha⁻¹), Badhasa (28.49 t ha⁻¹) and Batte (26.49 t ha⁻¹) at Hirna, closely followed by the marketable tuber yields of Chiro, Daddafa, Zemen and Badhasa cultivars at Haramaya. The proportion of marketable tuber yields of the four cultivars viz., Chiro, Daddafa, Zemen and Badhasa to their total tuber yields at Hirna amounted to as high as 85, 89, 76 and 80 %, respectively. This shows that the four cultivars produced more healthy and unblemished tubers than the others. The lowest marketable tuber yields were recorded for Chala, Mashenadima, Batte and Jarso at Haramaya and for Chala at both locations, which amounted to only 51, 62, 45, 74 and 43% of their corresponding total tuber yields, respectively. The higher marketable yields of the other cultivars were indicator of freedom of the tubers from diseases and physical blemishes.

Unmarketable tuber yields

The cultivars significantly differed also in the unmarketable tuber yields. The highest unmarketable tuber yields were recorded for the cultivar Chala at both Hirna (15.77 t ha⁻¹) and Haramaya (11.38 t ha⁻¹) (Table 2) whereas the lowest was recorded for the cultivars Zemen, Daddafa, Jarso and Chiro at Hirna, and Jarso at Haramaya. Chala produced the highest amount of unmarketable tuber yield. The highest unmarketable tuber weight of Chala could be attributed to the bacterial wilt disease (*Ralstonia solanacearum*), which infected the cultivar at both locations. The disease led to rotting of tubers thereby drastically reducing their yield. These bacterial wilt is a devastating disease of potato infecting and reducing tuber marketability drastically was described also by (Lung'aho *et al.*, 2007).

The cultivars that produced low yields such as Mashenadima, Jarso and Batte at Haramaya and Jarso, Daddafa and Mashenadima at Hirna were because these local cultivars were vulnerable to biotic environmental stresses particularly the late blight disease. However, Daddafa at Haramaya and Batte at Hirna performed relatively well. Corroborating this suggestion, Kidane-Mariam (1980) stated that local potato are generally low yielders and are susceptible to diseases and pests. Bekele and Hailu (2001) also stated that late blight caused by *Phytophthora infestans* is the most serious fungal disease of potato causing huge losses in yield. Therefore, application of proper management practices to enhance vigorous early growth is very important to achieve higher tuber yield from potato under these types of environmental conditions (Fotiveark, 1999; Caliskan *et al.*, 2004).

Table 2. The interaction effect of cultivars and location on marketable, unmarketable and total tuber yield of potato cultivars.

Location	Cultivar	Marketable	Unmarketable	Total Tuber yield (t/ha)
Haramaya	Badhasa	20.65 ^{c-f}	9.60 ^{bcd}	30.25 ^{cde}
	Batte	14.80 ^{fg}	8.49 ^{def}	23.29 ^{f-i}
	Chala	11.61 ^g	11.38 ^{bc}	22.99 ^{ghi}
	Jarso	16.19 ^{efg}	5.59 ^{gh}	21.78 ^{hi}
	Chiro	24.76 ^{bcd}	8.05 ^{def}	32.81 ^{a-d}
	Mashenadima	12.13 ^g	7.33 ^{efg}	19.46 ⁱ
	Zemen	21.34 ^{cde}	8.61 ^{def}	29.95 ^{cde}
	Daddafa	22.34 ^{cd}	9.19 ^{cde}	31.53 ^{bcd}
	Gabbisa	20.58 ^{c-f}	8.03 ^{def}	28.61 ^{def}
Hirna	Badhasa	28.49 ^{ab}	8.90 ^{cde}	37.39 ^a
	Batte	26.49 ^{abc}	6.64 ^{fg}	33.13 ^{a-d}
	Chala	11.91 ^g	15.77 ^a	27.68 ^{d-g}
	Jarso	19.31 ^{def}	5.62 ^{gh}	24.93 ^{e-i}
	Chiro	30.95 ^a	5.62 ^{gh}	36.57 ^{ab}
	Mashenadima	19.75 ^{def}	9.46 ^{bcd}	29.21 ^{cde}
	Zemen	30.49 ^{ab}	3.70 ^h	34.19 ^{abc}
	Daddafa	20.22 ^{def}	5.32 ^{gh}	25.54 ^{e-h}
	Gabbisa	20.30 ^{def}	10.84 ^{bc}	31.14 ^{bcd}
LSD (0.05)	6.07	2.08	5.49	
F-test	**	**	**	
CV (%)	17.67	15.25	11.44	

Treatment means followed by the same letter within a column are not significantly different. ** = significant at 1% probability level.

Tuber number

Marketable and total tuber numbers per hill were highly significantly influenced by the main effects of cultivar and location as well as by the interaction of the two factors. However, unmarketable tuber number per hill was influenced only by the main effect of cultivar and the interaction effect of cultivar and location.

Total tuber number

The cultivar Jarso (24.37) produced the highest total tuber number per hill at closely followed by Daddafa, Badhasa, Batte at Hirna and Jarso itself at Haramaya. The total tuber number produced by Jarso at Hirna significantly exceeded the total tuber numbers produced by Badhasa and Daddafa at the same location by about 37 and 36%, respectively. Mashenadima (7.04) at Haramaya produced the least tuber number and was exceeded the total tuber number produced by Jarso by about 187%. This shows that the cultivars markedly differ in the number of tubers they produce. The total tuber numbers produced by each of the cultivars at the two locations were in statistical parity. However, Batte, Gabbisa and Jarso produced significantly higher tuber number per hill at Hirna than at Haramaya (Table 3). This shows that the environment in which cultivar grow may also influence the number of tuber produced per hill. The results of this study are corroborated by the observation of Subarta and Upadhya (1997) that genetic and environmental factors play a vital role in stolon development and tuberization process. Yibekal (1998) also observed a significant difference in total tuber numbers produced between the improved cultivar AI-624 and the local check. Similarly, Allen (1978) also reported that the number of tubers set by plants was determined by stem density, variety, crop management and season.

Marketable tuber number

The cultivars significantly differed in marketable tuber numbers produced per hill. The highest marketable tuber numbers were recorded for Jarso (13.88), Chiro (13.18), Batte (11.91), Badhasa (11.43), Zemen (10.99) and Daddafa (9.88) at Hirna, closely followed by the marketable tuber numbers of Jarso and Daddafa at Haramaya. The lowest marketable tuber number was recorded for Chala (3.47) and Mashenadima (3.73) at Haramaya (Table 3). This shows that cultivars Jarso, Chiro, Batte, Badhasa, Zemen and Daddafa at Hirna were prolific and produced significantly higher, healthier and unblemished tuber numbers than the rest of the cultivars. Except Chala and Mashenadima, which produced marketable tuber numbers, which were in statistical parity at both locations, all other cultivars produced significantly higher marketable tuber numbers at Hirna than at Haramaya. This shows that Hirna is a more conducive environment for profitable potato production during the meher season.

Unmarketable tuber number

Similar to the differences observed in total tuber number and marketable tuber number, the cultivars also differed in unmarketable tuber numbers at both locations (Table 3). The highest unmarketable tuber number were recorded for cultivar Jarso (10.49) at Hirna and Badhasa (9.42) at Haramaya. However, the lowest unmarketable tuber number was recorded for cultivar Zemen (2.92) and Chiro (3.71) at Hirna closely followed by Mashenadima (farmers' cultivar) at Haramaya and Hirna, respectively.

Across the two locations, some cultivars produced significantly higher unmarketable tubers at Hirna than at Haramaya (Batte and Jarso). Others produced higher unmarketable tubers at Haramaya than at Hirna (Badhasa, Chiro and Zemen). The remaining cultivars all produced tuber numbers that were in statistical parity at both locations (Table 3).

This result indicates that number of tubers initiated by a particular cultivar could be influenced by the growing condition as well as the cultivars' inherent traits. Consistent with this suggestion, Subarta and Upadhyha (1997) reported that genetic and environmental factors play a vital role in stolon development and tuberization process. The highest unmarketable and total tuber number of Jarso could be related to late blight disease because of the inherent vulnerability of the cultivar to the disease. Consistent with this suggestion, Bekele and Hailu (2001) stated that Late blight is the major destructive and the most serious fungal disease of potato that affects tuber production

Table 3. The interaction effect of cultivar and location on marketable, unmarketable, total tuber number and Harvest index of potato cultivars.

Location	Cultivar	Tuber number per hill			Harvest index(%)
		Marketable	Unmarketable	Total	
Haramaya	Badhasa	6.37 ^{ij}	9.42 ^{ab}	15.79 ^{b-c}	75.86 ^{hig}
	Batte	7.03 ^{ghi}	3.35 ^{gh}	10.38 ^{hi}	73.34 ^{hij}
	Chala	3.47 ^k	7.13 ^{cd}	10.60 ^{hi}	76.84 ^{f-i}
	Jarso	9.74 ^{def}	7.10 ^{cd}	16.84 ^{bcd}	77.63 ^{e-h}
	Chiro	8.60 ^{fgh}	5.93 ^{de}	14.53 ^{def}	81.35 ^{b-f}
	Mashenadima	3.73 ^k	3.31 ^{fgh}	7.04 ^j	72.54 ^{ij}
	Zemen	6.70 ^{hij}	7.06 ^{cd}	13.76 ^{efg}	86.60 ^a
	Daddafa	8.87 ^{efg}	7.03 ^{cd}	15.90 ^{b-c}	75.97 ^{hij}
	Gabbisa	6.75 ^{g-j}	4.88 ^{efg}	11.63 ^{gh}	83.32 ^{a-d}
Hirna	Badhasa	11.43 ^{bcd}	6.31 ^{cde}	17.74 ^{bc}	82.06 ^{a-c}
	Batte	11.91 ^{abc}	5.27 ^{def}	17.18 ^{bcd}	79.55 ^{c-g}
	Chala	4.79 ^{kj}	8.06 ^{bc}	12.85 ^{fgh}	78.44 ^{d-g}
	Jarso	13.88 ^a	10.49 ^a	24.37 ^a	84.57 ^{ab}
	Chiro	13.18 ^{ab}	3.71 ^{fgh}	16.89 ^{bcd}	84.34 ^{abc}
	Mashenadima	3.75 ^k	4.50 ^{e-h}	8.25 ^{ij}	78.37 ^{efg}
	Zemen	10.99 ^{cde}	2.92 ^h	13.91 ^{efg}	79.48 ^{c-g}
	Daddafa	9.88 ^{c-f}	8.09 ^{bc}	17.97 ^b	70.10 ^j
	Gabbisa	8.84 ^{fg}	6.16 ^{de}	15.00 ^{c-f}	85.72 ^{ab}
LSD(0.05)	2.14	1.87	2.85	4.90	
F-test	**	**	**	**	
CV (%)	15.46	18.31	11.87	3.73	

Treatment means followed by the same letter within a column are not significantly different. ** = significant at 1% probability level.

Harvest index

The main as well as the interaction effects of location and cultivars significantly ($P < 0.01$) influenced harvest index. Zemen (86.60%) at Haramaya followed by Gabbisa, Jarso, Chiro and Badhasa at Hirna produced higher harvest indices that were similar and in statistical parity. The lowest harvest index was recorded for Daddafa (70.10%) at Hirna. On average, cultivar Zemen at Haramaya exceeded cultivar Daddafa by about 24%(table 3). As Radford, (1967) suggested this characterizes the allocation of dry matter to the economic part of the plant. The variations observed among the cultivars in harvest index may be attributed to their inherent genetic differences as well as the growing locations, which is evident from the significant interaction effect.

Table 4. The main effect of cultivars and location on average tuber mass.

	Average tuber mass (g/tuber)
Location	
Haramaya	69.53 ^a
Hirna	47.97 ^b
LSD(0.05)	6.28
F-test	**
Cultivars	
Badhasa	55.43 ^b
Batte	58.87 ^b
Chala	63.40 ^b
Jarso	31.37 ^c
Chiro	57.95 ^b
Mashenadima	88.93 ^a
Zemen	60.85 ^b
Daddafa	50.67 ^b
Gabbisa	61.32 ^b
LSD(0.05)	13.29
F-test	**
CV (%)	19.3

Average tuber weight

The average tuber weight was highly significantly ($P < 0.01$) influenced by cultivar and location. However, this parameter was not significantly influenced by the interaction effect of cultivar and location. The cultivar Mashenadima (88.93g/tuber) had significantly higher average tuber weight than the average tuber weights of all other cultivars. On the other hand, Jarso (31.37 g/tuber) produced the least average tuber weight. The remaining cultivars had average tubers weights that were in statistical parity with each other (Table 4). The average tuber weight of Mashenadima exceeded the average tuber weights of Chala, Gabbisa, Zemen, Batte, Chiro, Badhasa, Daddafa and Jarso by additional increment of about 40, 44, 46, 51, 53, 60, 75 and 183%, in the order listed here (Table 4). The tuber weight of most of the cultivars was in the medium category. The results of this study are in accord with those of Berga *et al.* (1994) and Bereke (1994) who observed significant differences between varieties for average tuber weight.

The results also revealed that heavier tubers were generally produced at Haramaya whereas larger numbers were produced at Hirna, indicating that tuber weight and numbers are inversely related and also influenced by environmental factors. In line with this result, Boyd *et al.* (2002); Struik (2007); Eremeev *et al.* (2008) also stated that the greater the number of tubers per plant, the smaller the average tuber mass, and vice versa.

Conclusion

The highest mean total and marketable tuber yields of about 30 t ha⁻¹ and 25 t ha⁻¹, respectively over both locations were recorded for the improved cultivars released by the Haramaya University, namely, Badhasa, Chiro, Zemen and Gabbisa. Among the farmers' cultivars, it was only Daddafa that competed with the cultivars released by the Haramaya University in terms of yielding ability by producing about 27 t ha⁻¹ of total and about 21 t ha⁻¹ of marketable mean tuber yields at both locations. However, cultivar Batte performed comparatively well like Daddafa, but only at Hirna and not at Haramaya. These results clearly indicate that the cultivars released by Haramaya University performed far better than the majority of farmers' cultivars.

Reference

- Allen, E.J., 1978. Plant density. pp. 279-324 In: P.M. Harris, (ed.). The Potato Crop: The Scientific Basis for Improvement of the Potato crop.
- Belay, S.C., W. Wortman and G. Hoogen boom, 1998. Haricot bean agro-ecology in Ethiopia: definition using agro-climatic and crop growth stimulation models. *African Crop Science Journal*. 6: 9-18.
- Bekele Kassa and Hailu Beyene, 2001. Efficacy and economics of fungicide spray in the control of late blight of potato in Ethiopia Ethiopian Agricultural Research Organization Holetta Agricultural Research Center. *African Crop Science Journal*. 9(1): 245-250.
- Bereke Tsehail, 1994. The utilization of true potato seed (TPS) as an alternative method of potato production. A

- Ph.D. Dissertation Presented to the Wageningen University, The Netherlands. 289p.
- Berga Lemaga, G. Hailemariam and G. Woldegiorgis, 1994. Prospects of seed potato production in Ethiopia. pp. 254-275. In: E. Hareth, D. Lemma (ed.). Proceedings of the Second National Horticultural Workshop of Ethiopia. EARO-FAO, Addis Ababa, Ethiopia
- Boyd, N.S., R. Gordon and R.C. Martin, 2002. Relationship between leaf area index and ground cover in potato under different management conditions. *Potato Research*. 45(2/4): 117-129.
- Caliskan, M.E., S. Caliskan and H. Arioglu, 2004. Effects of Pre-sprouting and Planting Date on Growth and Yield of Potato Crop in a Mediterranean Type Environment. *Annals of Botany*. 45: 34-47
- CSA (Central Statistical Agency), 2008/2009. Agricultural sample survey: Report on area and production of crops, Addis Ababa, Ethiopia. 126p.
- Endale Gebre, G. woldegeorgis and B. Lemaga, 2008. Potato seed management. pp. 53-78. In Root and tuber crops. *Ethiopian Institute of Agricultural Research*.
- Eremeev, V., A. Lohmus, P. Laaniste, J. Joudu, L. Talgre and E. Lauringson, 2008. Influence of seed potato pre-planting treatments on leaf area and tuber yield formation. *Annual Review of Plant Physiology*. 39: 355-378.
- FAO, 2008. Potato World: Africa-International Year of the Potato 2008. <http://www.potato2008.org/en/world/africa.html>.
- FAOSTAT, 2010. Agricultural data. Provisional 2009 Production and Production Indices Data. Crop primary. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Fotiveark, S., 1999. Early potatoes in Italy with particular reference to Sicily. *Potato Research*. 42: 229-240.
- Gildemacher, P.R., W. Kaguongo, O. Ortiz, A. Tesfaye, W. Gebremedhin, W.W. Wagoire, R. Kakuhenzire, P.M. Kinyae, M. Nyongesa, P.C. Struik and C. Leeuwis, 2009. Improving Potato Production in Kenya, Uganda and Ethiopia: A System Diagnosis. *Potato Research* 52: 173-205.
- HURC (Haramaya University Research Centre), 1996. Proceedings of the 13th annual Research and extension review meeting. pp. 26-28.
- Kidane-Mariam, H.M., 1980. Project Proposal for the Development of an Ethiopian Potato Program. Addis Ababa. Manuscript.
- Lung'aho, C., B. Lemaga, M. Nyongesa, P. Gildemacher, P. Kinyale, P. Demo and J. Kabira, 2007. Commercial seed potato production in eastern and central Africa. Kenya Agricultural Institute, Kenya. 140p.
- Mahmood, M.M., A. Hussain and K. Farooq, 2001. "Aallo Ki Kasht" 9 p. National Potato Programme, NARC. 59p.
- Mishra B.B., H.G. Kidan, K. Kibret, M. Assen and B. Eshetu, 2004. Soil and land resource inventory at Alemaya University research farm with reference to land evaluation for sustainable agricultural management and production: Synthesis of working papers, *Soil Science Bulletin*. Alemaya University, Ethiopia.
- MoARD (Ministry of Agriculture and Rural development), 2010. Animal and Plant Health Regulatory Directorate. Crop Variety Register. Crop Development Department, Issue No.13 June 2010, Addis Ababa, Ethiopia. 138p.
- Mondal, M.A.A., 2003. Improvement of potato (*Solanum tuberosum* L.) through hybridization and *in vitro* culture technique. A. Ph.D. Dissertation presented to Rajshahi University, Rajshahi, Bangladesh. 270p.
- Nebret Tadesse, 2011. The effect of Nitrogen and Sulfer on yield and yield component of common bean in Eastern Ethiopia. Unpublished M.Sc. Thesis presented to the school of graduate studies of Haramaya University. 13p.
- Radford, P.J., 1967. Growth analysis formulae-their use and abuse. *Crop Sciences Journal*. 7: 171-175.
- SAS (Statistical Analysis Software), 2007. Stat. Jahrbuch tuber Ernährung, Landwirtschaft und Forsten In German, Landwirtschaftsverlag Munster-Hiltrup, Germany, 2008.
- Simret Burga, 2010. Influence of inorganic nitrogen and potassium fertilizers on seed tuber yield and size distribution of potato (*Solanum tuberosum* L.). An. M. Sc Thesis Presented to the School of Graduate Studies of Haramaya University, Ethiopia. 65p.
- Struik, P.C. 2007. Response to the environment: temperature pp. 367-391. In: D. Vreugdenhill Bradshaw, J. Elsevier (eds.). Potato Biology and Biotechnology. Advances and Perspectives. Amsterdam, The Netherlands.
- Subarta, M. and M.O. Upadhya, 1997. Potato production in western Bengal. *Environmental. Ecology Journal*. 15: 646-900.
- Tekalign Tsegaw and P.S. Hammes, 2005. Growth and productivity of potato as influenced by cultivar and reproductive growth: II. Growth analysis, tuber yield and quality. *Scientia Horticulture Journal*. 105(1): 29-44.
- Tekalign Tsegaw, 2005. Response of potato to Paclobutrazol and Manipulation of Reproductive Growth under Tropical Conditions. A Ph.D. Dissertation presented to the department of production and soil science. University of Pretoria. 164p.

- Wiersema, S.G. and R. Cabello, 2002. Effect of potato mini tubers on the yield and production. Department of Physiology, Int. Potato Center, Lima, Peru. <http://www.springerlink.com/content/lru3208p21604732>.
- Yibekal Alemayehu, 1998. The effect of nitrogen and phosphorus on yield and yield components, and some quality traits of potato (*Solanum tuberosum* L.) grown on soils of Wondo Genet area. An M.Sc. Thesis presented to the school of graduate studies of Haramaya University.84p.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:
<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

