

Some Herbal Medicinal Plants Activity against *Candida* spp which Resistance to Antifungal Drugs

Mona Al-Terehi¹ Ali H. Al-Saadi² Zahraa Abed-Neama² Mohammed Al-Askeri³
Haider K.Zaidan² Rajaa Ali Habeeb² Noora Majed² Zahraa Haleem²

1. University of Kufa -College of Science

2. University of Babylon, College of Science

3. AL-Qadesia university

Abstract

Some medicinal plant that used as antibacterial in Iraq has been experimented as antifungal, five of these plants include *Zingiber officinale*, *Salvia officinalis*, *Origanum vulgare*, *Glycyrrhiza glabra*, *Punica granatum* were used against four species of *candida* spp. *C. paracitucus*, *C. albicans*, *C. tropicalis*, *C. kruzi*, which are resistance to antifungal drugs, 80% methanol was used to extract these plants, 5 mg/ml of each extracts used in nutrient broth for 4 days to evaluated anti-candida activity. The Results show that plants extracts causes complete inhibition, decreased in *candida* growth, don't effect on activated candida growth, *Zingiber officinale* causes complete inhibition to *candida* spp except *C. albicans*, *Salvia officinalis* was decreased in all *candida* spp, *Origanum vulgare*, effect on *C. albicans* and *C. kruzi* but it don't effect on other species. *Glycyrrhiza glabra* decreased all *candida* spp. *Punica granatum* decreased growth of *C. albicans* and *C. kruzi* only.

Keywords: *Candida* spp, plant extract, 80% methanol.

Introduction

Medicine plant has been used in different application especially to treated different pathogenesis of microorganisms in the world, in Iraq some of these organisms have resistance to wide spectrum of drugs thus it causes different infection and complication, the pathogenic include yeast such as *candida* is resistance to several drugs like cycloheximide, fluconazole, nystatin and gresofulvin, this resistance was increased in recent years because of several mechanisms development like genetic changes in some genes via mutation that causes changes in some enzyme pathway which leads to lesion in enzyme pathway or in multidrug protein transport which responsible of drug transport in side cell such as CDR1, CDR2 and CaMDR1 which have been played role in fluconazole resistance (Albertson *et al.*, 1996).

Studies are dealing with medicinal plants mechanism of action on microorganism suggested that these mechanisms may be affected on cell membrane which causes increased membrane preimpility associated with loss ions and reduction in membrane potentiality (Di Pasqua *et al.*, 2006; Turina *et al.*, 2006)

The disruption of the cell membrane causes defect in biological activity like energy conversion processes, nutrient processing, synthesis of structural macromolecules, and secretion of many growth regulators (Oussalah *et al.*, 2006).

Turina *et al.*, (2006) affirm the effectiveness of specific ions on plasma membrane has effective on the different process such as protons motive force, intracellular ATP content and overall activity of microbial cells like turgor pressure, solutes transport and metabolism regulation process.

Essential oils of many plants are composed of terpenes, terpenoids and other aromatic, aliphatic constituents can be penetrate and damage fungal cell wall and cytoplasmic membranes, permeable them and finally disrupted mitochondrial membranes, Changes in electron flow through the electron transport system in mitochondria lead to lipids, proteins and nucleic acid damage (Arnal-Schnebel *et al.*, 2004). The essential oils can hassle the depolarization of the mitochondrial membranes and decreased membrane potential, also affect Ca^{2+} and other ion channels, lowering pH and effect on the proton pump, ATP pool. changes in the fluidity of membranes resulted into the radicals leakage, cytochrome C, calcium ions and proteins. Thus, permeabilization of outer and inner mitochondrial membranes leads to cell death by apoptosis and necrosis (Yoon *et al.*, 2000).

Present study was suggested for using common plants that used in different medical application in Iraq to treated *candida* spp infection which is resistance to antifungal drugs, also these plants extracts don't have harmful side effect if it used under Specialists. Review of literature improved its ability to treated different infections such as pathogenic bacteria, fungal and viral because it contain phytochemicals compounds that have different mechanism to decreased infection or killed microorganism cells. Plants extract which used in present study choosing according to its ability to treated infection and low side effects and no cytotoxicity of its, also methods of extraction improved its ability to extract most of phytochemicals compounds from part of plants which used in present study. (Al-Saadi *et al.*, 2012, Al-Terehi *et al.*, 2012)

Materials and methods

1- Pathogenesis *Candida* spp : *candida* spp was isolation from (vaginal, mouth and urine then it diagnosis using macroscopic, microscopic and biochemical tests.

- 2- Antifungal susceptibility L these tests performed using Nystasin, Terbinafine, Griseofulvin and Fluconazole using disc diffusion method.
- 3- **Plants extract:** The plants extracts of plant was prepared according to the method of Sato *et al.* (1990) with some modification. Specific weight of the plant and it is mixed with the average 1 gm. to 3 ml of the solvent solution (20 %methanol: 80 % distilled water), the mixture is uniformed by electric blender for 30 minutes in room temperature. The solution is filtered by using gauze fabric for getting transudate solution. It is deride using incubator at 50 C° for 24 hours,
- 4- **Plant extract aliquot,** it prepares in concentration (0.1 g\ml) then it sterilized using mellipor paper 0.2 mm.
- 5- **Anti-candida activity:** this was performed using 50 mg/ml of plant extract, growth evaluation by optical density of growth culture according to Jabor *et al* (2013)

RESULTS

The results show that these extracts have anti-candida activity , these activity was differences between Candida types and types of plant extract as show in table(1) and figure (1)

Table (1) show plants extract activity against candida spp. *Zingiber officinale* was more effect than others plant extract it causes completed inhibition to *C. paracitucus*, *C.tropicus*, *C.kruzi*. Other plant extracts cause decreased in growth but these decreasing was Disparity from low to semi-complete inhibition.

In another hand some plants extract causes activate candida growth, *Punica granutum* with *C. paracitucus* and *C.tropicus*, and *Origanum vulgare* with *C.tropicus*. Other plants extract don't effect on *Candida spp.* growth *Origanum vulgare* with *C. paracitucus* as show in table (1).

Significance at (p > 0.05) was show in complete inhibition of growth in *C. paracitucus*, *C.tropicus* by *Zingiber officinale* and *C.kruzi* by *glycyrrhiza glabra* and *Punica granutum*

Table (1) Plants extracts activity against *Candida* spp.

candida Plant extract	<i>C. paracitucus</i>	<i>C. albicans</i>	<i>C.tropicus</i>	<i>C.kruzi</i>
Positive control	1.02±0.012	0.545±0.012	0.560±0.056	0.806±0.096
<i>Zingiber officinale</i>	0*	0.433±0.227	0*	0*
<i>Salvia officinalis</i>	0.157±0.157	0.355±0.355	0.433±0.433	0.283±0.283
<i>Origanum vulgare</i>	1.01±0.26	0	1.02±0.487	0.30±0.153
<i>glycyrrhiza glabra</i>	0.50±0.176	0.153±0.153	0.176±0.121	0.07±0.072*
<i>Punica granutum</i>	1.187±0.11	0.230±0.230	0.580±0.253	0.243±0.243*





Figure (1) Effect of plants extract types on *Candida* spp. A, *Candida paracitucus*; B, *C. albicans*; C, *C. tropicus* ; E, *C.kruzi*; 1, positive control ; 2, *Punica granutum*; 3, *Salvia officinalis*; 4, *Zingiber officinale*; 5, *Origanum vulgare*; 6, *glycyrrhiza glabra*.

Discussion

Results of present study show that plants extract have ability to decreased *candida* spp growth, complet inhibition, don't effect and activate candida growth, as show in table (1) Zingiber was most plant effect on *candida* spp it improved its ability against different pathogenic microorganism, Supreetha *et al* (2011) used ethanol to extract of ginger against *candida albicans* in vitro using disc diffusion methods at 24 and 48 hours they found that this extract have antifungal activity at 24 hours. Also the previous study recorded that ginger have different phytochemicals compounds which have antifungal antibacterial activity because it effect on different factors such as pH, nutrient ion, osmosis pressure and microenvironment of microorganisms, this compound may be causes disrupted of cellular activity of microorganism that lead to decrease of growth or killing it by oxidative damages (Hayes *et al* 2013) .

Origanum vulgare also inhibited *C. spp* growth in different level , it causes completed inhibition of *C albicans*, Vale-Silva and others (2011) clarified the mechanism of action of this inhibition using flow cytometry to tested different parts of *Origanum vulgare* phytochemicals compounds effect on cell membrane by direct effect on cell membrane lipids, thus it consider as fungicide for treated superficial infection. Essential oil of *Origanum vulgare* have significant inhibitory effect on *C. albicans* and *C. kruzi* using solid media and disc diffusion method (Souza *et al* ., 2007).

Liquors show decreased candida growth the review of literature recorded inhibition activity of different extract of licorice root against pathogenic bacteria and fungi Khanuja, K. (2006), this activity may be its phytochemical compound that interacted with microorganisms Al-Saadi *et al* (2012). *Salvia officinalis* oil was inhibitor pathogenesis *candida* spp using 15.6, 3.9, 31.3, 31.3 and 1.9 µg/m of its oil in Iran (Badiee *et al.*, 2012).

In Turkey Dulger and Hacıoglu (2008) used ethanol extract of sage also have anti-candida ability, sage extract contain from α -pinene, β -pinene, β -thujone, camphor, carvacrol, linalyl acetate ,sabinyl acetate and 1,8-cineole these extracts may be have anti-microbial activity which were improved by previous study .

Most of research using oil as antifungal thus methanol –water extract was used in present study, results improved efficacy against *candida* spp.

Pomegranate has been used as anti-bacterial and anti-fungal activity, other study used different parts of Pomegranate in extract using methanol, ethanol and acetone to extract phytochemicals, (Dahham *et al* 2010) used it as antifungal against *Aspergillus niger* growth.

Ahmed and Beg (2001) reported that a most of plants extracts methods including ethanol extracts of pomegranate showed antifungal activity against *Candida albicans*. In vitro studies have revealed that the extract of pomegranate inhibited the growth of oral bacteria and *candida* species. The present study also improved that used plant extract may be causes activated microorganisms as show in table (1) *Origanum vulgare* was activate *C.paraciticus* and *C. tropicus* also *Punica granatum* activate *C.paracciticus* , this may be because these extract have phytochemicals can activate cells nutrition, proliferation or activate some enzymatic pathway thus use herbal medicine must be in careful uses to avoid side effects.

REFERENCES

- 1- Albertson, G.D., Niimi, M., Cannon, R.D. and Jenkinson, H.F. (1996) Multiple eflux mechanisms are evolved in *Candida albicans* fluconazole resistance. *Antimicrob. Agents Chemother.* 40, 2835-2841.
- 2- Di Pasqua R, Hoskins N, Betts G, Mauriello G (2006). Changes in membrane fatty acids composition of microbial cells induced by addition of thymol, carvacrol, limonene, cinnamaldehyde, and eugenol in the growing media. *J. Agric. Food Chem.* 54(6): 2745-2749.
- 3- Turina AV, Nolan MV, Zygodlo JA, Perillo MA (2006). Natural terpenes: self-assembly and membrane partitioning. *Biophys. Chem.* 122(2): 101-113.
- 4- Oussalah M, Caillet S, Lacroix M (2006). Mechanism of action of Spanish oregano, Chinese cinnamon, and savory essential oils against cell membranes and walls of *Escherichia coli* O157:H7 and *Listeria monocytogenes*. *J. Food Prot.* 69(5): 1046-1055.
- 5- Arnal-Schnebel B, Hadji-Minaglou F, Peroteau JF, Ribeyre F, de Billerbeck VG (2004). Essential oils in infectious gynaecological disease: a statistical study of 658 cases. *Int. J. Aromather.* 14(4): 192-197.
- 6- Yoon HS, Moon SC, Kim ND, Park BS, Jeong MH, Yoo YH (2000). Genistein induces apoptosis of RPE-J cells by opening mitochondrial PTP. *Biochem. Biophys. Res. Commun.* 276(1): 151-156.
- 7- Al-Saadi. A.; Breesam. B. and Al-Turaihe. M. (2012) medical plants, AL radwan publisher 1st publish.
- 8- Al-Terehi, M. . Al Saadi , A. H. Al Ameri, Q. (2012). *In vivo* study of antimutagenic and antioxidant activity of *Glycyrrhiza glabra* root extract . *research in pharmacy* ,2,27-34
- 9- Sato, T. ;Onse, Y. ; Nagase, H. and Kito, H. (1990). Mechanism of antimutagenicity of aquatic plant extracts against (benzo (a)pyrene) in the *Salmonella* assay .*J. Mut. Res.* 241,283-290.
- 10- Jebor, M.; AL-saadi, A.; Hikmat, R.; AL-Terehi; Zaidan, H.K. and AL-Saadi. M.(2013).Characterization and antimicrobial activity of barley grain (*Hordeumvulgare*) extract. *Int.J.Curr.Microbiol.App.Sci* ,2(8): 41-

- 48.
- 11- Supreetha.S. , Sharadadevi Mannur , Sequeira Peter Simon , Jithesh Jain , Shreyas Tikare , Amit Mahuli (2011) Antifungal Activity of Ginger Extract on *Candida Albicans* :An In-vitro Study Journal of Dental Sciences and Research Vol. 2, Issue 2, Pages 1-5.
 - 12- Hayes BM, Bleackley, MR, Wiltshire JL, Anderson MA, Traven A, van der Weerden NL(2013) Identification and mechanism of action of the plant defensin NaD1 as a new member of the antifungal drug arsenal against *Candida albicans*. *Antimicrob Agents Chemother.* ;57(8):3667-75.
 - 13- Vale-Silva,L., Maria-Joaõ o Silva,1 Daniela Oliveira,1 Maria-José Gonc, alves,2 Carlos Cavaleiro,2 Lígia Salgueiro2 and Eugénia Pinto1 (2011) Correlation of the chemical composition of essential oils from *Origanum vulgare* subsp. *virens* with their in vitro activity against pathogenic yeasts and filamentous fungi *Journal of Medical Microbiology* (2012), 61, 252–260.
 - 14- Souzaa, E.L. , Stamforda, T.L.M, Limab, E.O. , Trajanob, V.N. (2007) Effectiveness of *Origanum vulgare* L. essential oil to inhibit the growth of food spoiling yeasts .*Food Control* Volume 18, Issue 5, May 2007, Pages 409–413.
 - 15- Khanuja, K. (2006) . Antimicrobial potential of *Glycyrrhiza glabra* Root. *CIMAP Com. J.* 39.
 - 16- Badieel, P, Nasirzadeh A, and Motaffaf, A. (2012) Comparison of *Salvia officinalis* L. essential oil and antifungal agents against *Candida* species *Journal of Pharmaceutical Technology& Drug Research*.
 - 17- Dulger,B. and Hacioglu N. (2008) Antifungal Activity of Endemic *Salvia tigrina* in Turkey *Tropical Journal of Pharmaceutical Research*, September 2008; 7 (3): 1051-1054.
 - 18- Ahmed, I. and A.Z. Beg, 2001. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J. Ethnopharmacol.*, 74: 113.
 - 19- Dahham, S.S; Ali,M.N. ; Tabassum; H. and Khan, M.(2010) Studies on Antibacterial and Antifungal Activity of Pomegranate (*Punicagranatum* L.). *American-Eurasian J. Agric. & Environ. Sci.*, 9 ,273-281.
 - 20- Yana H ,Velizar G ,Juergen W ,Leopold J ,Erich S, Tania G ,Andrei K,(2013) Chemical composition and antifungal activity of essential oil of *Salvia sclarea* L. from Bulgaria against clinical isolates of *Candida* species *J. BioSci. Biotech.* 2013, 2(1): 39-44.
 - 21- Al-Terehi, M. . Al Saadi , A. H. Al Ameri, Q. (2012). *In vivo* study of antimutagenic and antioxidant activity of *Glycyrrhiza glabra* root extract. *research in pharmacy* ,2,27-34

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

