

Prevalence of Major Gastrointestinal Helminthes Parasites of Small Ruminant in Enemay Woreda

Alemineh Shime* Samue Derso

Amhara National Regional State, East Gojjam Administration Zone, Gozamin Woreda Livestock and Fish Office, Animal Health Team

Abstract

A cross sectional study was conducted to determine the prevalence and risk factors associated with small ruminants GIT helminth parasites in Enemay Woreda, East Gojjamm, Northwest of Ethiopia from October, 2013 to April, 2014 based on coprological examination. A total of 384 small ruminants' fecal samples (248 sheep and 136 goats) were collected and examined using standard parasitological procedures of sedimentation and flotation techniques. The present study revealed that the overall prevalence of the major GIT helminthes parasite was 229 (59.63%). Out of 229 positive samples the species of parasite were found *Strongyle* (22.9%), *Fasciola* (14.1%), *Paramphistomum* (7.03%), *Monesia* (5.73%) and as mixed infection (9.9%). The study showed that 63.7% and 52.2 % of sheep and goats respectively were infected with one or more helminthes and higher prevalence was observed in sheep than goats and there was statically significant difference ($P<0.05$) between them. Female animals were found with higher prevalence of helminthes infection rate than male animals with a prevalence of 59.9% and 40.1% respectively and there was statically significant difference ($P<0.05$) between sex. Higher prevalence was observed in young animal than adult animal in this study and the prevalence was 67.9% and 53.6% respectively. There was statically significant difference ($P<0.05$) between age group. The study showed that higher prevalence of helminthic infection was observed in poor body condition animals as compared to medium and good body condition animals and their prevalence were 89.9%, 59% and 44% respectively. There was highly statically significant difference ($P<0.000$) between body condition of the animal. In conclusion the animal was affected by different helminth parasite infections which cause loss of production, reducing growth rate and death of small ruminants. So the animal owner should be deworming their small ruminants by different anthelmintics based on order of the Veterinarian to avoid drug resistance as recommendation.

Keywords: Enemay woreda, GIT helminthes, Prevalence, Risk factor, Small ruminants

INTRODUCTION

The livestock sector is a massive transformational state to meet increased demand of animal origin foods for increasing human population [1]. Ethiopia is believed to have the largest livestock population in Africa. This livestock sector has been contributing considerable portion to the economy of the country, and still promising to rally round the economic development of the country. It is eminent that livestock products and by-products in the form of meat, milk, honey, eggs, cheese, and butter supply etc. provide the needed animal proteins that contribute to the improvement of the nutritional status of the people. Livestock also plays an important role in providing export commodities, such as live animals, hides, and skins to earn foreign exchanges to the country. Ethiopia has an estimated of 53.4 million Cattle, 25.5 million sheep, 22.78 million goats, 2 million horses, 6.2 million donkeys, 0.38 million mules, about 1.1 million camels and 49.3 million poultries [2].

Sheep and goats are widely adapted to different climates and are found in all production system. They also have lower feed requirement as compared to cattle because of their small body size. This allows easy integration of small ruminants in to different farming system [3].

Parasitic helminthes or worms are important cause of disease in all species of animal. Although in many case they produce little serious damage to the host, these parasites are never beneficial in some case they can produce sever and even fatal disease [4].

In Ethiopia, 5-7 million sheep and goats die each year due to diseases including helminth infections. More significant, however, are losses resulting from inferior weight gains, condemnation of organs and carcasses and lower milk yields. The overall economic loss to the Ethiopian meat industry due to parasitic diseases is estimated at US\$ 400 million annually [5].

Sheep and goats harbor a variety of gastrointestinal tract (GIT) parasites, many of which are shared by both species. Among these parasites, helminthes are the most important GIT parasites that affect the growth as well as production of the animals. Gastrointestinal nematodes of *Trichostrongylidae* family are perhaps the most important parasites of small ruminants worldwide, causing significant morbidity and loss of production. Helminthic infections can be treated by anthelmintic chemotherapy, however, treatment is costly and drug resistance has evolved in all major parasite species [6].

In the varied agro-climatic zones of Ethiopia, small ruminants are important source of income for rural communities and are one of the nation's major sources of foreign currency from exports. In Ethiopia about 8 millions of small ruminants are slaughtered annually and providing more than 30% of domestic meat

consumption. The rich potential from the small ruminant sector is not efficiently exploited; however, due to several constraints, including malnutrition, inefficient management and diseases [7].

The sheep and goats provide cash income, meat and skin to the Enemay woreda society and to different hotels in Bichena town. The animal mostly affected by different disease due to suitability of the woreda to different disease epidemiology including helminthic infection and their productivity is low. But there is no enough information about the prevalence of major GIT helminth parasites of small ruminants in the woreda.

Therefore the objectives of this study in the study area were:

- To determine the prevalence of gastrointestinal parasite of small ruminants in the study area.
- To assess the major risk factors associated with prevalence of GIT parasites of small ruminants.

MATERIAL AND METHODS

Study area

The study was conducted from October 2013 to April, 2014 at Enemay Woreda which is found in East Gojjam administration zone, in Amhara region at 265km in Northwest of Addis Abeba. The mean annual temperature of the Woreda is 21°C and annual rain fall is 815-1440mm. It lies at 10°27'North 38°12'East latitude and longitude respectively and 2572 meter above sea level. Enemay Woreda has 108224 of livestock population of which 71432 small ruminants that managed under extensive management system. The animal used as cash income, draught power, and as food source in the form of meat, milk and egg [8].

Study animals

The study animals were small ruminants (sheep and goats) in Enemay Woreda which are managed under extensive management system. These animals are maintained in small households flocks of mixed age group and sex. All the sheep and goats that the sample collected was indigenous breed and the animal was classified as young (≤ 1 year) and adult (> 1 year) according to [9] and age was estimated based own owners knowledge and pattern of incisor eruption [10] and body condition can be classified as poor, medium and good according to [11] and body condition Scoring is based on feeling the level of muscling and fat deposition over and around the vertebrae in the loin region [12]

Sample size determination

The sample size required for this study was determined based on sample determination in random sampling with expected prevalence of major gastro intestinal helminthes parasite of small ruminant in the study area is 50% which no previous know prevalence and at 5% desire absolute precision and 95% confidence level according to [13]. Therefore, the sample size of 384 small ruminants (248 sheep and 136 goats) was obtained by using formula for sample size determination as given below as follow.

$$n = \frac{1.96^2 p \text{exp} (1 - p \text{exp})}{d^2}$$

Where:

n = required sample size,

pexp = expected prevalence = 50%

d = desired absolute precision = 5%

When this number substituted in the above formula the required sample size was 384.

Study design

The study design was cross-sectional which carried out to determine the prevalence of major GIT helminth parasites of small ruminants and to assess their prevalence based on coprological examination.

Sample collection and coprological examination

The sample was collected from 384 small ruminants (248 sheep and 136 goats) directly from the rectum which is placed on sample container bottle with 10% formalin as preservative. During sample collection, date, sex, species of animal, age, and body condition of the animal were properly recorded. After collecting the sample was examined by flotation and sedimentation technique at Enema Woreda Veterinary clinic with a standard parasitological procedure described by (14). Eggs of the different helminthes were identified on the basis of morphological appearance and size with the help of keys ([15]).

Data entry and analysis

All collected data were entered to Micro- Soft Excel sheet version 2007 and analyzed by SPSS version 20. Descriptive statistics was used to determine the prevalence of the parasites and Chi-square test was used to assess the association of the potential risk factors with the prevalence of the parasites. For statistical analysis a confidence level of 95% and P-values less than 5% ($p < 0.05$) was considered as significant.

RESULTS

Out of the total 384 (248 sheep and 136 goats) small ruminants examined over the study period, 229 (59.63%) were found to harbor one or more parasite species.

Out of the total of 248 (63.7%) of the sheep and 136 (52.2%) of the goats studied were found to harbor one or more parasite species. There was statically significant difference between the two species ($\chi^2=4.829$, and $P<0.05$ (Table 2).

The prevalence of major GIT helminthes parasite in relation to sex, 59.9% in female and 40.15% in male were observed. Higher prevalence was recorded in female (59.9%) than in male (40.1%) and there was statically significant between sex ($\chi^2=9.77$, and $P<0.05$ (Table 3).

The prevalence of major GIT helminthes parasite in different age group were 67.9% in young and 53.6% in adult sheep and goat and there was statically significant between age ($\chi^2=7.954$, and $P<0.05$ (Table 4).

Higher prevalence was observed in poor body condition (89.9%) as compared to medium (59.7%) and good (44%) body condition. There was also highly statically significant between body condition ($\chi^2=39.734$, and $P<0.000$ (Table 5).

The distribution of different classes of helminths parasites of small ruminant in the study area were nematodes (*Strongyle* type) followed by trematodes (*Fasciola* and *Paramphistomum*) and cestodes (*Monesia*) in both host species. The overall prevalence of the parasite based on specie of parasite was 22.92% *Strongyle*, 14.1% *Fasciola*, 7.03% *Paramphistomum*, 5.73% *Monesia* and 9.9% mixed infection (Table 6).

Table 2: prevalence of major GIT helminthes parasite based on species small ruminant

Species	No. exam	No.+ve	Prevalence (%)	χ^2	p-value
Sheep	248	158	63.7	4.829	0.028
Goat	136	71	52.2		
Total	384	229	59.63		

Table 3: prevalence of major GIT helminthes parasite based on sex of the animal

Sex	No. exam	No.+ve	Prevalence (%)	χ^2	p-value
Female	230	152	59.91	9.77	0.0002
Male	154	77	40.1		
Total	384	229	59.63		

Table 4: prevalence of major GIT helminthes parasite based on age

Age	No. exam	No.+ve	Prevalence (%)	χ^2	p-value
Young	162	110	67.9	7.954	0.005
Adult	222	119	53.6		
Total	384	229	59.63		

Table 5: prevalence of major GIT helminthes parasite based on body condition

Body condition	No. exam	No.+ve	Prevalence (%)	χ^2	p-value
Poor	69	62	89.9	39.734	0.000
Medium	181	108	59		
Good	134	59	44		
Total	384	229	59.63		

Table 6: Prevalence of major GIT helminthes parasite by study animal

Species	No. exam.	No.+ve	Strongyle (%)	Fasciola (%)	Paramphistomum (%)	Monesia (%)	Mixed (%)
Sheep	248	158(63.7)	61(24.6)	38(15.32)	20 (8.1)	17(6.9)	22(8.9)
Goat	136	71(52.2)	27(19.9)	16(11.8)	7(5.1)	5(3.7)	16(11.8)
Total	384	229	88(22.92)	54(14.1)	27(7.03)	22(5.73)	38(9.9)

DISCUSSIONS

The present study revealed that the overall prevalence of GIT helminth parasites was 59.63% in the small ruminants examined. This finding is comparable with the finding of [16] reported 61.4%, in Haremaya, South Eastern Ethiopia and lower than the results of other studies in sheep and goat carried out in different part of Ethiopia [17] 70.2% in Central Oremia, [18] 87.2% around Jima town, Western Ethiopia, [19] 87.3% in Gechi District, Southwest Ethiopia and elsewhere in the world (20) 96.0% in Tarai region of Uttarakhand, and [21] 69.7% in Ladakh, India. The current lower prevalence finding might be due to now a day the animal owner

manages their animal properly by regular deworming by different anthelmintics during different season of the year, proper feeding of their animal that helps the animal to protect themselves from different helminthes infection by developing rapid immune response to the parasite. Different parasite require different agro climate for multiplication and survival of the infective stage of the parasite and infect the animal and this area might be do not allow this things for the parasite.

The present study showed that 63.7% and 52.2 % of sheep and goats respectively are infected with one or more helminthes and higher prevalence was observed in sheep than goats which is agreed with other studies that reported higher prevalence in sheep than goats [22] 96.25% and 86% in Bokova, a rural area of Buea Sub Division, Cameroon, in sheep and goats respectively. This is higher prevalence in sheep might be due to the grazing habit of sheep when they graze closer to the ground might be consumed the infective stage of the parasite with the grass from the ground where as goats are mostly not grazing close to the ground rather they are brose the leaf of the tree which is above the ground that prevent themselves from exposure to the infective stage of the parasites.

Female animals were found with higher prevalence of helminthes infection rate than male animals and there was statically significant ($p < 0.05$) between them in the present study. The prevalence of GIT helminthes parasite in this study in female and male animal was 59.9% and 40.1% respectively. This finding agreed with other studies which are reported higher prevalence in female than male [23] 48.80% and 42.42% in North Gondar zone, Northwest Ethiopia in female and male animal respectively. The higher prevalence in female animals observed in the study due to male animals are slaughter early and more samples were collected from the female, and female animals immunity may be lowered than male animal during lactation and pregnancy and also male animals are kept indoor for the purpose of fattening where as female animals are not mange just like a male animal which are kept on communal grazing on the field.

Higher prevalence was observed in young animal than adult animal in this study and there was statically significant ($p < 0.05$) between age group. The prevalence of GIT helminthes parasite in this study young and adult animal was 67.9% and 53.6% respectively. This study is similar to other finding that reported higher prevalence in young animal than adult animal such as, [24] 79.6% and 62.4% in and around Asella, South Eastern Ethiopia.

This might be due to young animals are susceptible to different diseases including parasitic infection due to low development of immune response to the infection, lack of adaptation and resistance before they exposure to infection where as adult animals are resistant and adapted to infection due rapid response of immunity to the infection due to previous exposure of infection which remove the parasite before it attach to its predilection site.

The study showed that higher prevalence of helminthic infection was observed in poor body condition animals as compared to medium and good body condition animals and there was highly statically significant ($P < 0.000$) between body condition. The prevalence of helminthes parasite in these studs in relation to body conditions 89.9%, 59% and 44% in poor, medium and good body condition. This finding is similar to other studies which is [25] in and Around Mekelle Town, Northern Ethiopia, The higher prevalence in poor body conditions might be caused by due to malnutrition, other concurrent diseases or current parasitic infection that lead to lower the immune status of the animal to different diseases or infective stage of the parasites.

The major helminthe parasite that has been observed in this study were *Strongyle* type of species (Nematodes), *Fasciola* and *Paramphistomum* species (Trematode) and *Monesia* species (Cestode) parasites of small ruminant in this area. The overall prevalence of this parasite in this animal was 22.92 % *Strongyle*, 14.1% *Fasciola*, and 7.03% *Paramphistomum* 5.73% *Monesia* species of helminth parasite in small ruminants. This finding agreed with [26] in Ganderbal, Kashmir. The highest prevalence was seen in *Strongyle* type of parasite than other helminth parasites this might be due to the area is suitable to the survival of the infective stage of the parasite which means there was optimal moisture and temperature that helps to the egg of parasite to hatch and develop to the infective stage outside the definitive host. The development of larvae in the environment depends upon warm temperature and adequate moisture. In most tropical and sub-tropical countries, temperatures are permanently favourable for larval development in the environment.

CONCLUSION AND RECOMENDATIONS

Enemay Woreda has large number of small ruminant that are managed under extensive management system in mixed farming system that serve as source of food and cash income for rural society of the woreda. But the animal was affected by different helminth parasites such as *Strongyle* type, *Fasciola* species, *Paramphistomum* and *Monesia* specie of parasite and sometimes by mixed parasitic infection which causes loss of production, reducing growth rate and death of small ruminants due to lack of proper management like regular deworming, improper feeding, animals are keeping on communal grazing on the field and lack of adequate animal health and production extension workers that give to advise to the animal owner .

Based on the above conclusion the following recommendations are forwarded:

- ❖ The animal owner should be deworming their small ruminants by different anthelmintics based on order

of the Veterinarian to avoid drug resistance.

- ❖ The government should be creating awareness to the animal owners to avoid communal grazing and keep their animal indoor to improve the production and productivity of the animal.

REFERENCES

- 1.Karim, S.A., Tripath, M.K., Chaturved, O.H. and Shinde, A.K ., 2008. *Small ruminant production in India*. SSPH.
2. (CSA). 2011. Federal Democratic Republic of Ethiopia. Volume II: Report on livestock and livestock characteristic. Statistical bulletin. 505. Addis Ababa, Ethiopia.
- 3.Alemu, Y. and Merkel, R.C., 2008. Sheep and Goat Production Hand Book for Ethiopia
- 4.Jones, T.C, Hunt, R.D. and King, N.W., 1996. *Veterinary pathology*. 6th ed. Pp. 601.
- 5.MOARD. 2007. Ethiopia Sheep and Goat Productivity Improvement Program. Technical bulletin no.3. Control of internal parasites in sheep and goats.
- 6.Ijaz, M., Khan M. S., M., Avais, Ashraf, K., Ali, M. M. and Khan ,M. Z.U., 2009. Infection rate and chemotherapy of various helminthes in diarrhoeic sheep in and around Lahore. *The journal of animal & plant sciences*, 19 (1), p. 13-16.
- 7.Abebe, W. and Esayas, G., 2001. Survey of ovine and caprine gastro-intestinal helminthosis in eastern part of Ethiopia during the dry season of the year. *Revue Med.Vet.*, 152 (5), p. 379-384.
- 8.ERWARD. 2012. Enemay woreda agriculture and rural development office.
9. Fikru, R., Teshale, S., Reta D. and Yosef .K., 2006. Epidemiology of gastrointestinal parasites of ruminants in Western Oromia, Ethiopia . *Intern J Appl Res Vet Med*, 4(1)
- 10.MOARD., 2009. Ethiopia Sheep and Goat Productivity Improvement Program. Technical bulletin no. 23. Estimation of weight and age of sheep and goats.
- 11.Asmare, A., Assefa, K., and Tewdrose, F., 2012. Occurrence of small ruminant Ectoparasite in and around Bahir Dar Northwest of Ethiopia. *Advance in Biological Research*, 6 (5), p. 170-176.
- 12.Thompson, J. and Meyer, H., 1994. Oregon State University Extension Service offers educational programs, U.S. Department of Agriculture, and Oregon counties.
- 13.Thursfield, M., 2005. *Veterinary epidemiology*. 3rd ed. Black well. Pp .233.
- 14Hansen, J. and Perry, B., 1994. *The epidemiology, diagnosis and control of helminth parasites of ruminants: A hand book*. ILRAD. Nairobi, Kenya.
- 15.Urquhart, M.G, Armour, J., Duncan, L.J.; Dunn, M.A. and Janninss, W.F., 1996. *Veterinary parasitology*. 2nd ed. Black well science
- 16.Tesfaheywet Z., 2012. Helminthosis of sheep and goats in and around Haramaya, South eastern Ethiop. *Journal of Veterinary Medicine and Animal Health*, 4 (3), p. 48-55.
- 17.Berisa, K., Tigist, T., Teshal, S., Reta, D. and Bedru, H., 2011. Helminthes of sheep and goat in Central Oromia (Ethiopia) during dry season. *Journal of Animal and Veterinary advances*, 10, p. 1845-1849.
- 18.Nuraddis, I., Mulugeta, T., Mihreteab, B. and Sisay A., 2014. Prevalence of gastrointestinal parasites of small ruminants inand around Jimma town, Western Ethiopia. *Acta parasitologica globalis*, 5 (1), p. 26-32.
- 19.Bikila, E., Yeshitla, A., Worku,T., Teka, F. and Benti, D., 2013 Epidemiology of gastrointestinal parasites of small ruminants in Gechi District, Southwest Ethiopia. *Advances in Biological Research*, 7 (5), p. 169-174.
- 20.Pant, K., Rajput, M.K.S., Kumar, J., Sahu, S., Rajkumari, V. and Gangwar, P., 2009. Prevalence of helminthes in small ruminants in Tarai region of Uttarakhand. *veterinary world*, 2 (7), p.265-266.
- 21.Kuchai, J.A, Chishti, M.Z., Zaki, M.M, Ahmad, J., Rasool, M., Dar, S.A. and Tak, H., 2011. Epidemiology of helminth parasites in small ruminants of Ladakh, India. *Online Journal of Animal and Feed Research*, 1 (5), p. 239-242.
- 23.Shimelis, D., Asmare, A. and Wudu, T., 2011. Epidemiology of gastrointestinal helminthiasis of small ruminants in selected sites of North Gondar zone, Northwest Ethiopia. *Ethiopian Veterinary Journal*, 15 (2), p.57-68.
- 24..Diriba, L. and Birhanu, A., 2013. Prevalence of ovine gastrointestinal nematodes in and around Asella, South Eastern Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 5 (8), p. 222-228.
- 25.Welemehret N., Basaznew, B. and Mersha, C., 2012. Helminthe parasites in small ruminants: prevalence, species composition and associated risk factors in and around Mekelle town, Northern Ethiopia. *European Journal of Biological Sciences*, 4 (3), p.91-95.
- 26.Lone, B.A., Chishti, M.Z., Ahmad, F. and Tak, H., 2012. A Survey of gastrointestinal helminth parasites of slaughtered sheep and goats in Ganderbal, Kashmir. *Global Veterinaria*, 8 (4), p. 338-341. .

ACKNOWLEDGMENTS

First, I would like to praise almighty God for the patience he gave me to traverse this long journey and a

reason for the success of my educational career.

I have no words strong enough to express my heartfelt thanks to my advisor, Dr. Samuel Derso, for his intellectual guidance, provision of necessary materials, valuable and continuous advice and devotion his time to correct this thesis paper.

Final I would like to acknowledge all my family members and colleagues for their unforgettable financial and moral support throughout my work.