

Synthesis of New Complexes by Using Pyridine Acetohydrazone Derivatives

Bayan Al-Kilany¹ M.Moudar Al-Khuder²

1. Ph.D. student at Albaath university, faculty of science, department of inorganic chemistry

2. Prof. at Albaath university, faculty of science, department of inorganic chemistry

Abstract

A new ligand $L_1=[N-(1\text{-pyridine-2-yl})\text{Ethylidene AcetoHydrazide}]$; was synthesized by condensation of acetohydrazide with Acetyl pyridine. Then the complexes were synthesized by reacting the transition metals (Co^{2+} , Cr^{3+} , Fe^{2+}) with the ligand [with a metal to ligand ratio of (1:2) respectively] and this led to forming the following complexes : $[\text{Co}(L_1)_2] \text{Cl}_2$, $[\text{Cd}(L_1)\text{Cl}]$, $[\text{Fe}(L_2)\text{SO}_4]$. The ligands and complexes were characterized and studied on the basis of FT-IR, and U.V.-visible and the results were compatible with the proposed structures

Keywords: ligand, metal complexes, pyridine.

1. Introduction

Schiff bases are those compounds containing the active isomethyne group $\text{R}_1\text{R}_2\text{C}=\text{N}-$, which is due to multiple electronic and Stereotypical properties. The complexes formed by Schiff's bases with transitional metals ions were widely studied [1,2] and the spacial structure of a number of them was determined. The bases of Schiff with their properties are very useful compounds in the field of analysis and chemical separation. They are Clutch compounds used in copper calibration selectively [3]. They were also used as extraction factors for the determination of bi- copper as a photic[4] and chromatography to determine nickel in some natural food samples [5]. They were also used in the micro accurate determination of bi- cobalt photic[6], as well as in the process of extracting the ionic pair of binary metal cations [7]. The bases of Schiff derived from ketone condensation with the primary amines were named Ketimines [8]. The compounds derived from aldehydes condensation with amines were called Aldimines [9]. And in the case of condensation of hydrazidates with ketones or aldehydes in suitable Solvents, condensation products are called hydrazones [10].

2. Experimental

2.1 Materials


All the chemicals used were purchased from both Merck and Sigma Aldrich companies and used without further purification. FT-IR spectra was recorded using Jascow Japanese taype (A) Infrared Spectrophotometer Fourier Transform FT-IR-4100 (KBr). UV spectra was measured on Optizen spectrophotometer 200 -800nm.

3. Results and Discussion

3.1 Synthesis of a Schiff base ligand

The Schiff base ligand used in this study was prepared by refluxing an equimolar mixture of acetyl pyridine and the respective hydrazide in methanolic medium. A typical synthesis is described below.

30 ml methanol solution with acetyl pyridine (1.136 g, 10 mmol), aceto hydrazide (0.74g, 10mmole) with 50 ml methanol were added and the reaction mixture was stirred while refluxing for 3 h. Then the volume of the resultant was reduced to 30 ml. The solid resultant was filtered and thoroughly washed with ethanol (2×5 ml) followed by diethyl ether (2×5 ml). The solid was recrystallized by hot ethanol, Scheme 1.




Scheme 1. Synthesis route of the main reaction.

3.2 Syntheses of the complexes


All the complexes were prepared following a similar method of reacting a methanol solution with the metal salt [with a metal to ligand ratio of (1:2) respectively]. Preparation of one of the complexes is described below .

To a 50 ml methanol solution , the ligand (1 mmol), was added with stirring. When the dissolution is completed. Addition of MCl_2 anhydrous (0.5 mmol) when $[\text{M} = \text{Co}^{2+}, \text{Cr}^{3+}]$ and $[\text{FeSO}_4]$ anhydrous (0.5 mmol) in case of using (Fe^{2+}) to the colorless solution produced colored solution immediately. After 5 h of stirring the volume of the solution was reduced to 20 ml and filtered. The solid resultant was obtained and washed with

methanol (3 × 3 ml), followed by diethyl ether (2 × 5 ml), Scheme 2. Results of the compositional and spectroscopic analyses are shown in Tables 1-5.



Scheme 2. The main Synthesis of complexes.



Scheme 3. Reaction between the ligand and sulfate iron (II).

Table 1: Characterization of the Schiff base Ligands and metal(II) Complexes.

Compounds	Formulas	Color	m.p.°C	Yield (%)
L	C ₉ H ₁₁ N ₂ O ₁	White	160	50
[Co(L) ₂]Cl ₂	C ₁₈ H ₂₂ N ₄ O ₂ CoCl ₂	green	250-dec	40
[Cr (L) ₂]Cl	C ₁₈ H ₂₂ N ₄ O ₂ CrCl	green	210-dec	38
[Fe(L) ₂]SO ₄	C ₁₈ H ₂₂ N ₄ O ₆ FeS	black	280-dec	60

3.3 FT-IR spectra of the (L) ligand and its complexes:

The infrared spectra for the present compounds taken in the range 400-4000 cm⁻¹ helps to indicate regions of absorption vibrations. The main stretching modes are for ν(C=N), ν(C=C) and ν(C=O). The IR data of the spectra of Schiff base ligand (L) and its complexes are presented in Table 3. The IR spectra of the complexes were compared with those of the free ligands in order to determine the coordination sites that may be involved in chelation.

Spectrum of the (L) ligand shows a sharp band at (1677cm⁻¹) due to ν(C=N) azomethine group which has shifted to lower frequency about (70cm⁻¹) in the complexes indicating its participation in chelation through azomethine nitrogen. The lowering of band is due to the reduction of electron density in the azomethine link. Pyridine ν(C-N) shows absorption band at (1298cm⁻¹) in the Schiff base spectrum. This band makes shift by (1271cm⁻¹) in the complex. Azomethine group in the Schiff base (L) changed after complication .with Co(II) from (1677cm⁻¹) to (1638cm⁻¹) this indicate that involvement of Azomethine group in complications.

Table 2. Characteristic infrared absorption frequencies (cm⁻¹) of the ligand and complexes.

compounds	ν(NH)	ν(C=N)	ν(C=N) (C=N) _{py}	ν(C=O)	ν(C-H)SP ²
L	3192 _t	1617 _{st}	1577 _m	1677 _m	2924 _w
[Co(L) ₂]Cl ₂	3187 _{st}	1518 _{st}	1463 _m	1638 _m	2922 _w
[Cr (L) ₂]Cl	3213 _{st}	1602 _{st}	1557 _m	1673 _m	2925 _w
[Fe(L) ₂]SO ₄	3199 _{st}	1638 _{st}	1601 _m	1654 _m	2965 _w




Figure 1:FT-IR spectrum of ligand (L)




Figure 2: FT-IR spectrum of complex $[\text{Co}(\text{L})_2]\text{Cl}_2$




Figure 3: FT-IR spectrum of complex $[\text{Cr}(\text{L})_2]\text{Cl}$




Figure 4: FT-IR spectrum of complex $[\text{Fe}(\text{L})_2]\text{SO}_4$

3.4 Electronic spectral data:

The data of the electronic spectra of the ligand and its complexes are given in Table 3. The spectrum of Schiff base (L) presented two bands in the UV interval at 210nm and 286nm, assigned to $(\pi \rightarrow \pi^*)$ and $(n \rightarrow \pi^*)$ transitions respectively.

The electronic spectra of the complexes in methanol solution has two bands at (270, 370)nm. These bands may be assigned to the charge transitions (LMTC) of the form

$(n \rightarrow \pi^*)$ for azomethine group (C=N) and the pyridine ring was assigned to metal ion $\text{L} \rightarrow \text{M}$ [15].

The position of these bands suggests an octahedral environment to Co(II), Cr(II) and Fe(II).




Figure 5: UV absorption spectrum of ligand (L)




Figure 6: UV absorption spectrum of [Co(L)₂]Cl₂




Figure 7: UV absorption spectrum of [Cr(L)₂]Cl




Figure 8: UV absorption spectrum of [Fe(L)₂SO₄]

Table3: Magnetic moments, electronic bands and ligand filed parameters of L and its metal complexes [11].

compound	$\pi \rightarrow \pi^*$ (nm)	$n \rightarrow \pi^*$ (nm)	$L \rightarrow M$ (nm)	$d-d$ (nm)
L	210	286	---	---
[Co(L) ₂ Cl ₂]	211	270	370	---
[Cr(L) ₂ Cl]	215	230	290	---
[Fe(L) ₂ SO ₄]	210	220	280	---




Figure 9: Suggested structures for the L complex with Cr^{III}




Figure 10: Suggested structures for the L complex with Co^{II}




Figure 11: Suggested structures for the L complex with Fe^{II}

4. Conclusion

A new ligand $L_1 = [N-(1\text{-pyridine-2-yl}) \text{ Ethylidene AcitoHydrazide}]$; was synthesized by condensation of acetohydrazide with Acetyl pyridine. Then the complexes were synthesized by reacting the transition metals (Co^{2+} , Cr^{3+} , Fe^{2+}) with the ligand [with a metal to ligand ratio of (1:2) respectively] and this led to forming the following complexes: $[\text{Co}(L_1)_2] \text{Cl}_2$, $[\text{Cd}(L_1)\text{Cl}]$, $[\text{Fe}(L_2)\text{SO}_4]$. The ligands and complexes were characterized and studied on the basis of FT-IR, and U.V.-visible and the results were compatible with the proposed structures

5. Acknowledgment

The authors would like to express their thanks and appreciation to Albaath for supporting this research.

6. References

- [1]. Kaliyappan, T. Kannan, P. (2000) . " Co-ordination polymers Review Article, Prog. Polym ". Sci., V.25, pp.343-370.
- [2]. Yamada, S. (1999). "Advancement in stereochemical aspects of Schiff base metal complexes ", Coordin. Chem. Rev , V. (190-192), pp.537-555.
- [3]. Kormali, E. Kylic, E. (2002). " N, N'-disalicylidene-1,3-diaminopropane as a selective chelating titrant for copper(II) " Talanta, V. 58, pp.793-802.
- [4]. Cimerman, Z. Galic, N. Bosner, B. (1997) ." The Schiff bases of salicylaldehyde and aminopyridines as highly sensitive analytical reagents ", Anal. Chem. Acta, V. 343, pp.145-153.
- [5]. Fakhari, A. R. Khorrani, A. R., Naeimi, H. (2005) . "Synthesis and analytical application of a novel tetradentate N2O2 Schiff base as a chromogenic reagent for determination of nickel in some natural food samples ", Talanta, V. 66, pp.813-817.
- [6]. Khedr, A. M. Gaber, M. Issa, R. M. Erten, H. (2005). " Synthesis and spectral studies of (TA) and its Schiff bases with (TAAP) and (TAAH) , Dyes and Pigments ", V. 67, pp.117-126.
- [7]. Oshima, S. Hirayama, N. Kubono, K. Honjo, T. (2003) ." Ion-pair extraction behavior of divalent metal cations using neutral di-Schiff base ligands derived from 1,2-cyclohexanediamine and o-phenylenediamine ", Talanta, V.59, pp.867-874.
- [8]. W. R. Paryzek, M.T. Kaczmarek, V. Patroniak and I. P. Markiewicz, (2003) ,"Inorganic Chemistry Communications" 6, 26-29.
- [9]. G. Geindy, M. Omar and A. Haivdy, (2006) .Turk "J. Chem." 30, 361-382.
- [10]. A.K. Sen-Gupta and K. Gajela; "J. Ind. Chem." Soc., LVIII, 690 1981
- [11]. A Patra, A. K.; Mukherjee, R. Inorg. Chem. 1999, 38, 1388. (b)Ray, M. S.; Bhattacharya, R.; Chaudhuri, S.; Righi, L.; Bocelli, G.; Mukhopadhyay, G.; Ghosh, A. Polyhedron 2003, 22, 617. (c) Dey, M.; Rao, C. P.; Saarenketo, P.K.; Rissanen (2003) , K. Inorg. Chem. Commun. 2002, 5, 924. (d) Yang, C. T.; Vittal, J. J. Inorg. Chim. Acta, 344, 65.