

# MoO<sub>3</sub>-SiO<sub>2</sub> as an efficient and reusable heterogeneous catalyst for the synthesis of 5-substituted 1*H*-tetrazoles

S. Mohammad Sajadi<sup>1\*</sup>

1. Department of Petroleum Geoscience, Faculty of Sciences, University of Soran, PO box 624, Soran, Kurdistan Regional Government, Iraq, Tel:+9647503714550

\* E-mail of the corresponding author: [smohammad.sajadi@gmail.com](mailto:smohammad.sajadi@gmail.com), [mohammad.s@soranu.com](mailto:mohammad.s@soranu.com)

## Abstract

An efficient method for the preparation of 5-substituted 1*H*-tetrazole derivatives is reported using MoO<sub>3</sub>-SiO<sub>2</sub> as an efficient heterogeneous catalyst. This method has the advantages of high yields, simple methodology and easy work-up. This catalyst could be recycled very easily, which makes this methodology environmentally benign.

**Keywords:** 5-substituted 1*H*-tetrazole, [2+3] cycloaddition, MoO<sub>3</sub>-SiO<sub>2</sub>, solid acid

## 1. Introduction

Despite the scarcity of tetrazoles in natural systems, the chemistry of this heterocycles has gained increasing attention since the early quickly 1980 (Bulter 1996). Tetrazoles have a wide range of applications in pharmaceuticals as lipophilic spacers and carboxylic acid surrogates, in materials as specialty explosives and information recording systems, in coordination chemistry as ligands and also as precursors to a variety of nitrogen-containing compounds (Bulter 1996; Herr 2002; Holland 1967; Figdor 1967; Rhonnstad 2002; Klapötke 2009; John 1989; Modarresi-Alam 2007). However, the utility of these compounds is limited due to their insufficient synthetic availability.

Tetrazoles are commonly formed from nitriles or cyanamides and an azide source via a 1,3-dipolar cycloaddition (Kadaba 1973; Curran 1999; Huff 1993; Modarresi-Alam 2009; Nasrollahzadeh 2009; Habibi 2010) Unfortunately, each of those protocols suffers from some disadvantages: the use of both toxic metals and expensive reagents, drastic reaction conditions, water sensitivity, and the presence of dangerous hydrazoic acid. The use of hydrazoic acid presents considerable experimental difficulties because of its toxicity and tendency to explode (Duncia 1991; Carini 1991; Wittenberger 1993). Due to safety considerations, we required a method that did not use of hydrazoic acid or an azide source that produced hydrazoic acid in situ because of the associated hazards. Therefore, it is desirable to develop a more efficient and convenient method for the synthesis of 5-substituted tetrazoles.

Recently, Sharpless and co-workers reported an innovative and safe procedure for the preparation of 5-substituted 1*H*-tetrazoles from the corresponding nitriles and NaN<sub>3</sub> in the presence of a stoichiometric amount or 50 mol % of Zn(II) salts (Demko 2001). As the catalytic load on these homogeneous catalysts is high, the separation of the products and recycling of catalyst becomes difficult, shifting our preference in favor of high efficient heterogeneous catalysts. Since then there have been reports of the use of various heterogeneous catalysts (Matthews 2000; Kantam 2006; Amantini 2004; Schmidt 2007; Kantam 2005; Kantam 2006), but most of these catalysts contain zinc or aluminium. The heterogeneous catalysts effective in the synthesis of tetrazoles were believed to be limited to the compounds containing zinc or aluminium. The development of a catalytic synthetic method for tetrazoles still remains an active research area.

In recent years heterogeneous catalysts have gained significant importance in organic syntheses because of economic and environmental considerations. These catalysts are generally less expensive, eco-friendly, high reactive, easy to handle and recoverable (Habibi 2010; Habibi 2011; Nasrollahzadeh 2009; Kantam 2006; Mohammadi 2010; Modarresi-Alam 2007; Modarresi-Alam 2008). Among various silica-based heterogeneous catalysts, MoO<sub>3</sub>-SiO<sub>2</sub> has advantages of low cost, ease of preparation and can be recycled

(Dongare 2004; Maurya 2003; Umbarkar 2006). Molybdenum oxide supported on silica ( $\text{MoO}_3\text{-SiO}_2$ ) has proved to be an efficient catalyst in promoting various organic reactions such as Beckmann rearrangement and nitration of aromatics. However, the utility of this catalyst for the synthesis of tetrazoles has not been explored before. We herein report a new protocol for preparation of 5-substituted 1*H*-tetrazoles derivatives from nitriles using  $\text{MoO}_3\text{-SiO}_2$  as an efficient heterogeneous catalyst (Scheme 1).

## 2. Experimental

### 2.1. General

All reagents were purchased from Merck and Aldrich and used without further purification.  $^{13}\text{C}$  NMR and  $^1\text{H}$  NMR spectra were recorded on Bruker, 100 and 250 MHz using TMS as an internal standard. Chemical shifts are reported in ppm, and coupling constants are reported in Hz. IR spectra were recorded on a Shimadzu 470 spectrophotometer. TLC was performed on Merck-precoated silica gel 60-F254 plates. Nano  $\text{TiO}_2$  was prepared according to the literature (Hua 2007). Hua and co-workers studied the characterization of the Nano  $\text{TiO}_2$  using powder XRD and TEM (Hua 2007).

### 2.2. General procedure for preparation of 5-substituted 1*H*-tetrazoles

20%  $\text{MoO}_3\text{-SiO}_2$  was added to a mixture of nitrile (1.5 mmol), sodium azide (2.3 mmol) and distilled dimethylformamide (5 mL) and stirred at 120 °C for the appropriate time (Table 1). After completion of the reaction (as monitored by TLC), the catalyst was filtered and the filtrate was treated with ethyl acetate (35 mL) and 4*N* HCl (20 mL) and stirred vigorously. The resultant organic layer was separated and the aqueous layer was again extracted with ethyl acetate (25 mL). The combined organic layers were washed with water and concentrated to give the crude product. Column chromatography using silica gel gave pure product in high yield. All the products are known compounds and were characterized by comparing IR and  $^1\text{H}$  NMR spectral data as well as melting points with those reported in the literature. All yields refer to isolated products.

#### 5-Phenyl-1*H*-tetrazole (**2a**):

$^1\text{H}$  NMR:  $\delta$  7.58-7.64 (m, 3H); 8.01-8.06 (m, 2H).

#### 5-(4-Methylphenyl)-1*H*-tetrazole (**2c**):

$^1\text{H}$  NMR:  $\delta$  2.47 (s, 3H); 7.49 (d,  $J = 8.0$  Hz, 2H); 8.01 (d,  $J = 8.0$  Hz, 2H).

#### 5-(4-Nitrophenyl)-1*H*-tetrazole (**2d**):

$^1\text{H}$  NMR:  $\delta$  8.30 (d, 2H); 8.45 (d, 2H).

#### 5-(Naphthalen-2-yl)-1*H*-tetrazole (**2f**):

$^1\text{H}$  NMR:  $\delta$  7.59-7.68 (m, 2 H); 7.98-8.18 (m, 4H); 8.65 (s, 1H).

## 3. Result and Discussion

The general synthetic method is depicted in Scheme 1. 5-Substituted 1*H*-tetrazoles were obtained from the reaction of nitrile with sodium azide in the presence of 20 %  $\text{MoO}_3\text{-SiO}_2$  as an efficient heterogeneous catalyst as a solid acid catalyst at 120 °C for appropriate time in high yields.

To evaluate the scope and limitations of the methodology, reactions were carried out with various substituted benzonitriles including both electron-donating and electron-withdrawing substituents at *ortho*, *meta*, and *para* positions of the aromatic ring (Table 1). The results showed that there is no adverse effect of substituents, either electron-donating or electron-withdrawing, on the aromatic ring of benzonitriles on the product yield. All the products were characterized by comparing melting points with those of the reported compounds. The nature of the substituent on the benzonitrile did not affect the reaction time.

Interestingly 1,4-dicyanobenzene (Table 1, entry 8) afforded the mono-addition product, whereas in the reaction between sodium azide and 1,4-dicyanobenzene in the presence of Zn(II) salts the double-addition product was reported (Demko 2001). Reaction of the heteroaromatic nitrile, 3-pyridinecarbonitrile was complete at 120 °C after 6 h and gave the corresponding tetrazole in an excellent yield (Table 1, entry 9).

Because of environmental concerns, there is increasing need and interest in developing processes that minimize production of toxic hydrazoic acid. Here, because of direct use of sodium azide rather than hydrazoic acid in the synthesis of tetrazoles, a significant improvement in the synthesis is represented. Replacement of hydrazoic acid by inorganic azides, which are poorly soluble in organic solvents with low dielectric constants, causes an increase in reaction time and decrease in tetrazole yield (Herr 2002). Water was not a suitable solvent for this reaction. Not many organic solvents are stable at the high temperatures necessary for cycloaddition reactions (sometimes as high as 130 °C), and for this reason DMF is most commonly used for this purpose (Herr 2002; Kantam 2005; Kantam 2006).

Many solid acid catalysts have been used for the same transformation. Dongare and co-workers (Dongare 2004; Maurya 2003; Umbarkar 2006), have shown that 20% MoO<sub>3</sub>-SiO<sub>2</sub> is very strong solid acid catalyst with acidity equivalent to 0.94 mmol NH<sub>3</sub> desorbed per gram of catalyst with 180m<sup>2</sup>/g surface area and 71.64 Å average pore diameter. This catalyst contains molybdenum oxide nanoparticles of 1-2 nm dispersed on mesoporous silica. Fourier transform-infrared (FT-IR) spectra of adsorbed pyridine showed that the catalyst contains both Lewis as well as Brønsted acidity. Detailed Raman spectroscopic studies have shown the in situ formation of silicomolybdic acid on the catalyst surface in the presence of moisture. This in situ formed silicomolybdic acid imparts strong acidity to the catalyst which is responsible for very high catalytic activity. One of the main advantages of MoO<sub>3</sub>-SiO<sub>2</sub> as catalyst was clean reactions without any side product. The MoO<sub>3</sub>-SiO<sub>2</sub> catalyst could be recycled efficiently for three cycles without any appreciable loss in the yield. This reusability demonstrates the high stability and turnover of MoO<sub>3</sub>-SiO<sub>2</sub> under operating condition. The reusability of the catalysts is one of the most important benefits and makes them useful for commercial applications.

The simplicity, together with the use of inexpensive, non-toxic and environmentally benign catalyst under thermal conditions is remarkable features of the procedure. The products were characterized by IR, <sup>1</sup>H NMR and <sup>13</sup>C NMR spectroscopy and from melting points. The disappearance of one strong and sharp absorption band (CN stretching band), and the appearance of an NH stretching band in the IR spectra, were evidence for the formation of 5-substituted 1*H*-tetrazoles. <sup>13</sup>C NMR spectra display signals for tetrazole ring carbons of arylaminotetrazoles in the range of 154-157 ppm (depending on the nature of the substituents in the amino functionality) (Goljer 1983; Habibi 2010; Habibi 2011; Nasrollahzadeh 2009; Nasrollahzadeh 2010).

#### 4. Conclusion

In conclusion, we have developed a novel and highly efficient method for the synthesis of 5-substituted 1*H*-tetrazoles by treatment of nitriles with sodium azide in the presence of MoO<sub>3</sub>-SiO<sub>2</sub> as a reusable and efficient heterogeneous catalyst. The significant advantages of this methodology are high yields, elimination of dangerous and harmful hydrazoic acid and a simple work-up procedure. The catalyst can be recovered by simple filtration and reused without loss of activity. It also has excellent activity on an industrial scale and in most cases can be recovered from reaction mixtures and reused.

#### References

- Amantini, D., Beleggia, R., Fringuelli, F., Pizzo, F., Vaccoro, L. (2004), "TBAF-catalyzed synthesis of 5-substituted 1*H*-tetrazoles under solvent less conditions", *Journal of Organic Chemistry* **69**, 2896-2898.
- Bulter, R. N. (1996), "In Comprehensive Heterocyclic Chemistry II", Katritzky, A. R., Rens, C.W., Scriven, E. F. V. Eds., Pergamon: New York, 1996; Vol. 4, p 621.
- Carini, D. J., Duncia, J. V., Aldrich, P. E., Chui, A. T., Johnson, A. L., Pierce, M. E., Price, W. A., Santella, J. B. III., Wells, G. J., Wexler, R. R., Wong, P. C., Yoo, S.-E., Timmermans, P. B. M. W. M. (1991), "Nonpeptide angiotensin II receptor antagonists: the discovery of a series of

N-(biphenylmethyl)imidazoles as potent, orally active antihypertensives”, *Journal of Medicinal Chemistry* **34**, 2525-2547.

Curran, D. P., Hadida, S., Kim, S. Y. (1999), “Tris(2-perfluorohexylethyl)tin azide: A new reagent for preparation of 5-substituted tetrazoles from nitriles with purification by fluoruous/organic liquid-liquid extraction”, *Tetrahedron* **55**, 8997-9006.

Demko, P. Z. Sharpless, K. B. (2001), “Preparation of 5-substituted 1H-tetrazoles from nitriles in water”, *Journal of Medicinal Chemistry* **66**, 7945-7950.

Dongare, M. K., Bhagat, V. V., Ramana, C. V., Gurjar, M. K. (2004), “Silica-supported MoO<sub>3</sub>: A mild heterogeneous catalyst for the Beckmann rearrangement and its application to some sugar-derived ketoximes”, *Tetrahedron Letters* **45**, 4759-4762.

Duncia, J. V., Pierce, M. E., Santella, J. B. III. (1991), “Three synthetic routes to a sterically hindered tetrazole. A new one-step mild conversion of an amide into a tetrazole”, *Journal of Organic Chemistry* **56**, 2395-2400.

Figdor, S. K., Schach von Wittenau, M. (1967), “Metabolism of 5-(3-Pyridyl)tetrazole”, *Journal of Medicinal Chemistry* **10**, 1158-1159.

Goljer, I., Svetlik, J., Hrusovsky, I. (1983), “<sup>13</sup>C-NMR of substituted tetrazoles”, *Monatshefte fur Chemie* **114**, 65-70.

Habibi, D., Nasrollahzadeh, M., Faraji, A. R., Bayat, Y. (2010), “Efficient synthesis of arylaminotetrazoles in water”, *Tetrahedron* **66**, 3866-3870.

Habibi, D., Nasrollahzadeh, M. (2010), “Silica-Supported Ferric Chloride (FeCl<sub>3</sub>-SiO<sub>2</sub>): An Efficient and Recyclable Heterogeneous Catalyst for the Preparation of Arylaminotetrazoles”, *Synthetic Communication* **40**, 3159-3167.

Habibi, D., Nasrollahzadeh, M., Bayat, Y. (2011), “AlCl<sub>3</sub> as an effective Lewis acid for the synthesis of arylaminotetrazoles”, *Synthetic Communications* **41**, 2135-2145.

Habibi, D., Nasrollahzadeh, M., Kamali, T. A. (2011), “Green synthesis of the 1-substituted 1H-1,2,3,4-tetrazoles by application of the Natrolite zeolite as a new and reusable heterogeneous catalyst”, *Green Chemistry* **13**, 3499-3504.

Habibi, D., Nasrollahzadeh, M. (2011), “Synthesis of arylaminotetrazoles by ZnCl<sub>2</sub>/AlCl<sub>3</sub>/silica as an efficient heterogeneous catalyst”, *Monatshefte fur Chemie* DOI: 10.1007/s00706-011-0670-8.

Herr, R. (2002), “5-Substituted-1H-tetrazoles as carboxylic acid isosteres: medicinal chemistry and synthetic methods”, *Journal of Bioorganic Medicinal Chemistry* **10**, 3379-3393.

Holland, G. F., Pereira, J. N. (1967), “Heterocyclic tetrazoles, a new class of lipolysis inhibitors”, *Journal of Medicinal Chemistry* **10**, 149-154.

Huff, B. E., Staszak, M. A. (1993), “A new method for the preparation of tetrazoles from nitriles using trimethylsilylazide/trimethylaluminum”, *Tetrahedron Letters* **34**, 8011-8014.

John, E. O., Kirchmeier, R. L., Shreeve, J. M. (1989), “Reactions of 5-(Perfluoro-alkyl)tetrazolates with Cyanogen, Nitrosyl, and Cyanuric Chlorides”, *Inorganic Chemistry* **28**, 4629.


Kadaba, P. K. (1973), “Role of protic and dipolar aprotic solvents in heterocyclic syntheses via 1,3-dipolar cycloaddition reactions”, *Synthesis* 71-84.

Kantam, M. L., Balasubrahmanyam, V., Kumar, K. B. S. (2006), “Zinc Hydroxyapatite-Catalyzed Efficient Synthesis of 5-Substituted 1H-Tetrazoles”, *Synthetic Communications* **36**, 1809-1814.

Kantam, M. L., Shiva Kumar, K. B., Phani Raja, K. (2006), “An efficient synthesis of 5-substituted 1H-tetrazoles using Zn/Al hydrotalcite catalyst”, *Journal of Molecular Catalysis A: Chemical* **247**, 186-188.





















Kantam, M. L., Shiva Kumar, K. B., Sridhar, C. (2005), “Nanocrystalline ZnO as an efficient heterogeneous catalyst for the synthesis of 5-substituted 1H-tetrazoles”, *Advanced Synthesis and Catalysis* **347**, 1212-1214.

- Klapötke, T. M., Stierstorfer, J., Weber, B. (2009), "New energetic material: synthesis and characterization of copper 5-nitriminetetrazolates", *Inorganic Chemical Acta* **362**, 2311-2320.
- Matthews, D. P., Green, J. E., Shuker, A. J. (2000), "Parallel synthesis of alkyl tetrazole derivatives using solid support chemistry", *Journal of Combinatorial Chemistry* **2**, 19-23.
- Maurya, S. K., Gurjar, M. K., Malshe, K. M., Patil, P. T., Dongare, M. K., Kemnitz, E. (2003), "Solid acid catalyst for fluorotoluene nitration using nitric acid", *Green Chemistry* **5**, 720-723.
- Mihina, J. S., Herbst, R. M. (1950), "The reaction of nitriles with hydrazoic acid: synthesis of monosubstituted tetrazoles", *Journal of Organic Chemistry* **15**, 1082-1092.
- Modarresi-Alam, A. R., Khamooshi, F., Rostamizadeh, M., Keykha, H., Nasrollahzadeh, M., Bijanzadeh, H. R., Kleinpeter, E. (2007), "Dynamic <sup>1</sup>H NMR spectroscopic study of the restricted S-N rotation in aryl-N-(arylsulfonyl)-N-(triphenylphosphoranylidene)imidocarbamates", *Journal of Molecular Structure* **841**, 61-66.
- Modarresi-Alam, A. R., Nasrollahzadeh, M. (2009), "Synthesis of 5-arylamino-1H(2H)-tetrazoles and 5-amino-1-aryl-1H-tetrazoles from secondary arylcyanamides in glacial acetic acid: a simple and efficient method" *Turkish Journal of Chemistry* **33**, 1-14.
- Modarresi-Alam, A. R., Khamooshi, F., Nasrollahzadeh, M., Amirazizi, H. A. (2007), "Silica supported perchloric acid (HClO<sub>4</sub>-SiO<sub>2</sub>): an efficient reagent for the preparation of primary carbamates under solvent-free conditions", *Tetrahedron* **63**, 8723-8726.
- Modarresi-Alam, A. R., Nasrollahzadeh, M., Khamooshi, F. (2007), "Solvent-free preparation of primary carbamates using silica sulfuric acid as an efficient reagent", *Arkivoc* (**xvi**), 234-245.
- Modarresi-Alam, A. R., Nasrollahzadeh, M., Khamooshi, F. (2008), "Al(HSO<sub>4</sub>)<sub>3</sub> Mediated for the preparation of primary carbamates under solvent-Free conditions", *Scientia Iranica* **15**, 452-455.
- Mohammadi, B., Hosseini Jamkarani, S. M., Kamali, T. A., Nasrollahzadeh, M., Mohajeri, A. (2010), "Sulfonic acid-functionalized silica: a remarkably efficient heterogeneous reusable catalyst for the one-pot synthesis of 1,4-dihydropyridines", *Turkish Journal of Chemistry* **34**, 613-619.
- Nasrollahzadeh, M., Bayat, Y., Habibi, D., Moshae, S. (2009), "FeCl<sub>3</sub>-SiO<sub>2</sub> as a reusable heterogeneous catalyst for the synthesis of 5-substituted 1H-tetrazoles via [2+3] cycloaddition of nitriles and sodium azide", *Tetrahedron Letters* **50**, 4435-4438.
- Nasrollahzadeh, M., Habibi, D., Shahkarami, Z., Bayat, Y. (2009), "A general synthetic method for the formation of arylaminotetrazoles using natural natrolite zeolite as a new and reusable heterogeneous catalyst", *Tetrahedron* **65**, 10715-10719.
- Rhonnstad, P., Wensbo, D. (2002), "On the relative strength of the 1H-tetrazol-5-yl- and the 2-(triphenylmethyl)-2H-tetrazol-5-yl-group in directed ortho-lithiation", *Tetrahedron Letters* **43**, 3137-3139.
- Schmidt, B., Meid, D., Kieser, D. (2007), "Safe and fast tetrazole formation in ionic liquids", *Tetrahedron* **63**, 492-496.
- Umbarkar, S. B., Biradar, A. V., Mathew, S. M., Shelke, S. B., Malshe, K. M., Patil, P. T., Dagde, S. P., Niphadkar, S. P., Dongare, M. K. (2006), "Vapor phase nitration of benzene using mesoporous MoO<sub>3</sub>/SiO<sub>2</sub> solid acid catalyst", *Green Chemistry* **8**, 488-493.
- Wittenberger, S. J., Donner, B. G. (1993), "Dialkyltin oxide mediated addition of trimethylsilyl azide to nitriles. A novel preparation of 5-substituted tetrazols", *Journal of Organic Chemistry* **58**, 4139-4144.



Scheme 1. Conversion of nitriles to the corresponding 5-substituted 1H-tetrazoles using MoO<sub>3</sub>-SiO<sub>2</sub>

Table 1. Synthesis of various 5-substituted 1*H*-tetrazoles in the presence of MoO<sub>3</sub>-SiO<sub>2</sub> by reaction of sodium azide and nitriles at 120 °C

Entry	Substrate	Product	Time [h]	Yield [%] <sup>a</sup>	Ref.
1			14	86	(Kantam 2005)
2			14	83	(Kantam 2005)
3			14	84	(Nasrollahzadeh 2009)
4			14	84	(Amantini 2004)
5			14	81	(Kantam 2005)
6			14	83	(Demko 2001)
7			14	83	(Nasrollahzadeh 2009)
8			22	84	(Nasrollahzadeh 2009)
9			6	84	(Kantam 2005)
10			24	77	(Kantam 2005)
11			24	78	(Kantam 2005)
12			30	65	(Mihina 1950)

<sup>a</sup>Yield refer to the pure isolated product.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### **IISTE Knowledge Sharing Partners**

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

