

EFFICACY OF XYLOPIA AETHIOPICA (DUNAL) A. AND AFRAMOMUM MELEGUETA (ROSC) K. SCHUM. FRUIT ON SITOPHULIUS ZEA

¹*AJURUCHI, V.C. AND DURU, C. M.²

¹Department of Biological Sciences, Federal University of Technology, P. M. B. 1526, Owerri Imo State, Nigeria.

²Department of Biological Science, Federal University of Technology, P.M.B. 1526, Owerri Imo State, Nigeria.

ABSTRACT

The power of combined powder of dried fruits of *Xylopiya aethiopica*(Dunal) A. and *Aframomum melegueta* on the control of maize weevil *Sitophulus zea* were evaluated. The aim was to determine, their antifeedant potentials against the insect pests of stored maize. Oven dried fruits of X. *aethiopica* and A. *melegueta* was pulverized and mixed in equal proportion. Concentrations of 1,2,3,4,5,6,7,8,9,and10g were mixed with 100g of healthy grain, and incubated for 14 days with 20 weevils in each. Each treatment plus the control (0.0g) were replicated five times and laid out randomly. Results showed that mortality and % mortality of weevils increased progressively with the powder concentrations and the exposure period. Within the 14 days of exposure, mortality and % mortality were 4, 20% in the control and all, 100% in the test treatments. Reduction in seed damage increased significantly with the powder concentration ($P<0.05$). Weevil perforation index (WPI) showed that the combined powder had high protectant activity and therefore could serve as an alternative to the conventional chemical insect pesticides.

Key Words: *Xylopiya aethiopica*, *Aframomum melegueta*, *Sitophulus zea*, combined powder.

INTRODUCTION

High biotic pressure, on agro-food crops such as maize in storage, is known to be responsible for post harvest loss, experienced by farmers. Biodeterioration of stored grains, both marginal and comprehensive, have been traced to the activities of insect pests such as the maize weevils. (Rahman, et al., 2001, Ewete and Alamu, 1999). Maize weevil *Sitophilus zea* (S.zeamals. Motschulsky). S. *zea* is a relatively small, brown, thick winged, snouty beetle, belonging to the sub family Dryophthorinae, in the family, Curculionidae, orders Coleoptera, of the insecta class and phylum Arthropoda. This pest had been known to attack maize, both in the field and in storage. Maize is binomially known as *Zea mays* L. and is commonly called Corn. They belong to the grass family Poaceae formerly Gramineae. Of the monocotyledonous class, they are annuals, produce edible grains born on a woody cob. The cob is the axis of the spike flowers of the female inflorescence, born on axil of the isobilateral leaf. The male inflorescence is terminal. The Spikelets, are pairs of two flowers, one sessile one Pedicelled. (Nyananyo, 2006). Corn is a staple food, and forms the bulk of cereal food for both children and adults. It is grown commercially and stored in Granaries and

Silos. Unfortunately, due to the bio activities of weevils, farmers and corn dealers do suffer losses from time to time and from season to season. (Rahman et al., 2001). In other to address these losses, farmers had resorted to the use of chemical insecticidal pesticides. (El-Atta and Ahmed, 2002). However, these chemicals have attendant hazards like low affordability,,resistance,,environmental pollution, and other bio- hazards (UNEP,2002a, 2002b, Ayinde et al.,2006) The application of botanicals, as alternative substitutes to the conventional chemical agents has been reported by many researchers like Akueshi,et al.,2002, Ogonna et al.,2003,and Chomini et al.,(2010).

This study therefore is aimed at determining the antifeedant potentials of combined powder of the fruits and seeds of *Xylopi aethiopia* and *Afromomum melegueta* .

MATERIALS AND METHODS

Fruits of the test plants *Afromomum melegueta* and *Xylopi aethiopia* were collected from the relief market owerri.The maize grains were collected from dealers' from the same market. The weevils were isolated from bags of weevil infested grains from old stock, of maize in storage or in ware houses adjacent to the free zone area of the market. The specimen was taken to the Department of Biological Sciences of the Federal University of Technology, Owerri where the study was conducted. The weevils were then identified as *Sitophilus zea*.

Owerri is the capital city of Imo State, South Eastern Nigeria. It is located within latitudes 40 45- N and 7⁰ 05-N and longitude 6⁰ 50-E and 7⁰ 25-E in the rain forest belt of Nigeria. The study was conducted between the months of April to September 2013. The test fruits were oven dried to a constant weight at a temperature of 40^{0C}. Pulverized and kept dried and sterile. The maize grains were refrigerated for 72 hours, following, methods of Chomini et al.,(2006). The two powder were mixed in equal proportions and dispersed in concentrations of 0.0g, 1, 2, 3, 4, 5, 6 ,7, 8, 9, and 10g into Kilner jars , into which 100g of the disinfected grains had been dispensed. The powder and the seeds were thoroughly mixed through manual agitation for 5mins and allowed to stand for 60 minutes. 20 weevils were then introduced into the jars. The jars were then lined with muslin cloth.

Each treatment was replicated five times, and arranged randomly. Weevil mortality and % mortality were determined from the number of death at every 2 days, for 14 days after treatment. Seed damage at 14 days after treatment were determined using the relation,

$$WPI = \frac{\% \text{ treated seed perforated}}{\% \text{ control seed perforated}} \times \frac{100}{1}$$

Where WPI = weevil perforated index. (Fatope, et al., 1995)

Data was subjected to analysis of variance, using standard procedure (Kelly and Onyeka, 1992). Means were segregated using least significant difference (LSD) at 5%.

RESULTS AND DISCUSSIONS

Table 1: Mean and Mean % Mortality of *Sitophilus zea* treated with the combined fruit powder of *X. aethiopica* and *A. melegueta*.

Powder Conc.(g)	DAYS AFTER TREATMENT							DAYS AFTER TREATMENT						
	MM mean mortality							M % Mortality						
	2	4	6	8	10	12	14	2	4	6	8	10	12	14
0.00	0.00	1.0	2.00	2	2	4	4	0.00	10	10	10	10	20	20
1.00	4	6	10..2	12	16	17	20	20	30	51	60	80	85	100
2.00	9	13	14.2	16	18	20	20	45	65	71	80	90	100	100
3.00	10	18.2	20	20	20	20	20	50	91	100	100	100	100	100
4.00	12	18.6	20	20	20	20	20	60	93	100	100	100	100	100
5.00	12.2	20	20	20	20	20	20	61	100	100	100	100	100	100
6.00	13.0	20	20	20	20	20	20	65	100	100	100	100	100	100
7.00	14.6	20	20	20	20	20	20	73	100	100	100	100	100	100
8.00	16.0	20	20	20	20	20	20	80	100	100	100	100	100	100
9.00	16.2	20	20	20	20	20	20	81	100	100	100	100	100	100
10.00	17.4	20	20	20	20	20	20	87	100	100	100	100	100	100

Table 2: Mean number and % of seed perforation by *Sitophilus zea*, 14days after treatment with combined powder of *X. aethiopica* and *A. melegueta*.

Powder Conc.	Seed Perforation(sp)	No of seed	%Sp	WPI
0.00	95	500	19	
1.00	35	500	7	36.80
2.00	13	500	2.6	13.68
3.00	6.0	500	1.2	6.31
4.00	6.0	500	1.2	6.31
5.00	00	500	00	0.00
6.00	00	500	00	0.00
7.00	00	500	00	0.00
8.00	00	500	00	0.00
9.00	00	500	00	0.00
10.00	00	500	00	0.00

The insecticidal effect of the combined powder of *X. aethiopica* and the *A. melegueta*, on the test weevil, is shown in Table 1. Mortality and % mortality, increased with increase in concentration and the exposure period, of 2-14days. At 14 days, after treatment, all the

weevils had died significantly -100% mortality, except the control which had 20% mortality. Mortality rate, increased significantly, from 0.0g/100g seed concentration to 10g/100g seed concentration. Starting from the 5th degree, concentration, and all the weevils died within 4 days after exposure. Weevil perforation activities, against the concentration gradient are shown in Table 2. At the 14th day after treatment, the concentration of 5g/100g (w/w) to 10g/100g weevil perforation activity seized. The perforation activities decreased from 36.8% in 1g /100g to 6.3% in 4g/100g (w/w).

DISCUSSION

The use of combined powder of *X. aethiopica* and *A. melegueta* in this study was informed by the application of the botanicals, in the local preservation of grain crops. The test results on the mortality and %s mortality of the weevil in (table 1), corroborated with the findings of Chomini, et al., (2010). This showed increase in mortality and % mortality rate of bean weevils, *Callosobruchus maculatus*, with increase in concentration of *X. aethiopica* fruit powder. The efficacy of this combined powder is justified by the relative decrease in the weevil perforation index (table 2). This is akin to the finding of Yusuf, et al.,(2006) working on Neem and Mahogany bark extracts, against the bean beetle, *C. maculatus*. *X. aethiopica* is commonly called African pepper. It is a perennial tree, the fruits appear in clusters of numerous narrow carpel. The plant is a member of the Annonaceae family. *A. melegueta*, is commonly called “**Grain of paradise**” or alligator pepper. The fresh fruit is usually red with numerous brownish seeds embedded in whitish sugary pulp. Seed is usually chewed with kola nut, as spice. The plant belongs to Zingiberaceae family. Both fruits are sold in vegetable shops.

CONCLUSION:

To the extent of this study, it is evident that the combined powder of *X. aethiopica* and *A. melegueta* have high antifeedant activity against the test insects and therefore can be a good grain protectant, especially in storage.

REFERENCES

- Akueshi, C.O, Comfort O, Kadiri, E.U, Akueshi, S.E.A and Ngurukwem C (2002). Antimicrobial Potentials of *Hyptis suaveolens* poit (Lamiaceae) Nig .J. of Botany 15:37-41.
- Anjinde, I.A., Okuneye, P.A., Aromolaran, A. and Momoh, B.S. (2006). Health Implication of insecticide use in Cowpea production in Kano State, Nigeria: A Legit Analysis in Adepoju, S.O and Okuneye (eds) .Proceedings Of the 20th Annual national conference, FAMAN. FRIN, Federal College of Forestry Jos, 18th-21th September, 2006. Pp 607.
- Chomini ,M.S., Samndi, M.A., Akalusi, M.E., Okoiyele, L.O., Nwawkwushue, K.N. and Karma, I.B. (2006).Insecticidal effects of leaf, stem, bark and seed Kernel powdered extracts of Neem (*Azadirachta Indica* Ajuss.) Against Maize weevil (*Sitophilus zeamais matschulsky*).Nigerian Journal of Botany 19(2): 348-352.
- Chomini, M.S, Okoiyele, L.O, Chomini, A.E, and James, J.D. (2010) Efficacy of *Xylopiia aethiopica* Dunal. Powder in the control of Cowpea Brunchials, (*Callosobruchus macuratus*) FAB. Nigerian Journal of Botany 23(2) 235-244.

- Chomini, M.S., Akalusi, M.E., Okonkwo, M.C. and Musa, H. (2010). Effects of Cashew nutshell powder extract on *Callosobruchus maculatus* (F) in Stored cowpea. (*Vigna Unguiculata* L.) Walp. Nigerian Journal of Botany 23(2) 245-252.
- El-Atta, H.A., and Ahmed, A., (2002). Comparative effect of some botanicals for the control of the seed weevil *Carydo Serratus*. Oliver (col. Bruchidae.) Journal of Applied Entomology. 128 (10):577-583.
- Ewete, F.K and Alamu, O.T. (1999). Extracts of three Mahogany species, as grain Protectants against *Sitophilus Zeamais* Matschulsky. (Col:Curculinidae) Journal of tropical forest resources 157:22-29.
- Fatope, M.O, Mann, A. and Takeda, Y. (1995). Cowpea weevil bioassay. A simple Prescreen for plants with grain protectant effects. International Journal of Pest management 41:44-56.
- Kelly, M.G. and Onyeka, J.D.A. (1992). Introduction to Statistics and Experimental Design for life sciences. ABIC pub. Enugu, Nigeria 187pp.
- Nyananyo, B.L., (2006). Plants from Niger Delta. Onyoma research publications Nigeria pp 225
- Ogbonna, A.I, E nukora, E.I, Olorunfemi, P.O., Aguiyi, J.C., Ikwere, E.O, Akueshi, E.U, and Onyekwelu, N.A. (2003). Antibacterial activities of the aqueous Leaf extract of *Ximenia ameriesna*. Nigerian Journal of Botany 16:151-155.
- Rahman, S.A., Agada, J.E and idoko, O.J., (2001). Neem extract: The natural Insecticides for profitable and sustainable cowpea production in Northern Nigeria. Journal of Agricultural Technology 9:(1) 42-45.
- UNEP (2002). Integrated assessment of trade liberalization and trade related Policies: A country study on the export sector Nigeria UNEP/ETB/2002/a. United Nations publication, New York, Geneva 128pp.
- UNEP (2002b). Integrated assessment of trade liberalization and trade related Policies. A country study of on the cotton sector in China. United Nations Publication, New York, Geneva 66pp.
- Yusuf, S.R., Ahmed, B.I. and Galadima, B., (2006). Control of *Sitophilus zeamais* Mots. Adult population and damage on grain maize (*Zea mays* (L.) B. two Plant products compared with Pirimiphos Methyl in Adepoju, S.O and Okuneye, P.B. (Eds) Proceedings of 20th Annual Conference FAMAN. FRIN. Federal college of forestry Jos 18th-21st September, 2006 pp.621-624

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

