

Grain Yield Stability and Phenotypic Correlation Analysis of Bread Wheat (*Triticum aestivum* L.) Genotypes in North Western Ethiopia

Misganaw Ferede¹ Fisseha Worede²

1. Cereals Breeder, Amhara Regional Agricultural Research Institute, Adet Agricultural Research Center
P.O.Box 08, Bahirdar, Ethiopia

2. Lowland cereals Breeder, Amhara Regional Agricultural Research Institute, Sirinka Agricultural Research Center P.O.Box74, Woldia, Ethiopia

Abstract

An experiment was conducted at Adet, Simada and Debretabor experimental sites of Adet Agricultural Research Center in 2014 and 2015 cropping season under rain-fed condition to study stability and traits correlation of bread wheat genotypes. Twelve bread wheat genotypes were used as experimental treatments. The genotypes were laid out in randomized complete block design with three replications per site. The results of AMMI analysis depicted significant differences among genotypes across environments (locations and years) ($P \leq 0.001$). According to the study, the performances of genotypes grain yield were highly affected by environments (locations and years) and the genetic composition of genotypes. The highest variation was accounted for by location (29 %) followed by genotype (18%) and location by year (18 %) and genotype by year (12%) effects. Based on AMMI, GGE biplot and stability coefficient analyses, G4, G2, G11 and G9 were wide adaptable and relatively stable genotypes all over the test environments (locations and years) than the checks, G7 (TAY) and G12 (Kubsa). Therefore, based on the adaptability and stability and overall mean grain yield of genotypes, recently released genotypes Gambo (G4), Ogolcho (G2) and Tsehay (G9) and relatively older genotypes Shorima (G11) and TAY (G7) could be recommended for production at test environments in the Western Amhara Region. However, there is a need to study the effect of environmental variation on the occurrence of rust disease.

Keywords: AMMI, Correlation, GGE, Yield stability

1. INTRODUCTION

Ethiopia is frequently exposed to food shortage due to environmental variability, degradation of soil fertility, ever increasing of the population (Ashenafi, 2008) and inappropriate use of improved technologies (Zerihun, 2014). Even though the production and productivity of wheat crop in Ethiopia has increased in the last decades, the national average yield has not exceeded 2.54 tones/ha (CSA, 2014). It is lower than the world's average yield/ha which is about 3.3 tones/ha (FAO, 2014). This is due to factors such as use of unimproved low yielding varieties, uneven distribution of rainfall, poor agronomic practices and serious diseases like stem rusts (Dereje *et al.*, 2000).

Among the factors limiting wheat production and productivity, diseases are the foremost critical constraints all over the world. Particularly rust diseases cause highest yield loss of wheat production. A number of studies illustrated that wheat rusts could cause yield losses of 20-100 % on susceptible wheat genotypes (CIMMYT, 1989; Temesgen *et al.*, 1995; Emebet *et al.*, 2005; Stubbs, 1998; Marshall, 1988). Now that Ethiopia's wheat production covers only 75% of the national demand, the remaining 25% of the wheat is fortified through imported (Eyob *et al.*, 2014). Hence to overcome wheat yield losses and to cut down wheat national demand deficiency conducting considerable research works that contribute positive impact on wheat productivity and production are mandatory.

The process of variety development in the country is continuing year after year through various research institutes and universities. However, once released for production, the varieties are used for a long period of time continuously without considering their adaptation domain, grain yield stability and testing whether they are losing their potential or not. Now a day high yielding and rust disease resistant bread wheat varieties have recently been released in Ethiopia. However, farmers in Western Amhara Region commonly use relatively older bread wheat varieties such as Kubsa and TAY which were released in 1995 and 2005, respectively. Therefore, evaluation of recently released bread wheat varieties across environments and over years through different statistical methods enables to identify genotypes with better performance.

Hence, it is vital to evaluate grain yield stability and correlation of parameters of bread wheat genotypes used in the region with the following objectives: to evaluate the extent of grain yield stability, to assess the grain yield advantage of recently released varieties over farmers commonly used varieties and to investigate the relation of parameters to grain yield for future breeding program.

2. MATERIALS AND METHODS

2.1. Study Areas Description and Experimental Treatments

The experiment was conducted during 2014 and 2015 cropping season under rain-fed conditions at experimental sites of Adet Agricultural Research Center namely Adet, Simada and Debretabor. Twelve improved bread wheat genotypes were used as treatments for the study. The detail agro-ecological data of environments and the description of genotypes are listed in Table 1 and 2, respectively.

Table 1. Altitude, geographical location, soil type and climate data of the research sites

| Locations | Code | Cropping season | Code | Environments | Altitude (masl) | Geographical location | | Soil type | Climate data | |
|------------|------|-----------------|------|--------------|-----------------|-----------------------|----------------------|-----------|--------------|-----------------------|
| | | | | | | Latitude | Longitude | | RF(mm) | Temp(C ⁰) |
| Adet | Loc1 | 2014 | Yr1 | E1 | 2240m | 11 ⁰ 16'N | 37 ⁰ 29'E | Nitosol | 658.6 | 17.53 |
| Simada | Loc2 | 2014 | Yr1 | E2 | 2460m | 11 ⁰ 03'N | 37 ⁰ 30'E | Luvisol | 736.1 | 13.27 |
| Debretabor | Loc3 | 2014 | Yr1 | E3 | 2591m | 11 ⁰ 51'N | 38 ⁰ 01'E | Luvisol | 1102.7 | 15.48 |
| Adet | Loc1 | 2015 | Yr2 | E4 | 2240m | 11 ⁰ 16'N | 37 ⁰ 29'E | Nitosol | 948.9 | 19.4 |
| Simada | Loc2 | 2015 | Yr2 | E5 | 2460m | 11 ⁰ 03'N | 37 ⁰ 30'E | Luvisol | 770.6 | 15.07 |
| Debretabor | Loc3 | 2015 | Yr2 | E6 | 2591m | 11 ⁰ 51'N | 38 ⁰ 01'E | Luvisol | 958.1 | 15.94 |

Source: AARC (2014) and ANRSMA (2014 and 2015)

Note: Total amount of rainfall and average temperature from July to November

Table 2. Description of bread wheat genotypes used for the study

| Genotypes name | Code | Released by | Year of release | Grain yield (t/ha) during releasing time | | Adaptation Zone | |
|--------------------------------|------|-------------|-----------------|--|---------|-----------------|----------|
| | | | | On station | On farm | Altitude(masl) | RF(mm) |
| Hidase(ETBW 5795) | G1 | KARC | 2012 | 4.4-7 | 3.5-6 | 2200-2600 | >500 |
| Huluka(Flag 5) | G3 | KARC | 2012 | 4.4-7 | 3.8-6 | 2200-2600 | 500-800 |
| Ogolcho(ETBW 5520) | G2 | KARC | 2012 | 2.8-4 | 2.2-3.5 | 1600-2100 | 400-500 |
| Shorima(ETBW 5483) | G11 | KARC | 2011 | 2.9-7 | 2.3-4.4 | 2100-2700 | 700-1100 |
| Gambo(QUIAU#2) | G4 | KARC | 2011 | 3.5-5.7 | 4.5 | 750 | - |
| Tsehay(HAR 3837) | G9 | DBARC | 2011 | 3.8 | 2.8-3.5 | 2600-3100 | >900 |
| Danda'a(DANPHE#1) | G5 | KARC | 2010 | 3.5-5.5 | 2.5-5 | 2000-2600 | >600 |
| Bolo(HAR 3816) | G8 | DBARC | 2009 | 2.8-3.5 | 2.3-3.3 | 2580-3100 | >904 |
| Menze(HAR 3008) | G10 | DBARC | 2007 | 1.9-3.3 | 1.5-2.7 | 2800-3100 | >904 |
| Gassay(HAR 3730) | G6 | ADARC | 2007 | 4.4-5 | 3.5-4.7 | 1890-2800 | >700 |
| TAY (ET-12 D4/ HAR-604(1) (SC) | G7 | ADARC | 2005 | 2.5-6.1 | 3.4-5.8 | 1900-2800 | >700 |
| Kubsa(HAR 1685) (SC) | G12 | KARC | 1995 | 5.8-6.3 | 4-4.5 | 1850-2800 | 500-800 |

Source: MoA, Crop Variety Register (1995-2012)

ADARC- Adet Agricultural Research Center, DBARC- Debrebirhan Agricultural Research Center, KARC- Kulumsa Agricultural Research Center, SC-Standard Check, SC-Standard Check

2.2. Experimental Procedures and Method of Statistical Analysis

The treatments were laid out using a randomized complete block design with three replications per site and six rows per plot. Planting was done in the second week of July with seeding rate of 150 kg/ha on the plot area of 1.2 m*2.5 m with net harvested area of 0.8m*2.5m. Urea and DAP fertilizers as source of nitrogen and phosphorous were applied as per their recommendation rate and time of application for bread wheat specified to each experimental site. All other agronomic practices like weeding were applied uniformly for all the treatments at all experimental sites.

Grain yield was analyzed by using GenStat (17th Ed) software to compute analysis of genotypes and environments main and interaction effects, seasonal variation effects and grain yield stability of genotypes. Whenever the analysis results were highly-significant or significant, Fisher's LSD test at 1 % and 5% probability level, respectively, was used to separate the variable means of genotypes, environments and genotypes by environments interaction.

The AMMI analysis of variance summarizes most of the magnitude of genotype by environment interactions into one or few interaction principal component axes (IPCA) (Zobel *et al.* 1988, Crossa, 1990). The following AMMI model equation was used:

$$Y_{ger} - \mu - \alpha_g - \beta_e = \sum_n \lambda_n \tau_{gn} \delta_{en} + p_{ge} + \xi_{ger}$$

where Y_{ger} is the grain yield of genotype (g) in environment (e) for replicate (r), μ is the grand mean, α_g are genotype mean, β_e are the environment mean deviations, λ_n is the singular value for IPCA axis n, τ_{gn} are genotype eigenvector values for IPCA axis n, δ_{en} are the environment eigenvector values for (PCA) axisn, p_{ge} are the residuals and ξ_{ger} is the error term.

GGE biplot analysis was carried out to identify high yielding and stable varieties as well as representative and discriminating environments as per Yan (2001):

$$Y_{ger} - \beta_e = \sum_n \lambda_n \tau_{gn} \delta_{en} + p_{ge} + \xi_{ger}$$

where Y_{ger} is the grain yield of genotype (g) in environment (e) for replicate (r), β_e are the environment mean deviations, λ_n is the singular value for IPCA axis n, τ_{gn} are genotype eigenvector values for IPCA axis n, δ_{en} are the environment eigenvector values for (PCA) axis n, p_{ge} are the residuals and ξ_{ger} is the error term.

AMMI Stability Value (ASV) is the distance from the coordinate point to the origin in a two-dimensional plot of IPCA1 scores against IPCA2 scores in the AMMI model (Crossa, 1990). ASV was calculated for each genotype and each environment according to the relative contribution of IPCA1 to IPCA2 to the interaction sum of squares as follows:

$$ASV = \sqrt{\left[\frac{SS_{IPCA1}}{SS_{IPCA2}}(IPCA1score)\right]^2 + (IPCA2score)^2}$$

Lin and Binns (1988) defined the superiority measure (P_i) of the i^{th} test cultivar as the MS of distance between the i^{th} test cultivar and the maximum response as:

$$P_i = \left[n(\bar{X}_i - \bar{M})^2 + \left(\sum_{j=1}^n (X_{ij} - \bar{X}_i - M_j + \bar{M})^2 \right) / 2n \right]$$

Where X_{ij} is the average response of the i^{th} genotype in the j^{th} environment, \bar{X}_i is the mean deviation of genotype i, M_j is the genotype with maximum response among all genotypes in the j^{th} location, and n is the number of locations. The first term of the equation represents the genotype sum of squares and the second part represents the GE sum of squares.

Becker and Leon (1988) defined the concept of ecovalence (W_i) as the contribution of each genotype to the GEI sum of squares. The W_i or stability of the i^{th} genotype is its interaction with the environments, squared and summed across environments, and expressed as:

$$W_i = [Y_{ij} - Y_i - Y_{.j} - Y_{..}]^2$$

Where, Y_{ij} is the mean performance of genotype i in the j^{th} environment, Y_i and $Y_{.j}$ are the genotype and the environment mean deviations, respectively, and $Y_{..}$ is the overall mean.

According to Lin *et al.* (1986) the variance of genotype yields recorded across the test environments can be used as a measure of stability. For the genotype greatest stability is $S_i^2=0$. The formula is:

$$S_i^2 = \sum (R_{ij} - m_i)^2 / (e-1)$$

Where; S_i^2 = environmental variance, R_{ij} = observed genotype yield across environments, m_i = marginal means of genotypes, e = number of environments

3. RESULTS AND DISCUSSION

3.1. Impact of Genotypes, Locations and Years on grain yield

The analyses of variances depicted highly significant ($P \leq 0.001$) differences among genotypes, locations, years and their interactions for grain yield (Table 3). The highest variation was accounted for by location (29 %) followed by genotype (18%) and location by year (18 %) and genotype by year (12%) effects (Table 3). The grain yield of genotypes was highest at Adet in 2014 cropping season, and at Debretabor in 2015 cropping season. Similarly, grain yield of genotypes was lowest at Simada in 2015 (Table 4). Genotype 4 (Gambo) was the highest yielder both at the highest (Adet in 2014) and the lowest (Simada in 2015) yielding environments. Genotypes G4, G2, G11 and G9 showed 12.41, 10.22, 4 and 3.45 qt/ha grain yield advantage over standard check (G7) respectively, and 17.16, 14.97, 8.75 and 8.2 qt/ha grain yield advantage over standard check (G12), respectively (Table 4). Based on this study, grain yield response of genotypes was highly affected by location, environments (location by seasonal variation) and seasonal variation in line with Frey (1983) and Falconer (1990). As a result, screening and development of wide adaptable and relatively stable genotypes are determinant factor to increase bread wheat productivity and production.

Table 3. The analysis of variance of genotypes, locations, years and their interactions

| SV | DF | SS | MS | % SS (gen+loc+yr+gen*loc+gen*yr+loc*yr+gen*loc*yr+ residual) |
|------------|-----|----------|-----------|--|
| Gen | 11 | 9187.87 | 835.26** | 18 |
| Loc | 2 | 14393.4 | 7196.7** | 29 |
| Yr | 1 | 2764.46 | 2764.46** | 5.6 |
| Gen*Loc | 22 | 3798.1 | 172.64** | 7.6 |
| Gen*Yr | 11 | 5809.92 | 528.17** | 12 |
| Loc*Yr | 2 | 8998.73 | 4499.37** | 18 |
| Gen*Loc*Yr | 22 | 2591.46 | 117.79** | 5.2 |
| Residual | 142 | 2247.1 | 15.82 | 4.5 |
| Total | 213 | 49791.04 | | |

Gen-genotype, Loc- Location, Yr- Year, SV- Source of variation

Table 4. Genotypes mean grain yield (qt/ha) across locations and over years

| Genotypes | Loc1 | | Loc2 | | Loc3 | | Mean |
|-----------------|------------------|--------------|------------------|--------------|------------------|--------------|---------------------|
| | Yr1 | Yr2 | Yr1 | Yr2 | Yr1 | Yr2 | |
| G1 | 46.68 | 42.59 | 42.94 | 18.52 | 18.4 | 54.45 | 37.26 ^{de} |
| G2 | 64.67 | 58.19 | 51.96 | 25.91 | 48.03 | 49.13 | 49.65 ^a |
| G3 | 57 | 53.17 | 24.55 | 21.21 | 28 | 43.42 | 37.89 ^{cd} |
| G4 | 70.85 | 60.62 | 56.86 | 26.37 | 49.35 | 47 | 51.84 ^a |
| G5 | 52.98 | 30.86 | 44.86 | 16.73 | 37.01 | 36.47 | 36.48 ^{de} |
| G6 | 44.95 | 38.06 | 42.73 | 20.56 | 32.54 | 39.12 | 36.33 ^{de} |
| G7 | 59.15 | 52.24 | 27.98 | 17.84 | 30.96 | 48.74 | 39.49 ^e |
| G8 | 50.64 | 13.88 | 34.93 | 14.67 | 35.91 | 33.66 | 30.61 ^f |
| G9 | 66.78 | 45.48 | 45.85 | 17.74 | 35.31 | 46.11 | 42.88 ^b |
| G10 | 52.73 | 11.98 | 36.6 | 14.04 | 27.68 | 33.78 | 29.47 ^f |
| G11 | 56.13 | 51.55 | 46.12 | 23.52 | 38.22 | 45.06 | 43.43 ^b |
| G12 | 49.68 | 55.73 | 12.75 | 22.01 | 17.07 | 50.85 | 34.68 ^e |
| YrMean | 56.02 | 42.86 | 39.01 | 19.93 | 33.21 | 43.98 | |
| Loc mean | 49.44 | | 29.47 | | 38.59 | | 39.17 |
| LSD | 7.62 | | 4.38 | | 6.95 | | 6.42 |
| CV | 9.4 | | 9 | | 11 | | 10.2 |
| P level | <0.001 | | <0.001 | | <0.001 | | |

G1=Hidase, G2= Ogoloch, G3=Hulluka, G4= Ga'ambo, G5= Danad'a, G6= Gassay, G7=Tay, G8= Bolo, G9= Tsehay, G10=Menze, G11=Shorima, G12=Kubsa, Loc1=Adet, Loc2= Simada, Loc3= Debretabour, Yr=Year, Loc=Location, Yr1=2014, Yr2=2015

3.2. The Main and Interaction Effects of Genotypes and Environments

The AMMI analysis of grain yield showed highly significant differences among genotypes, environments and their interactions (Table 5). Environments depicted the highest variation on grain yield performance of genotypes which accounted for 41.9% followed by genotypes (14.7%), and genotype by environments interaction (19.5%) which was in agreement with the findings of Misganaw *et al.* (2015) (Table 5).

The partitioning of the genotype-environment interaction by employing AMMI model analysis showed that four of the Interaction Principal Component Axes (IPCA) were highly significant ($P < 0.001$). The four IPCAs accounted for 99.2% of the interaction sum of squares (SS), with 87.27% of the corresponding degrees of freedom (Table 5). Out of the four, the first interaction PCA captured 66.54% of the interaction SS with 27.27% of the corresponding degrees of freedom. Similarly, the second interaction PCA captured 15.72% of the interaction SS with 23.4% of the corresponding degrees of freedom. The two interaction PCAs explained 82.26% of the total variation in grain yield of the bread wheat genotypes. The AMMI1 biplot captured 43462 of the treatment SS of 60462. Therefore, it revealed 71.88% of the treatment SS. Approximately as much variation in grain yield was explained by the interaction term captured by IPCA1 as by the genotypic main effect. This showed that interaction is as important as genotypic main effect; implying that both specific and wide adaptations are important. In the biplot axes system, either main effects and IPCA1, or IPCA1 and IPCA2 are commonly used as abscissa and ordinates (Zobel *et al.*, 1988; Gauch, 1992).

The AMMI biplot showing the main and IPCA1 effects of both genotypes and environments on bread wheat grain yield is depicted in Figure 1. In such a system, distances along the abscissa (horizontal line) shows main effect differences, whereas the ordinate (vertical line) shows differences in interaction. In the present study,

G1, G3, G5 and G6 had more or less similar genotypic main effect, differing in interaction. However; G2 and G11, and G4 and G9 had nearly similar interaction effects, only differing in genotypic main effects. In the same manner, E4 and E2 had higher interaction effect; whereas E5 and E1 had minimum interaction effect, but they had higher differences in environmental main effect. According to Gauch and Zobel (1996) and Gauch and Furnas (1991), when a genotype and an environment have the same sign on the IPCA1 axis, their interaction is positive i. e., that particular genotype is adaptable and the environment is conducive for the genotype; if different, their interaction is negative. Accordingly G2, G4, G5, G6, G8, G9 and 10 were adapted to E1, E2 and E3. In contrast, G1, G3, G7, G11 and G12 were adapted to E4, E5 and E6.

Figure 2, in addition to delineating mega-environments, showed the interaction pattern of the 12 bread wheat genotypes with the six environments. The distances from the origin indicate the magnitude of interaction exerted by environments on genotypes, or vice versa (Voltas *et al.*, 2002). In other words, genotypes near the origin are not sensitive to environmental interaction, whereas genotypes distant from the origin are sensitive and have large interaction effects. Hence from this study, genotypes G11, G9, G6, G7 and G3 were weakly influenced by environmental factors while G10, G8, G12 and G4 were strongly affected by environmental factors. Therefore, genotypes which buffer variable environmental factors are more or less stable. Inversely, genotypes performance which varies due to environmental factors is specific adaptable.

Table 5. AMMI analysis of variance for genotypes, environments and their interactions based on Grain Yield Response

| SV | DF | SS | MS | % of SS (Gen+Env+ Gen*Env+IPCA1+2+3+4) |
|--------------|-----|-------|----------|--|
| Genotypes | 11 | 9188 | 835.3** | 14.7 |
| Environments | 5 | 26157 | 5231.3** | 41.9 |
| Block | 12 | 718 | 59.8** | 1.15 |
| Interactions | 55 | 12199 | 221.8** | 19.5 |
| IPCA 1 | 15 | 8117 | 541.1** | 13 |
| IPCA 2 | 13 | 1918 | 147.5** | 3.07 |
| IPCA 3 | 11 | 1476 | 134.2** | 2.37 |
| IPCA 4 | 9 | 592 | 65.8** | 0.95 |
| Residuals | 7 | 97 | 13.8 | 0.16 |
| Error | 132 | 1943 | 14.7 | 3.11 |
| Total | 270 | 62405 | 7265.3 | |

SV-Source of variation, Gen-Genotypes, Env- Environments, IPCA-Interaction principal component axes




Figure 1. AMMI biplot of main effects of genotypes and environments using symmetrical scaling
 G1=Hidase, G2= Ogocho, G3=Hulluka, G4= Gaambo, G5= Danda'a, G6= Gassay, G7=Tay, G8= Bolo, G9= Tshay, G10=Menze, G11=Shorima, G12=Kubsa, E1 and E4= Adet E2 and E5=Simada and E3 and E6= Debretabor, IPCA= Interaction Principal Component Axes

3.3. Clustering of Environments based on Genotypes Grain Yield Response

The GGE biplot is useful for identification of mega-environments, ideal genotype and test environments, among other things. Based on GGE biplot, environments were grouped into two mega environments. Mega environment one includes E1, E2, E3, E4 and E5 and mega environment two had singleton environment E6. Genotype G2 was the winning one across environments E1, E2, E3, E4 and E5 (Figure 2).

According to Yan *et al.* (2000) and Yan and Rajcan (2002), ideal genotypes are those having large PC1 scores (high grain yield) and small absolute PC2 scores (high stability). Accordingly G11 G6, G2 and G9 were better stable genotypes. Genotypes G4, G2, G11 and G9 were high-yielder in that order of importance. Though G6 was relatively stable genotype, it not preferable for production due its low-yielding capacity.

Ideal environments should be more representative of the entire set of environments and should have more genotype discriminating power, such environments should have small PC2 scores (absolute) and large PC1 scores (Yan *et al.*, 2000; Yan and Rajcan, 2002). Thus E5 and E1 are more representative, whereas E5, E4 and E1 are more discriminating environments.




Figure 2. GGE biplot analysis of genotypes and environments using environment scaling

G1=Hidase, G2= Oголоcho, G3=Hulluka, G4= Gaambo, G5= Danda'a, G6= Gassay, G7=Tay, G8= Bolo, G9= Tsehay, G10=Menze, G11=Shorima, G12=Kubsa, E1 and E4= Adet E2 and E5=Simada and E3 and E6= Debretabor, PC= Principal Component

3.4. Grain Yield Stability of Genotypes over Environments

AMMI and GGE biplot analyses show adaptable and stable genotypes in graphical forms. As a matter of fact, to know the stability of genotypes in numerical values further stability analysis works are pertinent to explore stable genotypes using different stability analyses methods. According to this study, grain yield stability of genotypes rank varies with the methods used. Based on statements of Becker and Leon (1988) and Crossa (1990) genotypes with a low W_i value have smaller deviations from the mean across environments and are thus more stable and the larger the ASV value, either negative or positive, the more specifically adapted a genotype is to certain environments while smaller ASV values indicate more stable genotypes across environments. Therefore G4, G2, G11 and G9 were relatively stable genotypes based on AMMI stability, cultivar superiority and static stability analyses values (Table 6).

Static stability analysis had a drawback which implies both higher and lower grain yielding genotypes as stable. AMMI stability value shows consistency of genotypes contribution to genotype by environment interactions. While cultivar superiority analysis only showed mean performance of genotypes across environments, nonetheless it is difficult to know consistency of genotypes yield response across environments. This shows the necessity of combined use of different stability analysis methods to properly identify stable genotypes both in potential and consistency of grain yield over environments.

Table 6. Grain yield stability of bread wheat genotypes based on different methods of analyses

| Genotypes | Mean | AMMI stability | | Stability coefficient analysis methods | | | |
|------------|--------------|----------------|----------|--|----------|------------------|-----------|
| | | ASV | R | Cultivar Superiority | R | Static stability | R |
| G1 | 37.26 | 4.28 | 6 | 176.9 | 9 | 230.4 | 7 |
| G2 | 49.65 | 1.99 | 3 | 8.2 | 2 | 174 | 4 |
| G3 | 37.89 | 9.91 | 9 | 157.9 | 6 | 236.7 | 9 |
| G4 | 51.84 | 3.34 | 5 | 4.6 | 1 | 228.3 | 6 |
| G5 | 36.48 | 8.7 | 8 | 159.8 | 7 | 152.8 | 3 |
| G6 | 36.33 | 3.23 | 4 | 160.9 | 8 | 77.8 | 1 |
| G7 | 39.49 | 8.4 | 7 | 123.7 | 5 | 261.7 | 11 |
| G8 | 30.61 | 12.7 | 11 | 318.6 | 11 | 198.3 | 5 |
| G9 | 42.88 | 1.89 | 2 | 59 | 4 | 257.3 | 10 |
| G10 | 29.47 | 12.4 | 10 | 346.1 | 12 | 231.5 | 8 |
| G11 | 43.43 | 1.17 | 1 | 52.9 | 3 | 132.1 | 2 |
| G12 | 34.68 | 18.9 | 12 | 291 | 10 | 376.2 | 12 |

G1=Hidase, G2= Ogolocho, G3=Hulluka, G4= Gaambo, G5= Danda'a, G6= Gassay, G7=TAY, G8= Bolo, G9= Tshay, G10=Menze, G11=Shorima, G12=Kubsa, ASV=AMMI stability value, R=Rank

3.5. Phenotypic Correlation Analysis of Parameters

Correlation study helps to identify important traits in a breeding program and practice effective selection method. In this study, plant height, dry biomass, harvest index, thousand-seed weight and hectoliter weight depicted highly significant positive correlation ($r= 0.54, 0.9, 0.61, 0.73$ and 0.66) respectively to grain yield; whereas days to maturity, spike length, numbers of spikelet per spike and numbers of seeds per spike showed positive non significant correlation to grain yield ($P \leq 0.05$) (Table 6). Therefore, selecting genotypes with tall plant types, heavier above ground mass, larger seed size, effective assimilate partitioning capacity and higher flour return rate is important in attaining higher yield in bread wheat. Tkachur *et al.* (1978), Tayyar (2010), Mohibullah *et al.* (2011) and Beheshtizadeh *et al.* (2013) stated that plant height, spike length, number of spikelet per spike, number of seeds per spike, biomass yield and 1000-grain weight had positive correlation with grain yield. Grain yield is the result of comprehensive effects of traits. Hence, for bread wheat breeding works to be successful, breeders should consider the traits which have strong positive relation to grain yield.

Table 7. Phenotypic correlation of bread wheat genotypes for 10 traits across test environments in 2014 and 2015

| | DM | PHT | SL | NSLPS | NSPS | DB | HI | TSW | HLW | GYLD |
|-------|----|--------|---------------------|--------|--------------------|---------------------|--------------------|--------------------|--------------------|--------------------------|
| DM | 1 | 0.39** | -0.25** | 0.19** | 0.34** | -0.03 ^{ns} | 0.12 ^{ns} | -0.07 | 0.09 ^{ns} | 0.01^{ns} |
| PHT | | 1 | -0.11 ^{ns} | 0.43** | 0.39** | 0.65** | 0.08 ^{ns} | 0.3** | 0.39** | 0.54** |
| SL | | | 1 | 0.17* | 0.11 ^{ns} | 0.04 ^{ns} | 0.17** | 0.06 ^{ns} | 0.08 ^{ns} | 0.12^{ns} |
| NSLPS | | | | 1 | 0.62** | 0.18** | -0.14 | 0.01 ^{ns} | 0.09 ^{ns} | 0.07^{ns} |
| NSPS | | | | | 1 | 0.11 ^{ns} | 0.09 ^{ns} | 0.06 ^{ns} | 0.09 ^{ns} | 0.11^{ns} |
| DB | | | | | | 1 | 0.24** | 0.54** | 0.45** | 0.9** |
| HI | | | | | | | 1 | 0.71** | 0.65** | 0.61** |
| TSW | | | | | | | | 1 | 0.66** | 0.73** |
| HLW | | | | | | | | | 1 | 0.66** |
| GYLD | | | | | | | | | | 1 |

** = Highly significant at $P \leq 0.01$, * = Significant at $P \leq 0.05$ and ns= nonsignificant

DM= Days to Maturity, PHT= Plant Height, SL= Spike Length, NSLPS= Number of Spikelets per Spike, NSPS= Number of Seeds per Spike, DB= Dry Biomass, HI= Harvest Index, TSW= Thousand Seed Weight, HLW=Hectoliter Weight, and GYLD=GrainYield

4. CONCLUSION AND RECOMMENDATIONS

According to the study, genotypes' grain yield was highly affected by environments (locations and years) and the genetic composition of genotypes. The highest variation was accounted for by location (29 %) followed by genotype (18%) and location by year (18 %) and genotype by year (12%) effects. Genotypes' grain yield response was highest at Adet in 2014 cropping season, and at Debreabor in 2015 cropping season. Genotypes G4, G2, G11 and G9 were showed 12.41, 10.22, 4 and 3.45 qt/ha grain yield advantage over standard check (G7) respectively and 17.16, 14.97, 8.75 and 8.2 qt/ha grain yield advantage over local check (G12) respectively. Depending on AMMI, GGE biplot and different coefficients of stability, G4, G2, G11 and G9 were wide adaptable and relatively stable genotypes all over the test environments (locations and years) than the two checks. Therefore, these genotypes that had a higher mean grain yield in a wide range of environments are important to improve production

and productivity and crossing purpose due to their buffering capacity of the variability of environmental factors. In this study, plant height, dry biomass, harvest index, thousand seed weight and hectoliter weight were depicted highly significant positive correlation to grain yield. These traits should, therefore, be considered to improve grain yield in bread wheat breeding programs.

Therefore, based on the adaptability and stability of overall mean grain yield, recently released genotypes Gambo (G4), Ogolcho (G2) and Tsehay (G9) and relatively older genotypes Shorima (G11) and TAY (G7) could be recommended for production at the test environments in the Western Amhara Region. However, there is a need to study the effect of environmental/seasonal variation on rust disease occurrence.

5. REFERENCES

- Akter A., Hassen J. M., Kulsum U. M., Islam M. R., Hossain K. and Rahman M. M. 2014. AMMI biplot analysis for stability of grain yield in hybrid rice (*Oryza sativa* L.). *J. rice research* 2(2):1-4.
- Amhara National Regional State Metreology Agency (ANRSMA), 2015 and 2016, Climate data for Adet, Simada and Debretabor, Unpublished, Bahir Dar, Ethiopia.
- Ashenafi Gedamu. 2008. Triticale production in Ethiopia-its impact on food security and poverty alleviation in the Amhara Region, Kassel University, German. Pp 1-208.
- Becker H. C. and Leon J. 1988. Stability analysis in plant breeding. *Plant breeding* 101:1-23.
- Beheshtizadeh H., Rezaie A. and Ghandi A. 2013. Principal component analysis and determination of selection criteria in bread wheat genotypes, *International journal of agriculture and crop science* 5 (8):2024-2027.
- Central Statistics Agency (CSA). 2014/2015 (2007 E.C.). *Agricultural Sample Survey, Volume I, Report on Area and Production of Major Crops (Private Peasant Holdings, Meher Season)*. Statistical Bulletin 278. Addis Ababa, Ethiopia.
- CIMMYT. 1989. The wheat revolution revisited: Recent trends and future challenges. In the 1987/88 CIMMYT World Wheat Facts and Trends. Mexico, D. F. CIMMYT. 1-10.
- Crossa J. 1990. Statistical analysis of multi-location trials. *Advances in agronomy* 44: 55- 85.
- Dereje G. and Yaynu H. 2000. Yield losses of crops due to disease in Ethiopia. *Pest Mgt Journal Eth.* 5:55-67.
- Emebet F., Belayneh A. and Kassaye Z. 2005. Identification of sources of resistance to stem rust of wheat, *P. graminis* sp. *tritici*. Report of completed research project from 1999-2004, Volume I Pathology. PPRC, Ambo.
- Eyob Bezabeh, Tesfaye Haregewoin, Dejene Hailegiorgis, Fitsum Daniel and Baye Belay. 2014. Change and growth rate analysis in area, yield and production of wheat in Ethiopia. *International journal of development research* 4 (10):1994-1995.
- Falconer D.S. 1990. Selection in different environments: Effects on environmental sensitivity (reaction norms) and on mean performance. *Genet. Res. Camb.* 56:57-70.
- Fetien Abay and Asmund Bjornstad. 2009. Identifying optimal testing environments of barley yield in the Northern Highlands of Ethiopia by Biplot Analysis. *Journal of the Drylands* 2(1): 40-47.
- Food and Agriculture Organization (FOA). 2014. Food Balance Sheets. FAOSTAT. Rome. (<http://faostat3.fao.org/download/FB/FBS/E>).
- Frey K.T. 1983. Plant population management and breeding. In: Wood D.R. (editor). *Crop breeding. America society of agronomy. mandison, USA.* 55-88.
- Gauch H.G. 1992. Statistical analysis of regional yield trials: AMMI analysis of factorial designs. Elsevier, Amsterdam, the Netherlands.
- Gauch H.G. and Furnas R.E. 1991. Statistical analysis of yield trials with MATMODEL. *Agronomy Journal* 83: 916-920.
- Gauch H. G. and Zobel R.W. 1996. AMMI analysis of yield trials. In: Karg M. S. and Gauch H.G. *Genotype by environment interactions*. CRC press. Bacarator, Florida.
- Hints Gebbru and Fetien Abay. 2013. Evaluation of bread wheat varieties for their adaptability in wheat growing areas of Tigray region, northern Ethiopia. *Journal of biodiversity and endangered species.* 1:12-18.
- Lin C.S., Binns M.R. and Lefkovitch L.P. 1986. Stability analysis. Where do we stand? *CropSci.* 26: 894-899.
- Lin C.S. and Binns M.R. 1988. A superiority measure of cultivar performance for cultivar x location data. *Can. J. Plant Sci.* 68: 193-198.
- Marshall D. 1988. Characteristics of the 1984-1985 wheat leaf rust epidemic in central Texas. *Plant Dis.* 72:239-241.
- Misganaw Ferede, Getachew Alemayehu and Fisseha Wored. 2015. Identification of Adaptable Improved Bread Wheat (*Triticumaestivum* L.) Genotypes in North Western Ethiopia, *Journal of Biology, Agriculture and Healthcare*, 5(23).
- Misra R. C., DAS S. and Patnaik M. C. 2010. AMMI analysis of stability and adaptability of late duration finger millet (*Eleusine coracana*) genotypes. *World Journal of Agriculture Science* 6(6):664-669.
- Mohammed N.E. 2013. Genotype by environment interactions for grain yield in bread wheat (*Triticumaestivum*),

- Journal of plant breeding and crop science.5(7):150-157.
- Mohibullah M., Rabbini M.A., Jehan S., Zakiullah, Amin A. and Ghazanfarullah.2011. Genetic variability and correlation analysis of bread wheat (*Triticumaestivum* l.)accessions. Pak.J. Bot. 43(6): 2717-2720.
- Stubbs R. W. 1988. Pathogenicity analysis of yellow (stripe) rust of wheat and its significance in a global context.Pp. 23-38 in NW. Simmonds and S. Rajaram, eds. Breeding Strategies for Resistance to the Rusts of Wheat. CIMMYT: Mexico.
- Tayyar S. 2010. Variation in grain yield and quality of Romanian bread wheat varieties compared to local varieties in Northwestern Turkey.Romanian biotechnological letters.15
- Temesgen K., Bekele G., Balcha Y. and Ayele B.1995. Status of wheat rusts in the major producing regions of Ethiopia. In:Danial D.L. Proceedings of a regional workshop for Eastern, Central and Southern Africa, Njoro, Kenya, October 2-6, 1994. pp: 180-184.
- Tkachuk R. and Kuzine F.D. 1979. Wheat: Relations between some physical and chemical properties.Can.J.plant Sci. 59:15-20.
- Voltas J., Van E.F., Igartua E., García del Moral L.F., Molina-Cano J.L., Romagosa I. 2002. Genotype by environment interaction and adaptation in barley breeding: Basic concepts and methods of analysis. In Slafer G.A., Molina-Cano J. L., Savin R., Araus J. L., Romagosa I. (eds.) Barley Science: Recent advances from molecular biology to agronomy of yield and quality. The Harworth Press Inc., New York, pp. 205-241.
- Yan W. 2001. GGE biplot: a Windows Application for Graphical Analysis of Multi environment trial Data and other Types of Two-Way Data. Agronomy 93: 1111-1118.
- Yan W., HuntL.A., Sheng Q. and SzlavnicZ. 2000. Cultivar evaluation and mega-environment investigation based on the GGE biplot. Crop Science 40:597-605.
- Yan W. and RajcanI. 2002.Biplot analysis of test sites and trait relations of soybean in Ontario. Crop Sci. 42:11-20.
- ZerihunTaddese. 2014. 4th West and Central Africa science and week.11th general assembly. Nigeria.
- Zobel R.W., Wright M.J and Gauch H.G. 1988. Statistical analysis of a yield trial. Agronomy Journal, 80:388-393.