

A Fixed Point Theorem In 2-Banach Space For Non-Expansive Mapping

Vishal Gupta (Corresponding Author)
Department of Mathematics
Maharishi Markandeshwar University
Mullana, Ambala-133001(Haryana), India
Tel. +91-94164-64210 E-mail: vishal.gmn@gmail.com

A.K.Tripathi
Department of Mathematics
Maharishi Markandeshwar University
Mullana, Ambala-133001(Haryana), India
E-mail: tripathi.adesh@gmail.com

Abstract:

Our object in this paper to discuss about fixed point theory in 2-Banach space also we established a fixed point theorem in 2- Banach space which generalized the result of many mathematician.

Key Words: Normed space, 2-normed space, 2- Banach space, Nonexpensive mappings.

1 Introduction

The concept of two banach space firstly introduced by (Gahler1964) This space was subsequently been studied by mathematician (Kirk1981) and (Kirk1983) in last years. (Badshah and Gupta2005) also proved some result in 2-Banach space.(Yadav et al 2007) prove the result in 2-Banach space for non contraction mapping. (Lal and Singh 1978)the analogue og Banach Contraction principle in 2-metric space for selfmap and in the present we prove a fixed point theorem in 2-Banach Spaces by taking nanexpansive mapping.

2 Preliminaries

2.1 Definition:

Let X be a real linear space and $\|.,.\|$ be a nongative real valued function defined on X satisfying the following condition:

- (i) $\|x, y\| = 0$ iff x and y are linearly dependent.
- (ii) $\|x, y\| = \|y, x\|$ for all x,y \in X.
- (iii) $\|x, ay\| = |a|\|x, y\|$, a being real, for all x,y \in X.
- (iv) $\|x, y + z\| = \|x, y\| + \|y, z\|$ for all x,y,z \in X.

then $\|\cdot, \cdot\|$ is called a 2-norm and the pair $(X, \|\cdot, \cdot\|)$ is called a linear 2-normed space.

So a 2-norm $\|x, y\|$ always satisfies $\|x, y + ax\| = \|x, y\|$ for all $x, y \in X$ and all scalars a .

2.2 Definition:

A Sequence $\{x_n\}$ in a 2-normed space $(X, \|\cdot, \cdot\|)$ is said to be a Cauchy sequence if $\lim_{m, n \rightarrow \infty} \|x_m - x_n, a\| = 0$ for all a in X .

2.3 Definition:

A Sequence $\{x_n\}$ in a 2-normed space $(X, \|\cdot, \cdot\|)$ is said to be convergent if there is a point x in X such that $\lim_{n \rightarrow \infty} \|x_n - x, y\| = 0$ for all y in X . If x_n converges to x , we write $x_n \rightarrow x$ as $n \rightarrow \infty$.

2.4 Definition:

A linear 2-normed space is said to be complete if every Cauchy sequence is convergent to an element of X . A complete 2-normed space X is called 2-Banach spaces.

2.5 Definition:

Let X be a 2-Banach space and T be a self mapping of X . T is said to be continuous at x if for every sequence $\{x_n\}$ in X , $\{x_n\} \rightarrow x$ as $n \rightarrow \infty$ implies $\{T(x_n)\} \rightarrow T(x)$ as $n \rightarrow \infty$.

2.6 Definition:

A function $f: R \rightarrow R$ is said to be upper semi continuous at a point $x \in R$ if given $\epsilon > 0$ there exist a neighbourhood N of x_0 in which $f(x) < f(x_0) + \epsilon$ for all $x \in N$.

2.7 Definition:

Let X be a 2-Banach space and C be non empty bounded closed and convex subset of X . A mapping $T: C \rightarrow X$ is said to be nonexpansive if

$$\|T(x) - T(y), a\| \leq \|x - y, a\| \text{ where } x, y \in C$$

3 Main Result

3.1 Theorem

Let F and G be two non expansive mapping of a 2-Banach space X into itself. F and G satisfy the following condition

(1)

$FG = G = I$ where I is identity map.

(2)

$$\|F(x) - G(y), a\| \leq \alpha \|x - F(x), a\| + \beta \|y - G(y), a\| + \gamma \|x - G(y), a\| + \delta \|y - F(x), a\| + \eta \|x - y, a\|$$

where $\alpha, \beta, \gamma, \delta, \eta, \geq 0 \quad \forall \quad x, y \in X$ where $2\alpha + 2\beta + 3\delta + \gamma + \eta \leq 2$ then F and G have common fixed point.

Proof:

Taking $y = \frac{1}{2} \|(F + I)x\|, z = G(y)u = 2y - z$, then

$$\|z - x, a\| = \|G(y) - FG(x), a\|$$

Now using (1) and (2) we get

$$\begin{aligned} \|z - x, a\| &= \|G(y) - G(F(x)), a\| \leq \alpha \|y - G(y), a\| + \beta \|F(x) - G(F(x)), a\| + \gamma \|y - G(F(x)), a\| + \delta \|F(x) - G(y), a\| + \eta \|y - F(x), a\| \\ &\leq \alpha \|y - G(y), a\| + \beta \|F(x) - x, a\| + \frac{1}{2} \gamma \|F(x) - x, a\| + \delta \|F(x) - y, a\| + \delta \|y - G(y), a\| + \frac{1}{2} \eta \|y - F(x), a\| \\ &\leq (\alpha + \delta) P_{y - G(y)} + (\beta + \frac{1}{2} \gamma + \frac{1}{2} \eta + \frac{1}{2} \delta) \|F(x) - x\| \end{aligned}$$

$$\text{Now } \|u - x, a\| = \|2y - z, a\| = \|G(y) - F(x), a\|$$

$$\begin{aligned} &\leq \alpha \|y - G(y), a\| + \beta \|x - F(x), a\| + \gamma \|y - F(x), a\| + \delta \|x - G(y), a\| + \eta \|y - x, a\| \\ &\leq \alpha \|y - G(y), a\| + \beta \|x - F(x), a\| + \frac{1}{2} \gamma \|x - F(x), a\| + \frac{1}{2} \delta \|x - F(x), a\| + \delta \|y - G(y), a\| + \frac{1}{2} \eta \|x - F(x), a\| \\ &\leq (\alpha + \delta) P_{y - G(y)} + (\beta + \frac{1}{2} \gamma + \frac{1}{2} \eta + \frac{1}{2} \delta) \|F(x) - x\| \end{aligned}$$

$$\|z - u, a\| \leq \|z - x, a\| + \|x - u, a\|$$

$$\leq (2\alpha + 2\delta) \|y - G(y), a\| + (2\beta + \gamma + \delta + \eta) \|x - F(x), a\|$$

$$\text{Now } \|z - u, a\| = \|G(y) - 2y - G(y), a\| = 2 \|y - G(y), a\|$$

$$\leq (2\alpha + 2\delta) \|y - G(y), a\| + (2\beta + \gamma + \delta + \eta) \|x - F(x), a\|$$

$$\Rightarrow 2(1 - \alpha - \delta) \|y - G(y), a\| \leq (2\beta + \gamma + \delta + \eta) \|x - F(x), a\|$$

$$\Rightarrow \|y - G(y), a\| \leq \frac{2\beta + \gamma + \delta + \eta}{2(1 - \alpha - \delta)} \|x - F(x), a\|$$

$$\Rightarrow \|y - G(y), a\| \leq S \|x - F(x), a\|$$

$$\text{where, } S = \frac{2\beta + \gamma + \delta + \eta}{2(1 - \alpha - \delta)} \leq 1$$

$$\text{and } 2\alpha + 2\beta + \gamma + 3\delta + \eta \leq 2$$

Let $T = \frac{1}{2}(F + I)$, then for any $x \in X$

$$\|T^2(x) - T(x), a\| = \|T(T(x)) - T(x), a\|$$

$$\Rightarrow \|T(y) - y, a\| = \frac{1}{2}\|y - F(y), a\|$$

$$\Rightarrow \frac{1}{2}\|FG(y) - F(y), a\| \leq \frac{1}{2}\|G(y) - y, a\|, \text{ because } F \text{ is nonexpensive function .}$$

So, $\|T^2(x) - T(x), a\| \leq \frac{S}{2}\|x - F(x), a\|$, by definition of S. We claim that $T^n(x)$ is a Cauchy sequence in X.

Also by completeness $T^n(x)$ converges to $T(x)$,

i.e $\lim_{n \rightarrow \infty} T^n(x) = x_0 \Rightarrow F(x_0) = x_0$ therefore x_0 is fixed point of F.

$$\text{Again } \|T^2(x) - T(x), a\| \leq \frac{S}{2}\|x - F(x), a\| = \frac{S}{2}\|FG(x) - F(x), a\| \leq \frac{S}{2}\|x - G(x), a\|$$

we can conclude that $G(x_0) = x_0$ that is x_0 is fixed point of G.

Therefore $F(x_0) = G(x_0) = x_0$, so x_0 is common fixed point of F and G.

The uniqueness part is obvious.

References

Badshah, V. H. & Gupta, O.P. (2005), Fixed point theorem in Banach and 2-Banach spaces, *Jananabha*, Vol.35, 73-78.

Gahlar, S. (1964), 2-Metreche raume and ihre topologicche structure, *Math Nadh.*, Vol.26, 115-148.

Kirk, W.A. (1983), Fixed point theorem for nonexpensive mappings, *Contemporary mathematics*, Vol.18, 121-140.

Kirk, W.A. (1981), Fixed point theorem for non expensive mappings, *Lecture notes in mathematics*, Springer-Verlag, Berlin and NewYork, Vol.886, 111-120.

Lal, S.N. & Singh, A.K. (1978), An analogous of Banach's contraction principle for 2-metric space, *Bullatin of Australian mathematical society*, Vol.18, 137-143.

Yadva, R.N., Rajput, S.S., Choudhary, S. & Bharwaj, R.K. (2007), Some Fixed point and common fixed theorem for non-contraction mapping on 2-Banach spaces, *Acta Ciencia Indica*, Vol. 33, No.3, 737-744.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

