

Effect of Partial Rootzone Drying Irrigation on Nutrient Concentration in Leaves, Photosynthesis, Photosynthesis Active Radiation and Yield of Sorghum Cultivar

*Adzemi Mat Arshad¹ and Ibrahim W²

¹Soil Science Laboratory, Food Crop Science Unit, School of Food Science and Technology, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Malaysia

²School of Agricultural Technology and Food Science, Department of Agricultural Technology, Ramat Polytechnic Maiduguri, Borno State, Nigeria.

*adzemi@umt.edu.my

Abstract

The effects of partial rootzone drying irrigation technique on nutrient concentration in leaves, photosynthesis, photosynthetic active radiation and yield of sorghum grown on Rhu Tapai and Rengam series soil were examined at Control Environment House, Faculty of Agrotechnology and Food Science, University Malaysia Terengganu. The experiments partial rootzone drying irrigation consisted of a factorial combination of irrigation regimes and soil types laid in a randomized complete block design with eight treatments. Irrigation regimes were at four levels namely: I₁₀₀, I₇₅, I₅₀ and I₂₅ and the soil types were at two levels namely: Rhu Tapai and Rengam series soil. The treatments were randomly assigned to experimental pots and replicated four times. All agronomic practices starting from planting of sorghum to harvesting were adhered to and photosynthesis, photosynthetic active radiation and yield parameters were recorded for the experiment. The result of the study shows that, sorghum performed better under partial rootzone drying technique. The results further revealed that, irrigation regimes I₁₀₀ and I₇₅ performed better in terms of photosynthesis, photosynthetic active radiation and yield parameters compared to I₅₀ and I₂₅ irrigation regimes. The study also revealed that there was no significant different between the two types of soil used for the study. The study, therefore, recommended the use of I₇₅ percent regulated deficit irrigation for optimizing sorghum yield production in semi-arid regions.

Keywords: Partial rootzone drying irrigation, nutrient concentration, photosynthesis, photosynthetic active radiation, yield, sorghum

1. Introduction

Sorghum (*sorghum bicolor* L. Moench) is the third important cereal crop grown in the United States and the fifth most important grain crop in the world after rice, maize and barley. It was originated in the region of the North-East Africa comprising Ethiopia, Sudan and East Africa (Doggett, 1988). The crop is well adapted to the range of environmental condition in semi-arid region of Africa with high variability (Doggett, 1988; Teshome *et al.* 1997; Rami *et al.* 1998). In 2010, Sorghum is a key crop in the warm low-rainfall regions of the tropics. Hence, it is adapted to a wide scope of agro-ecological conditions ranging from the high rainfall highland of Rwanda to the arid zones of Libya, where it is produced under irrigation. Sorghum is normally grown during the rainy season but on some soils it may also be sown at the beginning of the dry season using the residual soil moisture as in the northern regions of Cameroun and Nigeria. It is also highly battle to drought and salinity and has a remarkable yield potential even in trivial environments (Cosentino, 1996; Amaducci *et al.* 2004).

Water productivity (WP) is defined as crop yield per unit functional irrigation water that is the efficiency of applied irrigation water (Zhang, 2003). Partial stomatal closure and reduced leaf area occurred due to increased Abscisic acid. These are the main physiological responses to decrease transpiration in plants under PRD and enhance WP (Davies *et al.*, 2002) therefore, a higher WP (or WUE) is obtained (Morison *et al.*, 2008). Water productivity has been increased significantly by using partial rootzone drying on diverse crops (Davies *et al.*, 2002, Sepaskhah and Khajehabdollahi, 2005, Fereres and Soriano, 2007, Costa *et al.*, 2007, Geerts and Raes, 2009, Ahmadi *et al.*, 2010b). Recently in a meta-analysis Sadras (2009) confirmed that use of partial rootzone drying enhanced water productivity by 82% compared to full irrigation with no noteworthy reduction in yields. However, Liu *et al.* (2006b) indicated that partial rootzone drying was less efficient than deficit irrigation in enhancing water use efficiency. Wakrim *et al.* (2005) and Kirda *et al.* (2005) confirmed that partial rootzone drying resulted in lower water use efficiency than deficit irrigation in beans and maize respectively. Nevertheless more optimistic effect on fruit quality was occurred in partial rootzone drying than in deficit irrigation (Kang and Zhang, 2004, Kirda *et al.*, 2004, Leib *et al.*, 2006). De la Hera *et al.* (2007) and Ahmadi *et al.* (2010b) indicated that to investigate the effectiveness of partial rootzone drying compared to deficit irrigation it is necessary to

investigate hormonal changes resulted by long-term partial rootzone drying on reproductive development whether the chemical signaling in partial rootzone drying is different from deficit irrigation, the differences in the pattern of soil water uptake, root growth, and how the water redistribution from roots can influence chemical signaling in dry roots, and the duration and best timing for application of partial rootzone drying according to crop, soil, and site specifications. Sepaskhah and Ghasemi (2008) also reported findings from their study in partial rootzone drying conducted at Iran in semi- arid region resulted in an average of 28% reduction in sorghum grain yield with related reduction in applied water at customized 15-day irrigation intervals. Studied on the effects of every-other furrow and every-furrow irrigations on grain sorghum yield and water productivity at various irrigation intervals of 10, 15 and 20 days indicated that every-other furrow irrigation at 10 day intervals of every-other furrow abridged the applied water by 11% with no yield reduction compared every-furrow irrigation at 15day intervals.

To protract the growing world population, agricultural production will need to increase (Howell, 2001) yet the fraction of fresh water currently available for agriculture 72% is decreasing (Cai *et al*, 2003). Hence sustainable methods to increase crop water productivity are gaining in arid and semi-arid regions. Irrigated agriculture is the primary user of diverted water globally reaching a proportion that exceeds 70-80% of the total in the arid and semi-arid zones. It is therefore not surprising that irrigated agriculture is perceived in those areas as the primary source of water especially in the surfacing drought situations. Currently irrigated agriculture is stiff between two perceptions that are at variance, some perceive that agriculture is highly inefficient by growing water guzzling crops (Postel *et al*, 1996) while others emphasized that irrigation is essential for the production of sufficient food in the future, given the anticipated increases in food demand due to world population growth and changes in diets (Dyson, 1999). Globally, food production from irrigation represents >40% of the total and uses only 17% of the land area devoted to food production (Feres and Connor, 2004). Nevertheless, irrigated agriculture is still practiced in many areas in the world with complete disregard to basic principles of resources conservation and sustainability. Therefore irrigation water management in an era of water scarcity will have to be carried out most efficiently, aiming at saving water and at maximising the productivity. Deficit irrigation has widely been reported as a priceless strategy for dry regions (English, 1990, Feres and Soriano, 2007) where water is the limiting factor in crop cultivation. The main objective of the study was to evaluate the effect of partial rootzone drying irrigation on the nutrient concentration in leaves, photosynthesis, photosynthetic active radiation and yield of sorghum cultivar.

2. Materials and Methods

Experiment was conducted at Control Environment House, Faculty of Agrotechnology and Food Science, Universiti Malaysia Terengganu with Latitude and Longitude 5^o.20'N 103^o 5'E. The Altitude is 32m. The climate of the area is tropical rain-forest with a mean annual rainfall of 2911mm. (114.6 in). The average temperature in Terengganu is 26.7^oC (min 22^oC, max 32^oC) while the mean relative humidity for an average year is recorded as 71.7% and on a monthly basis it ranges from 68% in May and June to 79% in December. Sorghum (*Sorghum bicolor* L. Moench) cultivar Samsorg-KSV8 from Nigeria was used in this research. The plants were planted on Rengam series soil (Ultisol) and Rhu Tapai series soil (Sandy soil) or Bris (Beach ridges interspersed with swales soil).

Various treatments comprising of different regimes of irrigation namely: (i) 100% RDI (I₁₀₀) (ii) 75% RDI (I₇₅) (iii) 50% RDI (I₅₀) and (iv) 25% RDI (I₂₅) and one type of cultivar: SAMSORG14-KSV8 and two types of soil: Rengam (Ultisol) and Rhu Tapai (Sandy soil) respectively. All treatments were laid out by following a randomized complete block design (RCBD) with four replications. A total of thirty two polythene bags were used. The total area of experimental field is 185.81m² and four lateral line pipes, each were connected to the pipe with stoppers attached at each joints of laterals connecting to main pipes. Emitters or drippers were attached according to plant spacing of 75 by 50cm. Regulated deficit irrigation treatments were arranged at the front while partial rootzone drying irrigation followed behind. On each drippers were attached with clippers for regulating the irrigate

Soil samples were taken from the experimental field. The samples were used to determine physical properties of the soil. The properties include soil texture, bulk density, field capacity, permanent wilting point and available water (Table 1).

The fertilizer nutrients recommended for sorghum are 64kg N, 30kg of P₂O₅ and 30kg K₂O per hectare was applied. Single super-phosphate and muriate of potash were applied during polythene bags preparation and including poultry fertilizer. While urea as source of nitrogen was applied in split applications. 32kgN was applied three weeks after planting and the remaining half was applied six weeks after planting at 15cm from the plants. Weeding was carried out manually throughout the growing period to reduce competition for space, water,

light and nutrients between crops. For determination of N, P and K in plant tissues of third blade below ear of sorghum leaves, wet ashing was employed by digesting tissue samples in concentrated H₂SO₄ and H₂O₂ (Thomas et. al., 1967). The concentration of N, P and K were determined by an auto-analyzer. Dry ashing was employed for determination of Ca and Mg (Chapman and Pratt, 1961) in third blade ear of sorghum leaves. The concentrations were determined using the atomic absorption spectrophotometer.

Table 1: Physio-chemical properties of Rhu Tapai and Rengam Soil Series.

Soil properties	Rhu Tapai	Rengam
Particle size distribution		
Silt (%)	2.52	3.07
Sand (%)	67.35	30.28
Clay (%)	30.13	66.65
Texture	Sandy	Clay
Organic matter (%)	0.99	1.62
pH (1:1 suspension)	4.60	4.80
Bulk density (g/cm ³)	1.27	1.31
CEC (cmol (+) kg ⁻¹ soil)	9.53	7.14
Total nitrogen (%)	0.09	0.15
Exchangeable bases (cmol (+) kg ⁻¹ soil)		
Ca	0.20	0.17
Mg	0.02	0.10
K	0.01	0.10
% of water base on weight		
0.33 bar	6.50	23.50
1.0 bar	4.00	30.05

Data collection started after transplanting, physiological and yield parameters were recorded during the crop growth and development. Total yield per hectare were equally measured. Photosynthetic rate, photosynthesis also were measured. All data collected were analyzed using SAS statistical program (SAS Inst. 1999). Analysis of variance (ANOVA) test was conducted and significant differences among the treatments were determined using the Duncan New Multiple Range Test (DNMRT) at p≤0.05

3. Results and Discussion

3.1 Effect of Partial Rootzone Drying on Nutrient Concentration in Leaves

The result shows that there are no significant different in nutrient concentration of N, P, K, Ca and Mg in leaves of sorghum of different treatments. There is no interaction between the soils used in the experiment (Table 2). The N, P, K, Ca and Mg concentration in the leaves of sorghum are at adequate level for sorghum growth and yield production (Reuter and Robinson, 1986).

Table 2: Effect of Partial Rootzone Drying on Nutrient Concentration in Sorghum Leaves

Treatments	N	P	K	Ca	Mg
	(%)				
Irrigation					
I ₁₀₀	2.56a	0.44a	3.10a	0.35a	0.23a
I ₇₅	2.56a	0.44a	3.41a	0.33a	0.21a
I ₅₀	2.52a	0.56a	2.99a	0.34a	0.19a
I ₂₅	2.46a	0.47a	3.06a	0.33a	0.20a
Rhu Tapai Soil Series	2.55a	0.45a	3.10a	0.35a	0.21a
Rengam Soil Series	2.55a	0.46a	3.11a	0.35a	0.22a

Means followed by the same letter are not significantly different at p≤0.05 (DNMRT)

3.2 Effect of Partial Rootzone Drying on Photosynthesis

The result in Figure 1 revealed that at five leaves stage there was significant different In the partial rootzone drying irrigation regimes applied. The I₁₀₀% irrigation regime recorded highest numerical values when compared to the three other irrigation regimes (I₇₅%, I₅₀% and I₂₅%). The result also indicated that I₁₀₀% and I₇₅% partial rootzone drying irrigation regimes were not significantly different in all the growth stages except at five leaf stage as showed in Figure 2, 3 and 4 respectively.




Figure 1: Effect of Partial Rootzone Drying on Photosynthesis at Five Leaf Stage




Figure 2: Effect of Partial Rootzone Drying on Photosynthesis at Jointed Stage




Figure 3: Effect of Partial Rootzone Drying on Photosynthesis at Flowering Stage.




Figure 4: Effect of Partial Rootzone Drying on Photosynthesis at Dough Stage

3.3 Effect of Partial Rootzone Drying on Photosynthetic Active Radiation

All the effect of partial rootzone drying irrigation regimes on photosynthetically active radiation at various growth stages resulted in significant differences. The I₁₀₀% partial rootzone drying irrigation regime was significantly at par when compared to the other three regimes (I₇₅%, I₅₀% and I₂₅%). However, the other three partial rootzone drying irrigation regimes were not significantly different as indicated in Figure 5 and

furthermore Figure 6, 7 and 8 as showed that the lowest photosynthetically active radiation values were recorded under the I₂₅% partial rootzone drying irrigation regime.




Figure 5: Effect of Partial Rootzone Drying on Photosynthetic Active Radiation at Five Leaf Stage.




Figure 6: Effect of Partial Rootzone Drying on Photosynthetic Active Radiation at Jointed Stage




Figure 7: Effect of Partial Rootzone Drying on Photosynthetic Active Radiation at Flowering Stage.




Figure 8: Effect of Partial Rootzone Drying on Photosynthetic ally Active Radiation at Dough Stage.

3.4 Effect of Partial Rootzone Drying on Yield of Sorghum

Table 3 revealed that grain yield were higher in one hundred percent irrigation regime ($I_{100\%}$). The result further revealed that seventy five percent irrigation regime ($I_{75\%}$) was numerically higher than the other two irrigation regimes as shown in Table 3. However, partial rootzone drying irrigation resulting in also high yield in seventy five percent irrigation regime ($I_{75\%}$) and fifty percent irrigation regime ($I_{50\%}$). This result is benefit for sorghum. These results support the reputed drought tolerance of sorghum (Krieg and Lascano, 1990). This result is in conformity with findings of the followings, the different experiment results in partial rootzone drying have

shown that irrigation water may be reduced by approximately 30% to 50% in partial rootzone drying with no significant yield reduction (Kang and Zhang, 2004; Leib *et al*, 2006; Guang-cheng, *et al*, 2008; Ahmadi, 2009).

Table 3: Effect of Partial Rootzone Drying on Yield of Sorghum

Treatment	Yield (Kg/ha)
Irrigation	
I ₁₀₀	8037.1 ^a
I ₇₅	6939.4 ^b
I ₅₀	5613.3 ^c
I ₂₅	0.0 ^d
Rhu Tapai Soil Series	5254.4 ^a
Rengam Soil Series	5040.3 ^a

Means followed by the same letter are not significantly different at $p \leq 0.05$ (DNMRT)

Conclusion

The result of the study shows that sorghum performed better under partial rootzone drying irrigation technique. Irrigation regimes I₁₀₀ and I₇₅ performed better in terms of photosynthesis, photosynthetic active radiation and yield parameters compared to I₅₀ and I₂₅ irrigation regimes. The study also revealed that there was no significant difference between the two types of soil used for the study. The study, therefore, recommended the use of I₇₅ percent regulated deficit irrigation for optimizing sorghum yield production in semi-arid regions.

Acknowledgements

The authors would like to thank Universiti Malaysia Terengganu for giving permission to publish this paper

References

- Ahmadi, S.H. (2009). Agronomic and physiological studies of partial root-zone drying and deficit irrigation on potato in different soil textures. Published Ph.D. Thesis, Department of Basic Sciences and Environment, Faculty of Life Sciences University of Copenhagen, Denmark, 77p.
- Ahmadi, S.H., Andersen, M.N., Plauborg, F., Poulsen, R.T., Jensen, C.R., Sepaskhah, A.R., Hansen, S. (2010b). Effects of irrigation strategies and soils on field grown potatoes: Yield and water productivity. *Agri. Water Management*. DOI 10.1016/j.agwat.2010.07.007
- Amaducci, S. Monti, A. and Venturi, G. (2004). Non-structural carbohydrates and fibre components in sweet and fibre sorghum as affected by low and normal input techniques. *Ind. Crops Products*, 20: 237-248.
- Cai, X. Rosegrant, M. and Ringler, C. (2003). Physical and economic efficiency of water use in the river basin: Implications for water management. *Water Resources Res.* 39(1): 1013.
- Chapman, H.D. and Pratt, P. F. (1961). *Methods of analysis for soils, plants and waters*. Univ. of California, USA. 309pp
- Cosentino, S.L. (1996). Crop physiology of sweet sorghum. In: *Proceedings of First European Seminar on Sorghum for Energy and Industry*, Toulouse, pp. 30-41.
- Costa, J.M., Ortuno, M.F. Chaves, M.M. (2007). Deficit irrigation as a strategy to save water: physiology and potential application to horticulture. *Jour. of Integrative Plant Biology*, 49: 1421-1434.
- Davies, W.J., Hartung, W. (2004). Has extrapolation from biochemistry to crop functioning worked to sustain plant production under water scarcity? *Proceeding of the 4th International Crop Science Congress, Brisbane, Australia*, published on CDROM, Published on CDROM, <http://www.cropscience.org.au/icsc2004/>.
- Davies, W.J., Wilkinson, S., Loveys, B. R. (2002). Stomatal control by chemical signaling and the exploitation of this mechanism to increase water use efficiency in agriculture. *New phytologist*, 153: 449-460.
- De la Hera, M.L. Romero, P. Gomez-Plaza, E., Martinez, A. (2007). Is partial root-zone drying an effective irrigation technique to improve water use efficiency and fruit quality in field-grown wine grapes under semi arid conditions? *Agri. Water Management*, 87: 261-274.
- Doggett, H. (1988). *Sorghum*. Tropical Agriculture Series, 2nd ed. CTA, Wageningen, The Netherlands.

- Dyson, J. (1999). The Dyson Effect: Carbon "Off" Forestry and the privatization of the Atmosphere. Pub, The corner House, Sturminster Newton Dorset DT10 1YJ UK.
- English, M. (1990). Deficit irrigation, Analytical framework. Journal of Irrigation and Drainage Engineering 116: 399-412.
- English, M. J., Musick, J. T., and Murty, v. V. N. (1990). Deficit irrigation. Pp. 631-663. In G.J. Hoffman, T.A. Howell and K.H. Solomon (Eds) Management of farm irrigation systems. ASAEMono., Am. SocAgric.and Biol. Engr., St. Joseph, M L.
- Fereres, E. And Connor, D.J. (2004). Sustainable water management in agriculture. In: challenges of the New Water Policies for the XXI Century. (eds)..
- Fereres, E. And Evans, R .G. (2006). Irrigation of fruit trees and vines: an introduction. Irrigation Science 24, 55-57.
- Fereres, E. and Soriano, M.A. (2007). Deficit irrigation for reducing agricultural water use. J. Exp. Bot. 58: 147-159.
- Fereres E, Goldhamer D. A. (1990). Deciduous fruit and nut trees. In: Stewart BA, Nielsen DR, eds. Irrigation of agricultural crops, Agronomy 30. Madison, WI: ASA, CSSA, SSSA, 987-1017
- Geerts, S. and Raes, D. (2009). Deficit irrigation as an on-farm strategy to maximize crop water productivity in dry areas. Agri. Water Management, 96: 1275-1284.
- Guang-Cheng, S., Zhan-Yua, Z., Nac, L., Shuang-Ena, Y., Weng-Ganga, X. (2008). Comparative effects of deficitirrigation (DI) and partial rootzone drying (PRD) on soil water distribution, water use, growth and yield ingreenhouse grown hot pepper. Scientia Horti., 119: 11-16.
- Howell, T.A, 2001. Enhancing water use efficiency in irrigated agriculture. Agronomy Journal 93, 281-289.
- Kang, S. and Zhang, J. (2004). Controlled alternate root zone Irrigation. It's Physiological Consequences Impact on Water use efficiency. J. Experimental Botany. 55 (407): 2437-2446.
- Kang, S.Z., Zhang, J.H. (2004). Controlled alternate partial root-zone irrigation: its physiological consequences and impact on water use efficiency. Jour. of Experimental Botany, 55: 2437-2446.
- Kirda, C., Cetin, M., Dasgan, Y., Topcu, S., Kaman, H., Ekici, B., Dericci, M.R., Ozguven, A.I. (2004). Yield response of greenhouse grown tomato to partial root drying and conventional deficit irrigation. Agri. Water Management, 69: 191-201.
- Kirda, C., Topcu, S., Kaman, H., Ulger, A.C., Yazici, A., Cetin, M., Dericci, M.R. (2005). Grain yield response and N-fertiliser recovery of maize under deficit irrigation. Field Crops Research, 93: 132-141.
- Krieg, D. R and Lascano, R. J. (1990). Sorghum in irrigation of agricultural crops. Agron.monogr. 30. Stewart, E. Aand Nielson, D. R (eds.). ASA,CSSA, Madison, WI
- Leib, B.G., Caspari, H.W., Redulla, C.A., Andrews, P.K., Jabro, J. (2006). Partial rootzone drying and deficit irrigation of 'Fuji' apples in a semi-arid climate. Irrigation Science, 24: 85-99.
- Liu, F., Shahnazari, A., Andersen, M.N., Jacobsen, S.E., Jensen, C.R. (2006b). Effects of deficit irrigation (DI) and partial root drying (PRD) on gas exchange, biomass partitioning, and water use efficiency in potato. Scientia Horticulturae, 109: 113-117.
- Morison, J.I.L., Baker, N.R., Mullineaux, P.M., Davies, W.J. (2008). Improving water use in crop production.Philosophical Transactions of the Royal Society (London) B, 363: 639-658.
- Postel, S.L Daily, G.C .Ehrlich, P.R. (1996). Human appropriation of renewable Freshwater. Science 271:785-788.
- Rami, J.F., Dufour, P., Trouchen, G., Fliedel, G., Mestres, C., Davries, F. ,Blanchand, P. and Haman, P. (1998). Quantitative trait loci for grain quality, productivity, morphological and agronomical traits in Sorghum [*Sorghum bicolor* L. Moench].Theories of Applied Genetics 97:605-616.
- Reuter, D. J and Robinson, J. B. (1986). Plant analysis and interpretation manual. Inkata Press Melbourne, Aust. 219pp.
- Sadras, V.O. (2009). Does partial root-zone drying improve irrigation water productivity in the field? A meta-analysis. Irrigation Science 27: 183-190.
- SAS Institute. (1999). SAS user's guide: Statistics. SAS Inst., Cary, NC.
- Sepaskhah, A.R., Ghasemi, M.M. (2008). Every-other furrow irrigation with different irrigation intervals for Sorghum. Pakistan Journal of Biological Science, 11: 9. 1234-1239.
- Sepaskhah, A.R., Khajehabdollahi, M.H. (2005). Alternate furrow irrigation with different irrigation intervals for maize (*Zea mays* L.). Plant Production Science, 8: 592-600.
- Teshome, A., Baum, B.R., Fahrig, L., Torrance, J.K., Arnason, T.J. and Lambert, J.D. (1997). Sorghum [*Sorghum bicolor* (L) Moench] Landrace variation and classification in north shewa and south welo, Ethiopia. Eurphytica 97: 255-263.
- Thomos, R.L., Sheard, W.W. and Moyer, J.R. (1967). Comparison of conventional and automated procedures of N, P and K analyses of plant materials using a single digestion. Agron. Jour. 59:240-243
- Wakrim, R., Wahbi, S., Tah, H., Aganchich, B., Serraj, R. (2005). Comparative effects of partial root drying (PRD) and regulated deficit irrigation (RDI) on water relations and water use efficiency in common bean (*Phaseolus vulgaris* L.). Agri., Ecosystems & Environment, 106: 275-287.
- Zhang, H. (2003). Improving water productivity through deficit irrigation: Examples from Syria, the north China Plain and Oregon, USA. In: Water Productivity in Agriculture: Limits and Opportunities for Improvement. Kijne, J.W., Barker, R., and Molden, D. (eds). CABI publishing, 332p