

Impact of Climate Change on Livestock Health: A Review

Abdela Nejash

School of veterinary medicine, College of agriculture and veterinary medicine, Jimma University, Jimma
Ethiopia

ABSTRACT

This review work was conducted to explore the likely impacts of climate change on livestock health. Currently, the world is facing a number of challenges, of which climate change is a priority area. Climate change affects livestock health through several pathways involving both direct and indirect effect. The direct effects being most likely pronounced for diseases that are vector- borne, soil associated, water or flood associated, rodent associated, or air temperature/humidity associated and sensitive to climate. Furthermore, Climate change influences the emergence and proliferation of disease hosts or vectors and pathogens and their breeding, development and disease transmission. Consequently, it affects distributions and host–parasite relationships and its assemblages to new areas. Climate factors also influences habitat suitability, distribution, and abundance; intensity and temporal pattern of vector activity. Pathogens and parasites that are sensitive to moist or dry conditions may be affected by changes to precipitation and soil moisture. Higher temperatures resulting from climate change may increase the rate of development of certain pathogens or parasites that have one or more life cycle stages outside their animal host. This may shorten generation times and, possibly, increase the total number of generations per year, leading to higher pathogen/ parasite population sizes. Mammalian cellular immunity can be suppressed following heightened exposure to ultraviolet. In particular, there is depression of the number of T helper 1 lymphocytes, the cells involved in the immune response to intracellular pathogens. Therefore, successful adaptations may be shown as better way of coping with the negative consequences of climate change on livestock health.

Keywords: *Climate change, Livestock health, Livestock, Temperatures*

INTRODUCTIONS

Global climate change poses the threat of serious social upheaval, population displacement, economic hardships and environmental degradation (ESAP, 2009). Climate change and global warming now being accepted facts have affected all the ecosystems and will do so if left uncontrolled. Impacts on some components have gained more attention while others have been neglected. Animals belong to latter. Even among the aspects relating to impacts of climate change on animals, production related impacts have gained attention when the impacts on health in general and on infectious diseases in particular are neglected (Yatoo *et al.*, 2012). Climate change influences the emergence and proliferation of disease hosts or vectors and pathogens and their breeding, development and disease transmission. Consequently, it affects distributions and host–parasite relationships and its assemblages to new areas (ESAP, 2009)

Climate change, in particular global warming, is likely to greatly affect the health of animals, both directly and indirectly. The direct effects of climate on animal disease are likely to be most pronounced for diseases that are vector- borne, soil associated, water or flood associated, rodent associated, or air temperature/humidity associated and sensitive to climate (Grace *et al.*, 2015). Indirect impacts follow more intricate pathways and include those deriving from the attempt of animals to adapt to thermal environment or from the influence of climate on microbial populations, distribution of vector-borne diseases, and host resistance to infectious agents, feed and water shortages, or food-borne diseases (Yatoo *et al.*, 2012). Furthermore, Climatic changes can influence livestock health through a number of factors, including the range and abundance of vectors and wildlife reservoirs, the survival of pathogens in the environment (ESAP, 2009)

Higher temperatures resulting from climate change may increase the rate of development of certain pathogens or parasites that have one or more life cycle stages outside their animal host. This may shorten generation times and, possibly, increase the total number of generations per year, leading to higher pathogen/ parasite population sizes (Chauhan *et al.* 2014; Grace *et al.*, 2015). Increase in temperature results in the spatial distribution and intensity of existing pests, and diseases which in turn affect livestock productivity or may cause death of livestock in some extreme instances (Musemwa *et al.*, 2012). According to ECARD (2008), most diseases are transmitted by vectors such as ticks and flies, the development stages of which are often heavily dependent on temperature. Cattle, goats, horses and sheep are also vulnerable to an extensive range of nematode worm infections, most of which have their development stages influenced by climatic conditions more particular temperature. Understanding of the relation between climate change and livestock disease is critical for better management of animal health problems. Therefore, the objective of this paper to give an overview on the effects of climate changes on livestock health.

CLIMATE CHANGE AND ANIMAL DISEASES LINKAGE

The distribution of infectious diseases, (human, animal and plant) and the timing and intensity of disease outbreaks are often closely linked to climate. Climate change may affect livestock disease through several pathways both direct and indirect. The direct effects of climate on animal disease are likely to be most pronounced for diseases that are vector-borne, soil associated, water or flood associated, rodent associated, or air temperature/humidity associated and sensitive to climate (Grace *et al.*, 2015). These directly or indirectly effects by weather and climate may be spatial, with climate affecting distribution, *temporal* with weather affecting the timing of an outbreak, or relate to the *intensity* of an outbreak. Global climate change alters ecological construction which causes both the geographical and phenological shifts (Slenning, 2010). These shifts affect the efficiency and transmission pattern of the pathogen and increase their spectrum in the hosts (Brooks and Hoberg, 2007).

The increased spectrum of pathogens increases the disease susceptibility of the animal and thus, supports the pathogenicity of the causative agent. The livestock systems are susceptible to changes in severity and distribution of livestock diseases and parasites as potential consequences. Incidence of external parasite (43.3%) was first ranked as the problem in the warm temperate (Dhakal *et al.*, 2013). vector-borne diseases are especially sensitive to climate change. Changes in rainfall and temperature regimes may affect both the distribution and the abundance of disease vectors, as can changes in the frequency of extreme events. Arthropod vectors tend to be more active at higher temperatures; they therefore feed more regularly to sustain the increase in their metabolic functions, enhancing chances of infections being transmitted between hosts. Small changes in vector characteristics can produce substantial changes in disease (Grace *et al.*, 2015).

There is a link between climate and epidemiological conditions of disease agents. Temperature, precipitation, humidity, and other climatic factors are known to affect the reproduction, development, behavior, and population dynamics of the helminthes, arthropod vectors and the pathogen they carry. Climate change influences the emergence and proliferation of disease hosts or vectors and pathogens and their breeding, development and disease transmission (ESAP, 2009). The OIE Scientific Commission has concluded that climate changes are likely to be an important factor in determining the spread of some diseases, especially those that are vector-borne. The two most mentioned emerging and re-emerging cattle diseases in a recent OIE survey are Catarrhal fever (Bluetongue) and Rift Valley fever (OIE, 2008).

IMPACT OF CLIMATE CHANGE ON ANIMAL HEALTH

Impact of climate change on vectors

Arthropod vectors are cold-blooded (ectothermic) and thus especially sensitive to climatic factors. Temperature, precipitation, humidity, and other climatic factors influence the survival, production, development, behavior, and population dynamics of the arthropod vectors. Subsequently, climate factors influences habitat suitability, distribution, and abundance; intensity and temporal pattern of vector activity (particularly biting rates) throughout the year (ESAP, 2009)

There are several processes by which climate change might affect disease vectors. First, temperature and moisture frequently impose limits on their distribution. Often, low temperatures are limiting because of high winter mortality and a relatively slow rate of population recovery during warmer seasons. By contrast, high temperatures are limiting because they involve excessive moisture loss. Therefore, cooler regions which were previously too cold for certain vectors may begin to allow them to flourish with climate change. Warmer regions could become even warmer and yet remain permissive for vectors if there is also increased precipitation or humidity conversely, these regions may become less conducive to vectors if moisture levels remain unchanged or decrease, with concomitant increase in moisture-stress (Baylis and Githeko, 2006).

Changes in climate will influence arthropod vectors, their life cycles and life histories, resulting in changes in both vector and pathogen distribution and changes in the ability of arthropods to transmit pathogens. Therefore animals will be exposed to different parasites and/or diseases, as indicated by the predicted change in the distribution of, for example, the Tsetse fly in Africa, putting an even greater pressure on production and the survival of livestock breeds (Tabachnick, 2010).

The epidemiology of many diseases are based on transmission through vectors such as ticks, lice, mites, mosquitoes and flies, the developmental stages of which are often heavily dependent on temperature and humidity. Changes in rainfall and temperature regimes may affect both the distribution and the abundance of disease causing vectors, as can changes in the frequency of extreme events (Thornton *et al.* 2009). Research studies from India have found that meteorological parameters like temperature, humidity and rainfall explain 52 and 84% variations in the seasonality of Foot and Mouth (FMD) disease in cattle in hyper-endemic division of Andhra Pradesh and meso-endemic region of Maharashtra states, respectively (Ramarao 1988). The hot-humid weather conditions were found to aggravate the infestation of cattle ticks like, *Boophilus microplus*, *Haemaphysalis bispinosa* and *Hyalomma anatolicum* (Kumar *et al.* 2004).

The feeding frequency of arthropod vectors may also increase with rises in temperature. Many vectors

must feed twice on suitable hosts before transmission is possible - once to acquire the infection and, after the EIP, once to transmit it. For many blood-feeding arthropods, feeding frequency is determined by the time required for egg development. For example, *C. sonorensis* females feed every three days at 30 °C but only every ~14 days at 13 °C. At the warmer temperature, the vector is more likely to take the two feeds on suitable hosts that are required for successful transmission (Baylis and Githeko, 2006).

Impact of climate change on Pathogens

Higher temperatures and greater humidity generally increase the rate of development of parasites and pathogens that spend part of their life cycle outside the host. Changes to wind can affect the spread of pathogens. Flooding that follows extreme climate events provides suitable conditions for many water-borne pathogens. Drought and desiccation are inimical to most pathogens (Grace *et al.*, 2015). Increased the rate of development due higher temperatures may shorten generation times and, possibly, increase the total number of generations per year, leading to higher pathogen/ parasite population sizes. Conversely, some pathogens are sensitive to high temperatures and their survival may decrease with climate warming. Pathogens and parasites that are sensitive to moist or dry conditions may be affected by changes to precipitation, soil moisture and the frequency of loads. Changes to winds could affect the spread of certain pathogens and vectors. Some pathogens/parasites and many vectors experience significant mortality during cold winter conditions; warmer winters may increase the likelihood of successful overwintering (Harvell *et al.*, 2002).

Lengthening of the warm season may increase or decrease the number of cycles of infection possible within one year for warm- or cold-associated diseases respectively. Arthropod vectors tend to require warm weather so the infection season of arthropod-borne diseases may extend. Some pathogens/parasites and many vectors experience significant mortality during cold winter conditions; warmer winters may increase the likelihood of successful overwintering (Baylis and Githeko, 2006). Extreme weather events, for example, flooding can carry a risk of *Cryptosporidium* parasites and/or enterohaemorrhagic *Escherichia coli* emerging as diffuse pollution in a run-off from agricultural land. This poses an obvious threat to other livestock and is also a zoonotic risk to humans through contamination of water supplies. Future challenges in the control of parasitic zoonoses, including those related to climate change; deserve increasing attention alongside production-limiting disease (Polley and Thompson, 2009).

Impact of climate change on hosts

Some livestock will be exposed to new pathogens and vectors as their range increases and impacts can be severe. Climate stress (heat, inadequate food and water) can also lower host immunity (Grace *et al.*, 2015). Climate change may bring about substantial shifts in disease distribution, and outbreaks of severe disease could occur in previously unexposed animal populations (possibly with the breakdown of endemic stability) (Thornton *et al.* 2009). Endemic stability occurs when the disease is less severe in younger than older individuals, when the infection is common or endemic and when there is lifelong immunity after infection. Certain tick-borne diseases of livestock in Africa, such as anaplasmosis, babesiosis and cowdriosis, show a degree of endemic stability (Eisler *et al.* 2003).

Mammalian cellular immunity can be suppressed following heightened exposure to ultraviolet B (UV-B) radiation an expected outcome of stratospheric ozone depletion. In particular, there is depression of the number of T helper 1 lymphocytes, the cells involved in the immune response to intracellular pathogens. In terms of animal disease, such pathogens include viruses, rickettsia (such as *Cowdria* and *Anaplasma*, the causative agents of heartwater and anaplasmosis) and some bacteria, such as *Brucella*, the organism causing brucellosis (Baylis and Githeko, 2006).

CONCUSSIONS

Climate change has negative effect on livestock health in many aspects. Direct effects on animal disease are likely to be most pronounced for diseases that are vector- borne, soil associated, water or flood associated, rodent associated, or air temperature/humidity associated and sensitive to climate. Climate change can exacerbate disease in livestock, and some diseases are especially sensitive to climate change. Therefore, successful adaptations may be shown as better way of coping with the negative consequences of climate change and associated drivers of disease. Climatic changes may influence livestock health through a number of factors, including the range and abundance of vectors and wildlife reservoirs, the survival of pathogens in the environment.

REFERENCE

- Baylis, M., & Githeko, A. K. (2006). The effects of climate change on infectious diseases of animals. Report for the Foresight Project on Detection of Infectious Diseases, Department of Trade and Industry, UK Government, 35.
- Brooks, D. R. and Hoberg, E. P. 2007. How will global climate change affect parasite host assemblages. *Trends in Parasitology* 23: 571-574.

- Chauhan, D. S., & Ghosh, N. (2014). Impact of Climate Change on Livestock Production: A Review. *Journal of Animal Research*, 4(2), 223.
- Dhakal C. K., Regmi P. P., Dhakal I. P., Khanal B. Bhatta U. K, Barsila S. R. and Acharya B. 2013. Perception, Impact and Adaptation to Climate Change: An Analysis of Livestock System in Nepal. *J. Anim. Sci. Adv.*, 3(9): 462-471
- ECARD (European Commission Agriculture and Rural Development) (2008) "Climate change: the challenges for agriculture", European Commission Directorate-General for Agriculture and Rural Development
- Eisler, M.C., Torr, S.J., Coleman, P.G., Machila, N. and Morton, J.F. 2003. Integrated control of vector-borne diseases of livestock - pyrethroids: panacea or poison? *Trends in Parasitology*, 19: 341-345
- ESAP (Ethiopian Society of Animal Production), 2009. Climate change, livestock and people: Challenges, opportunities, and the way forward. Zelalem Yilma and Aynalem Haile (Eds). Proceedings of the 17th Annual conference of the Ethiopian Society of Animal Production (ESAP) held in Addis Ababa, Ethiopia, September 24 to 26, 2009. ESAP, Addis Ababa 300 pp
- Grace, D., Bett, B., Lindahl, J., & Robinson, T. (2015). Climate and Livestock Disease: assessing the vulnerability of agricultural systems to livestock pests under climate change scenarios. CCAFS Working Paper No. 116. Copenhagen, Denmark: CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS).
- Harvell, C.D., Mitchell, C.E. and Ward, J.R. 2002. Climate warming and disease risks or terrestrial and marine biota. *Sciences*, 296: 2158-2162.
- Kumar S., Prasad K.D. and Deb A. R 2004. Seasonal prevalence of different ectoparasites infecting cattle and buffaloes. *BAU. J. Res.* 16(1): 159-163
- Musemwa, L., Muchenje, V., Mushunje, A., & Zhou, L. (2012). The impact of climate change on livestock production amongst the resource-poor farmers of third world countries: a review. *Asian Journal of Agriculture and Rural Development*, 2(4), 621.
- OIE (2008). Report of the Meeting of the OIE Scientific Commission for Animal Diseases. http://www.oie.int/download/SC/2008/A_SCAD_feb2008.pdf.
- Polley, L., & Thompson, R. A. (2009). Parasite zoonoses and climate change: molecular tools for tracking shifting boundaries. *Trends in parasitology*, 25(6), 285-291.
- Ramarao D. 1988. Seasonal indices and meteorological correlates in the incidence of foot- and- mouth disease in Andhra Pradesh and Maharashtra. *Ind. J. Anim. Sci.* 58(4):432- 434.
- Slenning, B. D. 2010. Global climate change and implications for disease emergence. *Veterinary Pathology.*, 47(1):28-33
- Tabachnick, W. J. (2010). Challenges in predicting climate and environmental effects on vector- borne disease epizootics in a changing world. *The Journal of experimental biology*, 213(6), 946-954.
- Thornton, P., van de Steeg, J., Notenbaert, M.H and Herrero, M. 2009. The impacts of climate change on livestock and livestock systems in developing countries: A review of what we know and what we need to know. *Agri. Systems*, 101: 113-127.
- Yattoo, M. I., Kumar, P., Dimri, U., & Sharma, M. C. (2012). Effects of climate change on animal health and diseases. *International Journal of Livestock Research*, 2(3), 15-24.