

# Growth and Performance as affected by inclusion of *Moringa oleifera* leaf meal in Broiler chicks diet

Banjo, O.S.

Department of Agricultural Production and Management Science,  
Tai Solarin University of Education, P.M.B. 2118, Ijagun, Ogun State, Nigeria.

\*E-mail for correspondence: [banjowolepolo@yahoo.com](mailto:banjowolepolo@yahoo.com)

## Abstract

The experiment was carried out to investigate the inclusion of *Moringa oleifera* leaf as feed additive in broiler chicks. Eighty Anak 2000 Strains of two weeks old were allotted to four treatments with five replicates of four birds each in a completely randomized design. Four different diets with metabolizable energy levels ranging from 2800 to 2900kcal/kg diet were formulated and fed to the chicks for a period of four weeks. The level of inclusion of *Moringa oleifera* leaf meal ranged from 0% which served as the control, 1%, 2% and 3% in the diet. Inclusion of *Moringa oleifera* significantly ( $P < 0.05$ ) enhanced weight gain of birds at 2% level of inclusion. The inclusion of *Moringa* did not significantly ( $P < 0.05$ ) enhance feed intake and feed conversion.

**Keywords:** Broiler chicks, *Moringa oleifera*, growth, feed intake, feed conversion

## 1. Introduction

One of the practical solutions to some of the problems of poultry in the tropics is to pay attention to the areas of nutrient requirements of birds for maintenance and production and the nutrient composition of the available feed stuffs. The most logical step to take in solving the shortage and dwindling raw material supply is to direct efforts towards utilizing plants by-products and wastes for feeding poultry birds.

*Moringa oleifera* is one of the plants that can be utilized in the preparation of poultry feeds. The plant apart from being a good source of vitamins and amino acids, it has medicinal uses (Makkar and Bekker 1999; Francis *et al* , 2005). *Moringa oleifera*, otherwise regarded as a “miracle tree” has been used in the treatment of numerous diseases (Pal *et al*, 1995; Makomen *et al*, 1997; Gbasi *et al*, 2000 and Matthew *et al*, 2001) including heart disease and obesity due to its hypocholesterolemic property (Gbasi *et al*, 2001; Olugbemi *et al* 2010) also reported this quality. *Moringa oleifera* leaves have the calcium equivalent of 4 glasses of milk, 3 times the iron of spinach, 4 times the amount of vit A in carrot, and 2 times protein in milk (Loren, 2007). The leaves of *Moringa* are good source of protein, vitamins A, B and C and minerals such as calcium and iron (Dahot, 1988). The leaves of *Moringa* has high protein content which is between 20 – 33% on a dry weight basis, the protein is of high quality having significant qualities of all the essential amino acid as reported by Foidl and Paull (2008). Murro *et al* (2003) reported that the leaves contains a high level of vitamins A, B, C and calcium.

Kakengi *et al* (2003) reported that *Moringa oleifera* leaf meal was substituted for sunflower seed meal as a protein source for layers. The effects of substitution on feed intake, dry matter intake, weight, laying percentage and feed conversion ratio were investigated and they suggested that *Moringa* leaves could completely replace SSM up to 20% without detrimental effect on layers. However the crude fibre content if high can impair nutrient digestion and absorption (Aderemi, 2003; Omu 2011).

This study therefore considers the utilization of *Moringa oleifera* for improving the nutritional value of broilers and also to investigate the level of inclusion that will yield optimum performance of birds (broilers).

## 2. Materials and methods

The experiment was carried out at the Teaching and Research Farm of Tai Solarin University of Education, Ijagun, Ogun State, Nigeria. Feed ingredients were purchased from F.A feed (Nig) Limited, Ijebu – Ode, while *moringa oleifera* leaves were harvested from a farm in Ijebu – Oru, Ogun State. The *moringa oleifera* leaves were harvested and air dried under shade for 4 days and milled, after which the leaf meal was added into the diets at 0% to 3% level. A total of eighty unsexed two weeks old Anak 2000 strain of broiler chickens were allotted to four treatments with five (5) replicates of four birds each. The birds were assigned in a completely randomized design (CRD). Each

treatment group was fed one of four experimental diets containing 0, 1, 2 and 3% *Moringa Oleifera* leaf meal. The experimental birds were raised using deep litter systems, which was divided into experimental units. The records of growth rate and feed intake were taken weekly for a period of six weeks. The data collected were subjected to analysis of variance (ANOVA). Durcans's multiple range (DMR) Test was used to separate means where significant differences were observed.

### 3. Results and discussion

The result of the performance characteristics is shown in Table 3. The inclusion of *Moringa oleifera* leaf meal in the diet of the broilers significantly ( $P < 0.05$ ) enhanced their weight gain at 1% level which was significantly higher than the control.

The birds fed T2 recorded significant, ( $P < 0.05$ ) higher weight gain than T1 while those fed T3 diet recorded significantly ( $P < 0.05$ ) the highest body weight gain. The reason for the improved weight gain can be attributed to high protein content of *Moringa* leaf meal as claimed by (Danol, 1986) (Kakengi *et al* 2003) and (Olugbemi *et al.*, 2010).

The decrease in weight gain of birds fed T4 diet as compared to T2 and T3 despite the higher crude protein content may be due to higher crude fibre content which may impair nutrient digestion and absorption as claimed by (Otuma and Onu, 2008), while the reduced weight gain of broilers fed the control diet (T1) may be ascribed to low crude protein content of the diet compared to other diets.

The feed intake significantly ( $P < 0.05$ ) varied among the treatments. The reduced intake of diet T4 may be due to high crude fibre content which may invariably reduce palatability (Kakengi *et al* 2008).

The feed conversion ratio of the birds was significantly improved in all the treatments, while the diets produced no significant ( $p > 0.05$ ) impact on the protein efficiency ratio of the broiler birds.

### 4. Conclusion

The effect of *Moringa Oleifera* leaf meal used in this study was pronounced in the weight gain of the birds and it is also concluded that broilers can tolerate *Moringa oleifera* leaf meal up to 3% birds of inclusion without adverse effect on their growth.

### References

- Aderemi, F.A. (2003), "Effect of enzyme supplemented cassava siliate in cassava based diet on some visceral organs of pullet chicks". Proc. Of the 8<sup>th</sup> Annual Conference of the Animal Science Society of Nigeria Pp 57 – 59.
- Dahot, M.U. (1988), "Vitamin contents of flowers and seeds of *Moringa oleifera*". Biochemistry Pp 2122 – 2124.
- Francis, G., Makkar, H.P.S. and Becker, K. (2005), "Product from little research plants as aquaculture feed ingredients". Retrieved February 24, 2005 from <http://www.fao.org/DOCREO/ARTICLE/AGRIPPA/SSIEN.HTMLTOP>
- Foidl, N. and Paull, R. (2008), "*Moringa oleifera*". In: The Encyclopedia of Fruit and Nuti (ABI, Oxfordshire.uk Pp 509 – 512.
- Gbasi, S., Nwobodo, E. and Ofili, J.O. (2000), "Hypocholesterolemic affects of crude extract of leaf of *Moringa oleifera* lam in high fat diet fed wistar rats". *Journal of Ethonopharmacology* 69 (1): 21 – 25.
- Kakengi, A.M.V., Shen, M.N., Sarvert, S.V. and Fujihara, T. (2003), "Can *Moringa oleifera* be used as protein supplement to ruminant diet". *Asian – Australian Journal of Animal Science* 18 (1): 42 – 47.
- Makker, H.P.S. and Becker, K. (1999), "Plant toxins and detoxification methods to improve feed quality of tropical seeds review", *Asian-Australian Journal of Animal Science* (3): 467 - 480.
- Matthew, T., Matthew, Z., Taji, S.A. and Zachariah, S. (2001), "A review of viricidal Ayurvedic Herbs of India for poultry disease". *Journal of American Holistic Veterinary Medicine Association* 20 (1) 17 – 20.
- Makomen, E., Hunde, A. and Damecha, G. (1997). "Hypoglycaemic effect of *Moringa steropetala* aqueous extracts in rabbits". *Phyto-therapy Research* 11:147 – 148.

Murro, J.K, Mulikanmbele, V.R.M. and Sarwatt, S.V. (2002), “*Moringa oleifera* leaf meal can replace cotton seed cake in the concentrate mix feed with Rhodes grass (*Chloris gayana*) hay for growing sheep”. *Livestock Research for Rural Development* 15 (11).

Olugbemi, T.S., Mutayoba, S.K. and Lekule, F.P. (2010), “Effect of *Moringa oleifera* inclusion in cassava based diets fed to broiler chickens”. *International Journal of Poultry Science*, 9 (4):363 – 367.

Onu, P.N and Otuma, M.O. (2008), “Utilization of heat-treated sheep dropping in the diets of broiler finisher chicks”. *International journal of Poultry science*, 8 (10): 995 – 998.

Pal, S.K., Mukherjee, P.K. and Saha, B.P. (1995). “Studies on the anti-nuclear activity of *Moringa oleifera* leaf meal extract on gastric ulcer models in rat”. *Phytotherapy Research* 9:463 - 465.

**Table 1: Composition of experimental diets (%)**

Ingredients	Treatments			
	T1	T2	T3	T4
Maize	40	40	40	40
Soya meal	25	25	24	23
Groundnut cake	5	5	5	5
<i>Moringa oleifera</i>	-	1.0	2.0	3.0
Wheat offal	5	5	5	5
Orzel shell	5	5	5	5
Bone meal	2.0	2.0	2.0	2.0
Corn bran	14	13	13	13
Fish meal	2.5	2.5	2.5	2.5
L. lysine	0.25	0.25	0.25	0.25
Di. Methionine	0.35	0.35	0.35	0.35
Salt	0.4	0.4	0.4	0.4
Broiler premix	0.25	0.25	0.25	0.25
Mycofix	0.25	0.25	0.25	0.25
Total	100	100	100	100

**Table 2: Calculated nutrient**

	Diet 1	Diet 2	Diet 3	Diet 4
	Control Moringa	Moringa oleifera (1%)	Moringa oleifera (2%)	Moringa oleifera (3%)
Me (kal/kg)	2680.2	2506	2506.562	2506.563
Crude protein (%)	18.61	18.66	18.82	18.85
Crude fibre (%)	5.14	5.16	5.17	5.20
Calcium (%)	3.50	3.43	3.45	3.47
Phosphorus (%)	0.05	0.82	0.83	0.85
Methionine (%)	0.40	0.50	0.50	0.50
Lyshe (%)	0.75	0.80	0.80	0.80

**Table 3: Performance of broiler chicks fed *Moringa* leaf meal**

	T1	T2	T3	T4	S.E
Mean initial body weight (g)	145.25	140.60	150.20	148.50	1.36
Mean final body weight (g)	110.23 <sup>c</sup>	1415.124 <sup>ab</sup>	1521.56 <sup>a</sup>	1255.76 <sup>b</sup>	48.32
Mean body weight gain (g)	954.98 <sup>c</sup>	1274.52 <sup>a</sup>	1371.36 <sup>a</sup>	1107.46 <sup>b</sup>	46.55
Mean daily weight gain (g)	51.63 <sup>c</sup>	62.45 <sup>a</sup>	65.14 <sup>a</sup>	53.92 <sup>b1.92</sup>	
Mean total feed intake	2415.20 <sup>c</sup>	2822.40 <sup>a</sup>	28.73 <sup>a</sup>	22.046 <sup>b</sup>	45.73
Feed conversion ration	2.26 <sup>b</sup>	1.90 <sup>a</sup>	1.85 <sup>a</sup>	1.86 <sup>a</sup>	0.61
Protein efficiency ratio	2.08	2.58	2.46	2.57	0.14

Means having the same letter(s) in a column are not significantly ( $p < 0.05$ ) different

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

## CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

