

Characterization of Past Seasonal Climate Variability in Oromia Special Zone, North Eastern Ethiopia

Yimer Mohammed¹ Dr.Gashaw Bimrewu² Dr. Lisanwork Nigatu³

1.East Amhara Meteorology Service Center, Kombolcha, Ethiopia, P.O. Box 94

2.School of College of Agriculture and Environmental Sciences, BahrDar University, BahrDar, Ethiopia

3.School of Natural Resource Mang & Environmental Sciences ,Haramaya University. P.O.Box 138, Dire Dawa, Ethiopia

Abstract

This study was undertaken in oromia special zone of Amhara National Regional State to analyze past climate variability. Gridded climate data were obtained from National Meteorology Agency of Ethiopia. Standard stastical descriptors and statistical software like Instat V3.37, MAKESENS, XLSTAT 2014 and Arc GIS 10.1 were employed for the analysis and interpretation for results. The results indicated that the OSZ experienced moderate rainfall variability in kiremt (JJAS) season and very high variability in belg (FMAM) season . Increasing trend in kiremt rainfall while decreasing trend in belg rainfall. Except belg minimum temperature, also temperature trend increased. The median of the start of belg season did not between March and May. The risk of getting 5,7,10 and 15 days of dry spells based on the fitted first order Markov chain probability values indicate in belg season the probability of dry spell occurrence of 10 days' length was 60%. In kiremt season, the dry spell risk of being greater than10 day consecutive dry spells was less than 50%, and it is expected to occurred beteewen 178 DOY (June 26) to 262 DOY (Sept 18). District level of upcoming season climate information in terms of climate variability and encourage farmers to benefit from these services, apply adaptation and mitigation strategy are important for reduce the challenges of cropping practices in the study area.

Keywords:- Belg, Kiremt, Oromia, special Variavility

DOI: 10.7176/JBAH/10-1-06

Publication date: January 31st 2020

1. INTRODUCTION

Ethiopia's climate is prone to both extended rainfall deficits and excesses (Korecha, 2013). Studies in Ethiopia have shown that the causes for rainfall variability are erratic nature of rainfall distribution and late onset and early offset contribute to decline in crop yields with reasonable amount in almost all parts of the country (Godswill *et al.*, 2007). Assessing the characteristics of temperature and rainfall for a location is useful for choosing the most appropriate enterprises, and the most productive plant cultivars (Mavi and Tupper, 2004). In recent years, a case study made in parts of Ethiopia examined the impact of rainfall variability on the Ethiopia economy, and found that rainfall variability in the country led to a production deficit of 20%, and increase in poverty rates by 25% which costed by the economy over one-third of its growth potential (Hagos *et al.*, 2009). There are only a few studies on the effects of climate variability on crop production in Ethiopia (Bewket, 2009) either at national or regional scales which mask local scale variability like Oromia special zone.

Having knowledge on sequences of rainfall variability, events can assist acquiring specific information for agricultural planning (Mandal *et al.*, 2013). Within variable seasonal rainfall patterns, understanding the events of the occurrence of rain features like; onset and end date of rainy season, dry spells are crucial to decrease the adverse effects and exploit opportunities (Yemenu and Chemedda, 2013).The study of past inter seasonal rainfall variability in terms of onset, end date and length of rainy season, number of rainy days, length of dry spell within the growing period and its trend is important for agricultural purposes in the dry land area than annual and seasonal totals (Hadgu *et al.*, 2013).

Rainfall during a year occurs in different seasons, three seasons are known in Ethiopia, namely *bega*, *belg* and *kiremt* (NMA, 2007). So the main objective of this paper is to anylas two rainy season climate varavility of the special zone.

2 MATERIALS AND METHODS

2.1 DESCRIPTION OF THE STUDY AREA

The study area covers Oromia Special Zone found in North Eastern of Ethiopia. Geographically, it is located between 10^o 00' N to 11^o 30' N latitude and 39^o 40' E to 40^o20' E longitude (Figure 1). Kемisse is the administrative center of the Zone which is located at 326 km north east of Addis Ababa, the capital city of Ethiopia.




Figure 1. Map of the study area with selected stations

2.2 DATA COLLECTION

The choice of the four districts were based on representativeness of geographical settings in the study area (fig-1). The Special Zone has lack of long year observed climate data. As a result, gridded daily rainfall and dekadal temperature data of four weather station points for the time period 1986 to 2015 were collected from National Meteorology Agency of Ethiopia used for these analyses.

2.3 DATA ANALYSIS

2.3.1 VARIABILITY ANALYSIS USING STATISTICAL PARAMETERS

Analysis of two rainy season rainfall and temperature and involved characterizing long-term mean values, and calculation of indices of variability, and trends at seasonal and annual time scales.

Heterogeneity of monthly rainfall amount were investigated using the precipitation Concentration Index (PCI).

The seasonal PCI values were calculated as given by Oliver (1980) as follows:

PCI on seasonal scale calculated as:

$$PCI_{seasonal} = \frac{100}{3} * \frac{\sum_{i=1}^4 p_i^2}{\sum_{i=1}^4 (p_i)^2} \quad (Eq 1)$$

Where, P_i is the rainfall amount of the i^{th} month; and Σ = summation over the 12 months incase of annual time scale and four months incase of seasonal time scale.

According to this classification, Oliver (1980) suggested that PCI values less than 10 represents a uniform precipitation distribution (i.e., low precipitation concentration); PCI values from 11 to 15 denote a moderate precipitation concentration; values from 16 to 20 denote irregular distribution, and values above 20 represent a strong irregularity (i.e., high precipitation concentration) of precipitation distribution.

Coefficient of variation, standard deviation, and mean were used to analyze the variation in explanatory variables (rainfall characteristics and temperature).

Scientifically, using the following formula.

$$CV = \frac{SD}{\bar{x}} * 100 \quad (Eq 2)$$

Where, CV is coefficient of variation, SD is the standard deviation and \bar{x} mean of observed data. According to

Hare (1983), CV (%) values are classified as follows: < 20% as less variable, 20- 30% as moderately variable, and > 30% as highly variable.

Standard deviation was calculated from the observed data, mean and number of the observations as follows:

$$SD = \sqrt{\frac{\sum_{i=1}^n \left(x_i - \bar{x} \right)^2}{n}} \quad (Eq 3)$$

Where, SD is the standard deviation, x is an observed value, \bar{x} is mean of observation data and n is number of observations (years). Using the classification of Reddy (1990), the stability of rainfall is examined as follows:

when standard deviation <10 as very high stabilities, 10-20 as high stability, and 20-40 as moderate stability and >40 as less stability. In addition standard statistical tool and software like instat V3.37, XLSTAT 2014 and Microsoft Excel 2016 were used for the purpose of any analysis.

2.3.2 SEASONAL RAINFALL ANOMALY

As described by Agnew and Chappel (1999) the standardized rainfall anomaly (Z_{ij}) were used to characterize the annual and seasonal drought frequency and intensity, and inter seasonal fluctuations of rainfall and crop yield as given below.

$$Z_{ij} = \frac{x - \bar{x}}{SD} \quad (\text{Eq 4})$$

Where, Z_{ij} is normalized rainfall total for station i during a year (or season) j ; X is an observed annual and seasonal rainfall value, \bar{x} is mean and SD is the standard deviation. This statistic enables us to determine the dry

(-ve values) and wet (+ve values) years in the records. The drought severity classes are extreme drought ($Z_{ij} < -1.65$), severe drought ($-1.28 > Z_{ij} > -1.65$), moderate drought ($-0.84 > Z_{ij} > -1.28$), and no drought ($Z_{ij} > -0.84$) (Agnew and Chappel, 1999).

2.3.3 TREND ANALYSES

In this study, non-parametric Mann-Kendall's trend test were used for the assessment of past and future climate trend at annual and seasonal time scales. Mann-Kendall's trend test which is less sensitive to outliers and test for a trend in a time series without specifying whether the trend is linear or non-linear

The Mann-Kendall's test statistic was given as:

$$S = \sum_{i=1}^{N-1} \sum_{j=i+1}^N \text{Sgn} (X_j - X_i) \quad (\text{Eq 5})$$

Where, S was the Mann-Kendal's test statistics; X_i and X_j were the sequential data values of the time series in the years i and j ($j > i$) and N was the length of the time series. A positive S value indicates an increasing trend and a negative value indicates a decreasing trend in the data series.

The sign function was computed as:

$$\text{Sgn} (X_j - X_i) = \begin{cases} + 1 \text{ if } (X_j - X_i) > 0 \\ 0 \text{ if } (X_j - X_i) = 0 \\ - 1 \text{ if } (X_j - X_i) < 0 \end{cases} \quad (\text{Eq 6})$$

The variance of S , for the situation where there may be ties (that is, equal values) in the x values is given by:

$$\text{Var} (S) = \frac{1}{18} \left[N(N-1)(2N+5) - \sum_{i=1}^m t_i(t_i-1)(2t_i+5) \right] \quad (\text{Eq 7})$$

Where, m was the number of tied groups in the data set and t_i was the number of data points in the i^{th} tied group. For N larger than 10, Z_{MK} approximates the standard normal distribution (Yenigun *et al.*, 2008) and computed as follows:

$$Z_{MK} = \begin{cases} \frac{S - 1}{\sqrt{\text{Var} (S)}} \text{ if } S > 0 \\ 0 \text{ if } S = 0 \\ \frac{S + 1}{\sqrt{\text{Var} (S)}} \text{ if } S < 0 \end{cases} \quad (\text{Eq 8})$$

The presence of a statistically significant trend was evaluated using the Z_{MK} value. In a two-sided test for trend, the null hypothesis H_0 was accepted if $|Z_{MK}| < Z_{1-\alpha/2}$ at a given level of significance. $Z_{1-\alpha/2}$ was the critical value of Z_{MK} from the standard normal table. For example: for 5% significance level, the value of $Z_{1-\alpha/2}$ is 1.96. A positive value of Z_{MK} indicates an increasing trend while a negative value indicates a decreasing trend. In the present study, the significance of the observed change was examined at $p \leq 0.05, 0.01$, and 0.001 significance level. The Mann-Kendall's trend test done by using Excel template MAKESENS (Mann-Kendall test for trend and Sen's slope estimates) 1.0.

To analyse spatial distribution of climatic elements were rasterized to generate grided point data by the simple kriging interpolation technique using ARCGIS 10.1 software. This is because simple kriging interpolation technique takes account of the spatial correlation pattern with the least interpolation error (Beck *et al.*, 2005).

2.4 DETERMINATION OF RAINFALL CHARACTERISTICS

2.4.1 START , END AND LENGTH OF THE GROWING SEASON

In setting an onset date of the past records, many different criteria could be use for different crops exhibiting different maturity plus drought tolerance levels and soil types. Here, the one with 20 mm of total rainfall

received over three consecutive days that were not followed by greater than 10 days of dry spell length within 30 days from planting was adopted (Raman, 1974).

On the other hand, the end of growing season is mainly dictated by the water stored in soil and its availability to the crop after the rain stops. In this study 100 mm/meter of the plant available soil water and site specific daily reference evapotranspiration (ET_o) values were considered, and the end of the growing season was taken defined as any day after 1st of September for *kiremt* seasons when the soil water balance reaches zero (Stern *et al.*, 1982).

In determining the end date, set an estimated evapotranspiration showed in table 1 and 100 mm/m of the plant available soil water were considered. For the present study, Hargreaves and Samani (HS) method was applied to estimate ET_o based on observed daily temperature data after calibration and validation of HS model against of FAO Penman Monteith cropwat 8.0 calibrated by Bati station. Estimat ET_o at Bati, Kemisse, Artuma and take an average value for Senbete district.

Table 1. Estimated mean *kiremt* and *belg* season ET_o in mm/day after calibration

Seasons/ Stations	Bati	Kemisse	Artuma
Period	1995-2015	2004-2015	2008-2015
<i>Kiremt</i>	4.29	4.67	4.45
<i>Belg</i>	4.12	4.75	4.24

The onset and cessation of rainfall date are analyzed using an Instat version 3.37 package developed by the Statistical Services Centre of the University of Reading (Stern *et al.*, 2006).

Length of *kiremt* growing season (LGP) were determined as the difference between the end and start of rainy seasons. Mesay (2006), Hadgu *et al.* (2013) and Hadgu *et al.* (2014) used the same method to determined LGP.

2.4.2 RAINFALL TOTALS AND NUMBER OF RAINY DAYS

In the context of Ethiopia, Segele and Lamb (2005) employed three rainfall thresholds to define a rainy day (0.1mm, 0.5mm and 1mm) but a threshold value of 1mm were used to define days as wet or dry; because < 1mm of rainfall value almost has no effect on growth of crops (Robel *et al.*, 2013). Thus, in present study, number of rainy days were determined by counting all days with rainfall greater or equal to 1.0 mm as outlined by (NMSA, 2001). Seasonal (for *kiremt* June to September and for *belg* February to end of May) rainfall totals were determined as sum of rainfall of each day with greater or equal to 1 mm. Different researchers used the same methods (Segele and Lamb, 2005; Mesay, 2006; Hadgu *et al.*, 2013; Muluneh, 2015).

2.4.3 PROBABILITY OF DRY SPELL

The dry spell probabilities were determined as consecutive number of days with rainfall less than 1 mm per day exceeding 5, 7, 10 and 15 consecutive days. Dry spell length was analyzed by first order Markov Chain analysis (Stern *et al.*, 2006) using INSTAT v3.37 software.

3 RESULT AND DISCUSSIONS

3.1 DISTRIBUTION OF SEASONAL RAIN FALL

Figure 2. presents the monthly rainfall distribution of the study area. The highest rainfall was observed in the month of August across the Special Zone and the amount increased from Bati (209.7 mm) to Senbete (312.2 mm). While in *belg* season the highest monthly rainfall was observed in the month March at Bati and April at Kemisse, Artuma and Senbete (figure 3).




Figure 2. Mean monthly rainfall distribution at Oromia Special Zone, during 1986-2015




Figure 3. Spatial distribution mean seasonal rainfall 1986-2015.

Figure 3 shows that, in kiremt season lowest rainfall observed in northern part of the region (Bati) and highest rainfall recorded in southern part (Senbete) of the Special zone. But in belg season lowest rainfall performed at central part (Kemisse and Artuma) and highest rainfall records in the southern part (Senbete) of the special zone (figure 3).

3.2 SEASONAL RAINFALL VARIABILITY

Kiremt season rainfall variability In all the study stations as revealed in Table 2, the main rainy season (*kiremt*) contributes the maximum to the annual rainfall with the highest contribution in Kemisse (71.8%) and the lowest in Bati (64.2%). Similar result was also reported by Muluneh (2015) with contribution of seasonal *kiremt* rainfall to annual total rainfall varied from 60% at Srinka to 76% at Lalibela in NEA during the period of 1992-2012.

The mean total *kiremt* season rainfall varied from 484.1mm at Bati to 681.7mm at Senbete of the Special Zone with CV ranged from 22.6 % to 30.4% and SD ranged from ± 109.4 mm to ± 207.2 mm, respectively (table 2). The result showed that, the Special Zone experienced moderate rainfall variability and less stability of rainfall *kiremt* season. This less stability showed that the *kiremt* rainfall totals were not easily predictable and could result in difficult decision regarding rain fed crop production during the study period. Similar results were also reported by Muluneh (2015), Hadgu *et al.* (2013) and Bewket (2009) in the north and north eastern parts of Ethiopia. The PCI of the *kiremt* season revealed values between 11 and 16, which that implies the Special Zone had moderate monthly rainfall distribution for the *kiremt* season (Table 2).

Table 2. Past seasonal rainfall Variability at OSZ, during (1986-2015)

Seasons	Districts	Statistics						
		Min (mm)	Max (mm)	Mean (mm)	SD (mm)	CV%	CT%	PCI
<i>Kiremt</i>	Bati	301.0	728.0	484.1	109.4	22.6	64.2	13.1
	Kemisse	310.0	866.0	594.1	157.3	26.5	71.8	13.4
	Artuma	273.0	852.0	546.0	132.8	24.3	68.7	13.5
	senbete	185.4	1040.0	681.7	207.2	30.4	68.7	13.0
<i>Belg</i>	Bati	35.0	472.0	184.6	108.4	58.7	24.5	15.1
	Kemisse	35.0	419.0	164.9	105.8	64.2	19.9	15.3
	Artuma	10.0	511.0	164.4	118.4	72.1	20.7	14.9
	Senbete	48.0	611.0	216.6	129.5	59.8	21.8	13.9

Where, SD is standard deviation, CV is coefficient of variation, CT is percent of of seasonal totals contribution for annual total rainfall and PCI is precipitating concentration index.

Belg season rainfall variability The contribution of *belg* season (FMAM) to the annual rainfall totals varied from 19.9% in Kemisse to 24.5% in Bati (Table 2). This indicated that the contribution of the seasonal *belg* rainfall to the annual total was less than half of the contribution of the *kiremt* season rainfall. This result also agreed with the findings of Muluneh (2015) and Bewket and Alemayehu (2016) who reported the same comparable result.

The mean seasonal *belg* rainfall totals ranged from 164.4mm at Artuma to 216.6mm at Senbete station of the Special Zone with CV ranged from 58.7% to 72.1%, and the SD value ranged from ± 105.8 mm to ± 129.5 mm, respectively (table 2). This implies that the seasonal *belg* rainfall was characterized by high variability as the classification of Hare (1983). Similar result were it has been reported by Muluneh (2015) also reported 150.5mm at Kobo and 271.8 mm at sirinka in NEA during 1992-2012. Likewise, high seasonal *belg* rainfall variability was reported by Bewket (2009), over the Amhara Region, and Bewket and Alemayehu (2016) over North Shewa for the time period of 1975-2003, 1979-2008, and 1983-2013, respectively.

The PCI value of the *belg* seasons varied between 11 and 16, that indicates the zone received small amount rainfall with moderate monthly rainfall distribution in the study period (Table 2). The small amount rainfall observed with less stability and high variability indicate that the rainfall was not dependable and easily predictable, and the season was unlikely to sustain agricultural crop production on rainfall during 1986-2015.

3.3 SEASONAL RAINFALL ANOMALY

Kiremt seasonal rainfall anomaly The *kiremt* rainfall anomaly showed that 43.3% of the 30 years occurrence of rainfall deficit and 46.7% of 30 years indicated positive departure from the normal or occurrence of good rainfall during the study years (Figure 4A). Among the 13 dry years moderate to severe drought years were observed in 1987, 1989, 1990, 1991, 1993 and 2015. Among 14 positive years 1988, 1998, 1999, 2000, 2007 and 2010 were moderate to extreme wettest years in *kiremt* season of the study period (1986-2015). Korecha (2013) has also found that, the years 1987, 1991, 1997, and 2009 recorded negative anomaly and the years 1988, 2006 and 2007 recorded positive anomaly of *kiremt* rainfall over Ethiopia. This implies could be challenged that rain fed crop production in the Special Zone has been challenged by risk of dry years during the study period. In agreement with this, Bewket (2009), Ayalew *et al.* (2012) and Muluneh (2015) noticed that the rain fed agriculture was highly at risk in the NEA.

Belg seasonal rainfall anomaly The *belg* rainfall anomaly showed that 50% of the 30 years experienced negative departure from normal (rainfall deficit) and 33.3% of the 30 years showed positive departure from normal (good rainfall) (Figure 4B). Among those 15 dry years moderate to severe drought years are observed 1994, 1999, 2008, 2009 and 2013, and among the 10 wet years 1987, 1993, 1995 and 1996 recorded moderate to extreme wettest years in *belg* season of the study period (1986-2015). This indicates that drought frequencies were high in *belg* season than the *kiremt* and the annual records. Comparable result were also reported by Muluneh (2015), where the years, 1992, 1999, 2000, 2008, 2009 and 2011 recorded below mean rain fall in NEA during 1992-2012.




Figure 4. Areal mean annual and seasonal rainfall anomalies at OSZ, during 1986-2015

3.4 RAINFALL TREND

The Mann–Kendall trend test showed non-significantly increasing trends of *kiremt* rainfall at all stations while there was decreasing trend for *belg* total rainfall with statistically significant trends at Kemisse and Artuma (Table 3). In line with the result, Muluneh (2015) found that, increasing trend *kiremt* rainfall over the majority of the study area and the *belg* rainfall has shown a declining trend across the NEA during 1992-2012. Likewise, Hadgu *et al.* (2013) reported a increasing *kiremt* rainfall trend at at Alamata and Edagahamus and decreasing trend in *belg* rainfall total at all study stations during the period 1980-2009.

Table 3. Trend of past seasonal rainfall at OSZ, during 1986-2015.

Stations	<i>Kiremt</i>		<i>Belg</i>	
	Z _{MK}	Slop	Z _{MK}	Slop
Bati	1.18 ^{ns}	3.75	-0.82 ^{ns}	-2.40
Kemisse	0.25 ^{ns}	1.19	-1.98 ^c	-4.14
Artuma	0.62 ^{ns}	1.67	-1.98 ^c	-4.80
Senbete	1.86 ^{ns}	9.15	-1.70 ^{ns}	-4.67

Where, Slope (Sen’s slope) is the rate of change mm/year, ^{ns} is non-significant at all level of significance level and ^c is trend at 0.05 level of significance.

3.5 RAINFALL CHARACTERIZATION

3.5.1 PAST START AND END OF SEASONS

Start of *belg* season The *belg* season fulfilled the criteria of start of season in the past 30 years only for 14 years (46.7%), 10 (33.3 %) years, 8 (26.7 %) years and 15 (50%) years at Bati, Kemisse, Artuma and Senbete, respectively. Except at Senbete the median of the start of *belg* season disclose between March and May (Table 4). During 1986-2015, the *belg* mean started on DOY 141(May 20), 156 (Jun 4), 162 (Jun 10) and 138 (May17) at Bati, Kemisse, Artuma and Senbete, respectively experiencing high variability (>30 %) and low stability (>50 days). This indicated that, the start of *belg* season show low predictability of effective onset date/erratic and difficult to adopt fixed recommendations (date of sowing, cultivars, planting density, fertilizer rates and time of application) because of failure of start date of growing season across the study area. This shows that there was high risk of crop failure that is planting in *belg* season in the study area during 1986-2015.

Table 4. Past start date of *belg* season at OSZ, during 1986-2015

Statistic	<i>Belg</i> SOS			
	Bati	Kemisse	Artuma	Senbete
Early (DOY)	62 (Mar 2)	65 (Mar 5)	65 (Mar 5)	65 (Mar 5)
1 st Qrt (DOY)	79 (Mar 19)	98 (Apr 7)	126 (May 5)	85 (Mar 25)
Median (DOY)	179 (Jun 27)	186 (Jul 4)	189(Jul 7)	131(May 10)
3 rd Qrt (DOY)	191(Jul 9)	192 (Jul 10)	193(Jul 11)	192(Jul 10)
Late (DOY)	213(Jul 31)	219(Aug 6)	219 (Aug 6)	221 (Aug 8)
Mean (DOY)	141(May 20)	156 (Jun 4)	162 (Jun 10)	138 (May17)
SD (DOY)	56.6	51.8	51.1	56.1
CV (%)	40.2	33.1	31.5	40.7

Start *Kiremt* Season Summary statistics for the past start of *kiremt* growing season during 1986-2015 in the OSZ is described in Figure 5. The median for onset of *kiremt* rainfall in the OSZ were 190 DOY (Jul 8) at Kemisse and Senbete, and 192 DOY (Jul 10) at Bati and Artuma. In line with the present result, Muluneh (2015) also presented the median SOS of *kiremt* growing season of 187 DOY (Jul-5), 190 DOY (Jul-8), 187 DOY (Jul-5) and 187 DOY (Jul-5) at Kombolcha, Kobo, Lalibela and Srinka, respectively with CV ranging from 4% to 6% during 1992-2012. Ayalew *et al.* (2012) has also found that 189 DOY (July-9) and 186 DOY (July-4) as the median SOS of the *kiremt* growing season at Srinka and Kombolcha, respectively, for the period 1978-2008.




Figure 5. Box and whisker plots for start and end of *Kiremt* growing season at OSZ, during 1986-2015.

End of *Kiremt* Season This also shown in Figure 5, the Special Zone, the median end date of *kiremt* season (EOS) was on DOY 291 (Oct 17), 299 (Oct 25), 301 (Oct 27) and 307(Nov 2) at Bati, Kemisse, Artuma and Senbete, respectively with SD ranges from ± 13.1 days to ± 14.5 days and CV ranges from 4.3% to 5% at Senbete and Bati, respectively. The main rainy season terminated during the last days of 280 DOY (Oct 6), 290 DOY (Oct 16), 293 DOY (Oct19) and 299 (Oct 25) once in four years' and terminated earlier than 300 DOY (Oct 26), 307 DOY (Nov 2), 309 DOY (Nov 4) and 315 DOY (Nov 10) at Bati and Senbete in three out of four years, respectively (Figure 5). Similar to this study, Ayalew *et al.* (2012) found that. the average ending date of *kiremt* rainfall ranged from DOY 266 (September 22) at Mehal Meda to DOY 304 (October 30) at Debark at ANRS.

3.5.2 LENGTH OF SEASONS AND NUMBER OF RAINY DAYS

A further note could also be made from Table 5 that rainfall season duration was dependent mainly on the onset date in the northern part of the zone (at Bati), rainfall duration was lower than 88 days in only 25 % of the years, while it was lower than 111 days in 75% of the years. Similarly, the lower quartile for rainfall duration in the southern part of the special zone (at Senbete) was below 102 days in 25% and below 128 days for 75% of the study years; while in the central part of the zone the lower and upper quartiles were 99 and 121 at Kemisse and 96 and 116 at Artuma, respectively.

Table 5. Past length of growing period and number of rainy days in *kiremt* growing season at OSZ, during 1986-2015.

Statistic	Length of Growing Period				Number of Rainy Day			
	Bati	Kemisse	Artuma	Senbete	Bati	Kemisse	Artuma	Senbete
Short (days)	58	76	76	57	29	21	23	10
1 st Qrt (days)	83	99	96	102	39	45	40	45
Median (days)	101	107	107	119	45	50	43	51
3 rd Qrt (days)	111	121	116	128	48	56	49	57
Long (days)	129	133	139	146	61	73	75	71
Mean (days)	98	108	106	115	43	50	45	50
SD (days)	17.5	15.6	14.8	18.4	7.2	10.3	9.7	11.8
CV (%)	17.8	14.4	13.9	16.0	16.8	20.8	21.6	23.5

The variability of LGP ranged from 13.9% at Artuma to 17.8 at Bati (Table 5). This clearly showed that the central and southern part of the special zone have stable and longer length of growing period than the north part of the zone. Similar to this study, Ayalew *et al.* (2012) reported that Sirinka, Mahil Meda and Kombolcha experienced the least duration (<100 days) of *kiremt* rainfall, where as, Motta, Bahir Dar, Debark and Debre Markos experienced long duration (>140 days) of *kiremt* rainfall.

Similar to LGP, number of rainy days (NRD) between start and end of seasons (Table 5), indicated that at Bati the number of rainy days less than 39 days in only 25 % of the years, while it was lower than 48 days in 75% of the years with midian of 45 days.

This showed that NRD increased from north to south, but the variability was reversed across the special zone.

3.5.3 PROBABILITY OF DRY SPELL

The risk of getting 5,7,10 and 15 days of dry spells based on the fitted first order Markov chain probability values at OSZ during 1986-2015 is depicted in Figure 6. In *belg* season the probability of dry spell occurrence of 10 days' length was 60% at Bati, Kemisse, and Senbete and 70% at Artuma during the peak rainfall months (March and April) in *belg* season only between 88 DOY (March 28) and 118 DOY (April 27), respectively. Means that, the risk of dry spell occurrence of 5,7 and 10 days are greater than 60% during the *belg* season. On the other hand, the observed probability of dry spell occurrence greater than 15 days' length was less than 60% at Bati, Kemisse and senbete between 85 DOY (mar 25) and 121 DOY (April 30) and less than 50% at Artuma between 82 DOY (Mar 22) and 130 DOY (May 9).




Figure 6. Probabilities of maximum dry spells exceeding 5, 7, 10 and 15 days' length in the OSZ, during 1986-2015

In *kiremt* season, the dry spell risk of being greater than 10 day consecutive dry spells was less than 50%, and it is expected to occur at Bati from 184 DOY (July 2) to 256 DOY (Sep 12), at Kemisse 178 DOY (June 26) to 253 DOY (Sept 9), at Artuma 184 DOY (July 2) to 259 DOY (Sep 15), and at Senbete 169 DOY (June 17) to 262 DOY (Sept 18) (Figure 4). This information is very important to farmers in the study area to decide on crop types to be cultivated and on planning sowing dates as a function of observed from the onset dates. In rain-fed farming, the intermittent dry spell becomes critical, particularly for the seedling establishment during the first 30 days or so after planting. Also used in decision making with respect to selecting dry spell resistance crop and field operations within the farming system.

3.6 TEMPERATURE VARIABILITY AND TREND

3.6.1 SPATIAL VARIABILITY TEMPERATURE

Maximum temperature (Tmax) The special zone experienced in *kiremt* season temperature of 29.3 °C to 33.1 °C, and in *belg* season temperature of 28.4 °C to 31.6 °C (Figure 7). As is apparent from figure 7, the special zone has experienced warmer in *kiremt* season than the *belg* season. The special zone with mean Tmax 28.3 °C at Bati, 31.5 °C at Kemisse and Artum and 29.8 °C at senbete. This shows that the central part is warmer than the north and southern part of the Special Zone.




Figure 7. Spatial variability of mean annual and seasonal maximum temperature at OSZ, during 1986-2015

Minimum Temperature (Tmin) In the study area the of *kiremt* season Tmin was from 15.1 °C to 17.8 °C, and the *belg* season temperature varied from 14.3 °C to 16.0 °C (Figure 8). Among the district in the study Zone, Bati district was colder in terms of *belg* season Tmin, while during the *kiremt* season Artuma district colder over other districts




Figure 8. Spatial variability of mean annual and seasonal minimum temperature at OSZ, during 1986-2015

3.6.2 SEASONAL TEMPERATURE TREND

Kiremt season temperature trend The *kiremt* Tmax showed increasing trend at all stations studied in with statistically significant trend at Bati, Kemisse and Artuma stations. The *Kiremt* Tmax increment varied from

0.09 °C per decade in the central part of the zone to 0.40 °C per decade in north part of the special zone. Observed Tmin of *kiremt* also showed positive trend but the trends were non-significant (Table 6).

Table 6. Trend of mean annual and seasonal maximum and minimum temperature at OSZ, during 1986–2015

Districts	<i>Kiremt</i>		<i>Belg</i>		
	Z _{MK}	Slope	Z _{MK}	Slope	
Maximum Temperature	Bati	2.32 ^c	0.040	4.07 ^a	0.082
	Kemisse	2.57 ^c	0.038	4.60 ^a	0.082
	Artuma	2.18 ^c	0.009	3.28 ^b	0.018
	Senbete	1.75 ^{ns}	0.023	4.21 ^a	0.070
Minimum Temperature	Bati	0.54 ^{ns}	0.007	-2.28 ^c	-0.045
	Kamisse	1.78 ^{ns}	0.019	-0.25 ^{ns}	-0.003
	Artuma	0.96 ^{ns}	0.006	-1.39 ^{ns}	-0.007
	Senbete	1.43 ^{ns}	0.017	-0.68 ^{ns}	-0.007

Where, Z_{MK} is Mann–Kendall test statistic, Slope (Sen's slope) is the rate of change (Degree Celsius) per year; **a** is 0.001 level of probability; **b** is 0.01 level of probability; **c** is 0.05 level of probability. and **ns** is non-probability at 0.05 probability level.

Belg season temperature trend The *belg* Tmax showed increasing trend at all stations studied years in the zone with statistically significant. The increment varied from 0.18 °C per decade to 0.82 °C per decade. Conversely, the Tmin of *belg* showed negative trend at all studied stations of the special zone but non-significant except in the north part of the special zone (Table 6).

CONCLUSIONS

This study was undertaken to analyze the characteristics of historical girded climate data in OSZ of ANRS. Analysis of climate characteristics for the study area 1986-2015 for past. The area received mean total rainfall in *kiremt* season varied from 484.1mm to 681.7mm and in *belg* season it received varied from 164.4mm to 216.6mm, respectively during 1986-2015. The past three decades seasonal rainfall trend test showed increasing in *kiremt* rainfall while decreasing in *belg* rainfall at all studied station of the special zone for that past. Regarding temperature analyses, except *belg* minimum temperature increasing trends in the study area during the study period.

During 1986-2015, the *belg* mean start of season 141 DOY (May 20), 156 DOY (Jun 4), 162 DOY (Jun 10) and 138 DOY (May17) at Bati, Kemisse, Artuma and Senbete, respectively with experienced high variability (>30 %) and low stability (>50 days). The median of *kiremt* season onsets are 190 DOY (Jul 8) at Kemisse and Senbete and 192 DOY (Jul 10) at Bati and Artuma with CV ranges 4.8 % at Artuma and 7.0% at Senbete and SD ranges from ± 9.3 days at Artuma and ± 13.3 days at Senbete, respectively. The variability of LGP ranges from 13.9% at Artuma to 17.8% at Bati and NRD ranges from 16.8% at Bati to 23.5 % at Senbete of the Special Zone.

The occurrence of dry spell probability 5,7 and 10 days are greater than 60% during the *belg* season and in *kiremt* season, the dry spell risk of being 10 Day consecutive dry less than 50% be fallen at Bati from 184 DOY (July 2) to 256 DOY (Sep 12), at Kemisse 178 DOY (June 26) to 253 DOY (Sept 9), at Artuma 184 DOY to 259 DOY (Sep 15), and at Senbete 169 DOY (June17) to 262 DOY (Sept 18). This information is very important to the farmers to decide on crop types to be cultivated and on planning sowing dates as a function of observed from the onset dates. Also used decision making with respect to selecting dry spell resistance crop and field operations within the farming system.

REFERENCE

- Agnew, C.T. and A. Chappel. 1999. Drought in the sahel. *GeoJournal* 48: 299-311. agrometeorological indices. *Journal of Agro Meteorology (1): 111 – 114.*
- Ayalew Dereje, Kindie Tesfaye, Girma Mamo, Birru Yitafaru and Wondimu Bayu. 2012. Variability of rainfall and its current trend in Amhara Region, Ethiopia. *African Journal of Agricultural Research,7(10):1475–1486.*
- Beck, C., Grieser, J., Rudolf, B., 2005. A new monthly precipitation climatology for the global land areas for the period 1951 to 2000. *German Weather Service, Climate status report 2004, pp. 181–190.*
- Bewket Weldeamlak. 2009. Rainfall variability and crop production in Ethiopia; Case study in the Amhara region; *proceeding of the 16th International Conference of Ethiopian Studies, Addis Ababa, Ethiopia.*
- Bewket Woldeamlak and Alemayehu Arragaw. 2016. Local climate variability and crop production in highlands

- of Ethiopia. *Environmental Development*, 19: 36 – 48. <http://dx.doi.org/10.1016/j.envdev.2016.06.002>
- Godswill, M., Dawit, K. and Dejene, A. 2007. A comparative analysis of rain fed and irrigated agricultural production in Ethiopia. *Journal of Irrigation and Drainage. Vol 21 No 1. Springer Netherland.*
- Hadgu Gebre, Tesfaye Kindie and Mamo Girma. 2014. Analysis of climate change in Northern Ethiopia: implications for agricultural production. *Theoretical and Applied Climatology ISSN 0177798X.Vol.117: P 3-17.*
- Hadgu Gebre, Tesfaye Kindie, Mamo Girma and Kassa Belay. 2013. Trend and variability of rainfall in Tigray, Northern Ethiopia: Analysis of meteorological data and farmers' perception. *Academically Journal Environmental Science, 1(8): 159-171.*
- Hagos, F., Makombe, G., Namara, R.E. and Awulachew, S.B. 2009. Importance of irrigated agriculture to the Ethiopian economy: Capturing the direct net benefits of irrigation. Colombo, Sri Lanka: *Int. Water Manage. Inst. p.37.*
- Hare, F.K. 1983. Climate and desertification. Revised analysis (WMO-UNDP) WCP-44: 5-20. Geneva, Switzerland.
- Korecha, D. 2013. Characterizing the predictability of seasonal climate in Ethiopia. Doctoral dissertation in meteorology. University of Bergen, Norway.
- Mandal, K.G., Kumar, A., Ghosh, S., Kundu, D.K., Panda, R.K., Mohanty, R.K., Raychaudhuri, M., Padhi, J., Majihi, P. and Sahoo, D.K. 2013. Analysis of rainfall and soil characteristics of kuanria canal command for water resource management research bulletin.
- Mavi, H.S. and Tupper, G.J. 2004. Agro meteorology: principles and applications of climate studies in agriculture. *Food Products Press®, an imprint of the Haworth Press, Inc., 13: 904-1580.*
- Mesay Abebe. 2006. The onset, cessation and dry spells of the small Rainy Season (*belg*) of Ethiopia. National Meteorological Agency of Ethiopia, Addis Ababa. 10-13P.
- Muluneh Getahun. 2015. Analysis of past and future intra-seasonal rainfall variability and its implications for crop production in the North Eastern Amhara Region, MSc. Thesis Haramaya University, Haramaya, Ethiopia.
- NMA (National Meteorology Agency). 2007. Climate change national adaptation programme of action (NAPA) of Ethiopia: technical report, National Meteorological Agency, Addis Abeba, pp. 85
- NMSA (National Meteorology Service Agency). 2001. Initial national communication of Ethiopia to the united nations framework convention on climate change (UNFCCC), Addis Ababa, Ethiopia, pp. 113.
- Oliver, J.E., 1980. Monthly precipitation distribution: a comparative index. *The Professional Geographer*, 32: 300-309.
- Raman, C.R.V. (1974). *Analysis of Commencement of Monsoon Rains over Maharashtra State for Agricultural Planning*. Published Scientific Report No. 216. Poona, India: India Meteorological Department.
- Robel, A. A., E. DeGiuli, C. Schoof, and E. Tziperman. 2013. Dynamics of ice stream temporal variability: Modes, scales, and hysteresis, *J. Geophys. Res. Earth Surf., 118: 925–936.*
- Segele, Z. T. and P. J. Lamb. 2005. Characterization and variability of Kiremt rainy season over Ethiopia. *Meteorology and Atmospheric Physics, 89(1): 153–180.*
- Stern, R., Rijks, D., Dale, I. and Knoch, J. 2006. *Instat climatic guide*. Statistical services center, Reading University, UK.
- Stern, R.D. and Coe, R. 1982. The use of rainfall models in agricultural planning. *Agric. Meteorol., 26: 35-50.*
- Yemenu, F. and Chemed, D. 2013. Dry and wet analysis of two rainy seasons for decision support in agricultural water management for crop production in central highlands of Ethiopia. *Journal of Biological and Agricultural Health care, 3(11):1-6*
- Yenigun K, Gumus V, Bulut H (2008) Trends in stream flow of the Euphrates basin. Turkey. *Proc Inst Civil Eng Water Manag* 161: 189–198. doi:10.1680/wama.2008.161.4.189