

Pre-extension Demonstration of Dolichos lablab (Lablab purpureus) under sown in Maize at Dugda and Lume Districts of East Shoa Zone, Oromia, Ethiopia

Daniel Wana Dawit Abate Nabi Husen Meseret Tilahun
Oromia Agricultural Research Institute,
Adami Tulu Agricultural Research Center, P.O. Box 35, Batu, Ethiopia

Abstract

The activity was conducted in Bekele-Girisa of Dugda district. The objectives were to demonstrate forage production from lablab under sown in maize and to evaluate the lablab-maize intercropping practices with farmers participation. Accordingly, Farmers' Research and Extension Groups (FREGs) with 18 members were established at Bekele Girisa kebele. Four trial farmers were selected from each FREGs for forage production based on the criteria including interest of the farmers in producing forage on their land, farmer having enough land for forage production and who have milking cows. BH-540 maize variety was intercropped with Lablab purpureus on plot a size of 32*32m². Pure stands of maize with similar plot size was also established for comparison. Lablab purpureus was intercropped in maize at two weeks period after maize planting. Farmers were used as replication with participatory approach. The results revealed that the total biomass (Stover + lablab) and maize grain yields were significantly different ($P < 0.05$) among the practices at Dugda site. Even though statistically not significant ($P > 0.05$) between the two practices, there was total biomass and maize grain yield increment in lablab under-sown in maize than pure stand of maize. Similarly, crude protein content of the maize Stover under sown with lablab was improved. Farmers were also very interested for forage production from lablab intercropping in maize as compared to sole maize production practice. Therefore, it is recommended to extend the forage production strategy such as lablab intercropping in maize to enhance the production of high biomass of forage with good quality in the study area.

Keywords: biomass yield, FREG, intercropping, Lablab

DOI: 10.7176/JBAH/11-15-04

Publication date: August 31st 2021

Introduction

In mixed crop-livestock systems, livestock feed supply is mainly dependent on crop residues, natural pastures, and other agricultural by-products. However, the quantity and quality of the available feed resources is declining from time to time as most of the available land is cultivated for crop production (Tolera *et al.*, 2012).

Forage legumes integrated with food crops and livestock is often advocated to minimize external inputs as well as to improve the productivity and sustainability of crop-livestock production in developing countries (Giller., 2001; Peters and Lascano., 2003). Forage legumes provide food, feed and facilitate soil nutrient management.

Intercropping is a type of mixed cropping agricultural practice of cultivating two or more crops in the same space at the same time. Intercropping of cereals with legumes has been popular in tropics (Hauggaard-Nielsen *et al.*, 2001; Tsubo *et al.*, 2005) and rain-fed areas of the world (Banik *et al.*, 2000; Ghosh 2004; Agegnehu *et al.*, 2006; Dhima *et al.*, 2007) due to its advantages for soil Conservation (Anil *et al.*, 1998), weed control (Poggio 2005; Banik *et al.*, 2006), lodging resistance, yield increase (Anil *et al.*, 1998; Chen., 2004), and legume root parasite infections control (Fenandez-Aparicio *et al.*, 2007). Different studies also indicated that forage legumes integration through intercropping did not have a significant effect on maize grain and biomass yield (Mergia Abera, 2014). The feasibility of intercropping lablab in maize for additional feed source was investigated and promising results were obtained and recommended for the end users (Diriba Geleti and Lemma Gizachew 2003).

However, this intercropping practice was not demonstrated to small scale farmers and evaluated at on-farm condition with farmers' participation. Therefore, this study was designed to evaluate and demonstrate the practice of lablab intercropping in maize to improve livestock feed production in the study area.

Materials and Methods

Description of the study area

The study was carried out at Dugda districts of East Shoa zone. One kebele; Bekele-Girisa was selected from the District based on the livestock population potential, severity of feed shortage and cropping system (maize dominant). Geographically Dugda district is located between 8⁰⁰1'N to 8⁰¹0'North latitude and 38⁰³1'E to 38⁰⁵7'E longitude. Meki, the capital city of Dugda district, is located 134 km to the South East of Addis Ababa

on the main road to Ziway town. The altitude of the study area ranges from 500 to 2000 (m.a.s.l). The area receive an erratic, unreliable and low rainfall, averaging between 500 and 900 mm per annum. The rain fall is bi-modal with the long rain lasting from June to September (Abule *et al.*, 1999).

Farmers' selection

The activity was carried out using Farmers' Research and Extension Groups (FREGs) of smallholder farmers. Accordingly, FREGs with 18 members were established at Bekele Girisa kebeles. Detailed analysis of the problem and potential benefits of improved forage production and utilization were discussed with farmers. Four trial farmers were selected from each site for the forage production based on the criteria including interest of the farmers in producing forage on their land, farmer having enough land for forage production and farmers who have milking cows.

Trial establishment and management

BH-540 maize variety was planted as intercrop with *Lablab purpureus* on plot size of 32*32m². With farmers practice, pure stands of maize crop of the same variety with similar plot size was also established as comparison. Seed rate of 25 kg/ha with 75cm of spacing between the rows, and 25 cm among the plants were used for maize crop. *Lablab purpureus* was intercropped between the maize rows at seed rate of 15kg/ha (half of the recommended seed rate for sole production) two weeks after maize planting. Trial farmers were used as replication with participatory approach. NPS fertilizer at rate of 100kg/ha was applied at planting. All other recommended agronomic practices were done for all plots uniformly.

Farmer's training and evaluation of forage development technologies

Theoretical training was given for group members on forage production and utilization before planting. Then practical training was given for the group at each farm; where the trial was conducted to address the crop establishment, general management, harvesting and feeding system. Neighbors were encouraged to attain the training. Farmers carried out qualitative evaluation of the forage intercropping system through matrix ranking. They critically evaluated forage production strategy based on their criteria. Farmers of the two districts used almost similar criteria for evaluation of the forage production strategy. The major criteria considered in the evaluation includes; herbage biomass yield, multipurpose use of the technology, protection of soil erosion, ability of drought tolerance, improvement of soil fertility and compatibility of the technology with the existing production system, Finally they selected forage production strategies suitable to their farming condition.

Biomass yield advantage determination

Biomass yield advantages of the forages were determined by comparing the biomass yield obtained from forage intercropped (maize-lablab) and sole maize farming practices using the following formula:

$$\text{Biomass yield advantage \%} = \frac{\text{Yield of intercrop (t/ha)} - \text{Yield of sole (t/ha)} \times 100}{\text{Yield of sole (t/ha)}}$$

Data collection and analysis

Relevant agronomic and yield including plant height, biomass yield of lablab, maize Stover and seed yield were collected. The data was organized and analyzed to describe various variables using Microsoft Excel and Statistical Package for Social Sciences (SPSS 20). The student t-test was used for mean separation.

Result and discussions

Agronomic and yield performances of Lablab under-sown in maize

Agronomic and yield performances of lablab under-sown in maize at Bekele Girisa site of Dugda district is presented in table 1. At Dugda site the total biomass yield and seed yield differ significantly ($p < 0.05$). Other tested parameters like plant height (cm) and crude protein (cp) were not significantly ($p > 0.05$) different among the two practices. The total biomass yield (7.20 t/ha) and maize grain yield (52.7 quintal/ ha) were recorded for maize-lablab intercropping practice. Due to the lablab intercropping the total biomass was increased by 9.1% at study area site.

The higher maize grain yield recorded for the intercropping could be due to the better maize crop management since the recommended seeding rate and spacing were used for the intercropping practice. In farmers practice (sole maize production), farmers used lower spacing between rows and plants that could be a cause for lower maize grain yield recorded. In addition, the under sown forage legumes help in suppressing the growth of unwanted weeds and conserve moisture in the soil. This result is in agreement with Mergia Abera (2014), where inclusion of vetch, cowpea and lablab increased grain yield of maize by 7.4%, 5.9%, and 5%, respectively. However, the results of this study is contrary to those reported by Aklilu *et al.* (2007) and Mpairwe *et al.* (2002) where the inclusion of forage legumes depressed grain yield of companion cereals by 3.6 to 9%.

Abubeker (2006) also reported that simultaneous planting of lablab significantly ($P < 0.05$) reduced grain and stover yield but increased forage dry matter (DM) yield. However, delayed planting, did not affect ($P > 0.05$) grain, Stover, forage dry matter (DM) or total fodder yields.

Even though it is not significantly ($p > 0.05$) different, the crude protein content of maize stover under sown with lablab was greater than that of maize Stover from pure stands (sole maize). Similarly, Mergia Abera (2014) also stating that crude protein content was not significantly different ($p > 0.05$) among the maize stovers samples taken from maize-lablab intercropped and sole maize treatments. Crude protein content of most cereal crop residues are lower than 7 % which is the critical level of microbial protein synthesis of feed intake (Adugna *et al.*, 1999). However, due to lablab-intercropping in maize the crude protein content of maize stover was above the critical level. This indicates that maize under-sown with forage legumes improve the crude protein quality of stover than pure stand(sole) maize sown.

Table1: Agronomic and yield performance of maize-lablab intercropping at Bekele Girisa site of Dugda district.

Practices	PH(cm)	DMY (tone/ha)			MSY (quintal /ha)	CPMS(%)
		Stover	Lablab	Total		
Sole maize	226.50	6.60	-	6.60	51.40	7.47
Maize –lablab intercropping	227.70	6.15	1.05	7.20	52.7	7.67
Mean	227.10	6.38	-	7.18	52.05	7.57
Standard Error	5.22	0.36	-	0.35	0.41	0.21
Sig. level	Ns	Ns	-	*	*	Ns

PH=plant height of Maize; DMY= Dry matter yield; MSY= Maize seed yield; CPMS= Crude protein of maize Stover.

Biomass yield advantages

The biomass yield advantage of maize lablab intercropping practices were 9.1% at Dugda site. This indicates that the intercropping practice was more advantageous than sole maize cropping. Intercropping of forage legumes and cereals generally results in higher fodder protein yield than cereal alone. However, fairly high yielder and quality forage legumes are needed to augment the cereal residues in order to produce a feed which would be capable of meeting the basal nutritional requirements of ruminants.

Training

Theoretical and practical training was given for FREGs and neighbors farmers on forage production and utilization before forage technology establishment and during forage harvesting. A total of 50 farmers (32 males and 12 females) participated in training on forage production and utilization practices. The training was mainly focused on forage crop establishment, general management, harvesting, storage and feeding system.

The reaction of participating farmers in terms of the advantages and drawbacks of the forage production by under-sowing forage legumes in maize crop as compared to pure maize production practice (monoculture) were indicated in table 3. According to the participant farmers, district animal feed experts and development agents maize-lablab intercropping was found as better strategy for forage and maize production as compared to sole maize cropping. This is mainly due to the benefits of under-sowing forage legumes in maize crop including additional quality feed production from lablab, soil fertility improvement, protection of soil erosion, ability of drought tolerance and compatibility of the technology to the existing production system.

All participant farmers were very much impressed and interested to grow lablab forage in maize crop after they have realized the benefits of the intercropping practice. They also understood that one can produce forage crops by under-sowing without competing land for crop production. Farmers also had obtained good awareness regarding improved forage production and utilization practices. They were encouraged in participation of the forage production and promotes the adoption of improved forage technologies in the study area. On the other side, there was increased realization on the part of researcher and extension workers that the technology became effective and acceptable by the farmers when the farmers themselves are involved in the research and extension program. It also benefited the researchers and extension workers in gaining and understanding of farmer's evaluation criteria and created good opportunity to communication with farmers.



Figure 1 During the training was given at Dugda district.

Table 3: Farmer's criteria for evaluation of lablab legumes under sown in maize and pure stand maize production practices (High score = 5 and least score = 1) and number of evaluating farmers =50

Evaluation parameters	Sole maize	Maize + lablab
Biomass yield.	4	5
Multi-purpose use as food & feed.	3	5
Protection of soil(water runoff protection)	3	5
Drought tolerance.	4	5
Moisture conservation and soil fertility improvement	3	5
Maize grain yield improvement	3	4
Total score	20	29
Rank	2nd	1st

Conclusion

The result of the current study indicated that the total biomass (maize + lablab) and maize seed yields of lablab forage legume under-sown in maize were performed better than the pure stand of maize production. Similarly, the changes in crude protein of forage were more pronounced in maize-legume intercrops than in pure cropped maize. Even though, the amount of biomass yield obtained from lablab legumes was low, the fact that the yield

obtained was without affecting maize grain yield makes the technology of lablab under-sowing in maize strategy attractive. Hence, those farmers practicing maize-legume intercropping could obtain more benefits in terms of food and animal feed than those practicing mono cropping.

Farmers evaluation result showed that the participating farmers were also very much interested in lablab under-sowing in maize crop as forage development strategy to solve animal feed shortages of the study area. Hence, small holder farmers are encouraged to produce lablab by under sowing in maize to enhance dry season feed availability and quality. Moreover, further studies on other forage legume species should be evaluated for their compatibility when under-sown/intercropped in food crops.

ACKNOWLEDGMENTS

In depth grateful thanks for Agricultural Growth Programs (AGP-II) for funding the research work. We are also gratitude appreciation to Adami Tulu Agricultural Research Center for their support in logistic facilitation and all Animal feed and nutrition research team members who were actively involved in field management and data collection during the research work.

References

- Abubeker Hassen., Lemma, G., and N.F.G. Rethman.2006. Effect of Lablab purpureus and Vicia atropurpuria as an intercrop, or in a crop rotation, on grain and forage yields of maize in Ethiopia. *Tropical Grasslands* (2006) Volume 40, 111–118.
- Abule E, Gremew E, Aliye H 1999. Adami Tulu Agricultural research Center, Bulletin No.1, Oromia Agricultural Development Bureau. Fifnfine, Ethiopia.
- Adugna Tolera, T. Berg and F. Sundstol, 1999. The effect of variety on maize grain and crop residue yield and nutritive value of the stover. *Animal Feed Science and Technology*. 79: 165-177.
- Agegnehu, G., Ghizam, A. and Sinebo, W. 2006. Yield Performance and Land-Use Efficiency of Barley and Faba Bean Mixed Cropping in Ethiopian Highlands. *European Journal of Agronomy*, 25, 202-207.
- Aklilu Mekasha, Ashebre Tegegn, Abraham Abera and Mandefro Nigussie, 2007. Compatibility and performance of different annual forage legumes under-sown to different maize varieties. *Pastoral livestock systems: opportunities and challenges as a livelihood strategy*. Proceedings of the 15th annual conference of the Ethiopian Society of Animal Production, Addis Ababa, Ethiopia.
- Anil, L., Park, J., Phipps, R.H. and Miller, F.A. 1998. Temperate Intercropping of Cereals for Forage: A Review of the Potential for Growth and Utilization with Particular Reference to the UK. *Grass and Forage Science*, 53, 301-317. <https://doi.org/10.1046/j.1365-2494.1998.00144.x>
- Banik, P., Midya, A., Sarkar, B.K. and Ghose, S.S. 2006. Wheat and Chickpea Intercropping Systems in an Additive Series Experiment: Advantages and Weed Smothering. *European Journal of Agronomy*, 24, 325-332.
- Banik, P., Sasmal, T., Ghosal, P.K. and Bagchi, D.K. 2000. Evaluation of Mustard (*Brassica campestris* var. Toria) and Legume in 1:1 and 2:1 Replacement Series System. *Journal of Agronomy and Crop Science*, 185, 9-14. <https://doi.org/10.1046/j.1439-037X.2000.00388.x>
- Chen, C., Westcott, M., Neill, K., Wichman, D. and Knox, M. 2004. Row Configuration and Nitrogen Application for Barley-Pea Intercropping in Montana. *Agronomy Journal*, 96, 1730-1738. <https://doi.org/10.2134/agronj2004.1730>
- Dhima, K.V., Lithourgidis, A.A., Vasilakoglou, I.B. and Dordas, C.A. 2007. Competition Indices of Common Vetch and Cereal Intercrops in Two Seeding Ratio. *Field Crops Research*, 100, 249-256.
- Diriba Geleti and Lemma Gizachew 2003. Integration of forage legumes in to maize based cropping systems in Western Ethiopia: Effect of intercropping of *Lablab purpureus* and *Vicia atropurpurea* on maize grain and total forage yields. Challenges and Opportunities of Livestock Marketing in Ethiopia. Yilma Jobre and Getachew Gebru (Eds). In: Proceedings of 10th Annual conference of the Ethiopian Society of Animal Production (ESAP) held in Addis Ababa, Ethiopia, August 22-24, 2002. ESAP, Addis Ababa. 407pp
- Fenandez-Aparicio, M., Josefina, C. and Sillero, D.R. 2007. Intercropping with Cereals Reduces Infection by *Orobanche crenata* in Legumes. *Crop Protection*, 26, 1166-1172.
- Ghosh, P.K. 2004 Growth, Yield, Competition and Economics of Groundnut/Cereal Fodder Intercropping Systems in the Semi-Arid Tropics of India. *Field Crops Research*, 88, 227-237.
- Giller, K.E. 2001. Nitrogen fixation in tropical cropping systems, 2nd ed. Wallingford, UK: CAB International.
- Hauggaard-Nielsen, H., Ambus, P. and Jensen, E.S. 2001. Evaluating Pea and Barley Cultivars for Complementary in Intercropping at Different Levels of Soil N Availability. *Field Crops Research*, 72, 185-196. [https://doi.org/10.1016/S0378-4290\(01\)00176-9](https://doi.org/10.1016/S0378-4290(01)00176-9)
- Mergia Abera 2014. The Effect of Under Sowing of Forage Legumes in Maize on Dry Matter Yield and Nutritional Value of the Fodder in Baresa Watershed, Ethiopia. *International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064*. Volume 3 Issue 8, August 2014.

- Mpairwe, D.R., Sabiitti, E.N., Ummina, N.N., Tegene, A. and Osuji, P., 2002. Effect of intercropping cereal crops 17. with forage legumes and source of nutrients on cereal grain yield and fodder dry matter yields. *African Crop Science Journal* 10(1): 81-97.
- Peters, M., and E.C. Lascano 2003. Forage technology adoption: linking on-station research with participatory methods. *Tropical Grasslands* 37: 197-203.
- Poggio, S.L. 2005. Structure of Weed Communities Occurring in Monoculture and Intercropping of Field Pea and Barley. *Agriculture, Ecosystems & Environment*, 109, 48-58.
- Tolera Adugna, Yami Alemu, Alemu Dawit 2012. Livestock feed resources in Ethiopia: Challenges, Opportunities and the need for transformation. Ethiopia Animal Feed Industry Association, Addis Ababa, Ethiopia.
- Tsubo, M., Walker, S. and Ogindo, H.O. 2005. A Simulation Model of Cereal-Legume Intercropping Systems for Semi-Arid Regions. II. Model Application. *Field Crops Research*, 93, 23-33.