

Impact of Different Nitrogen Amount and Seeding Rates on Growth and Yield of NERICA 4 Rice Varieties in Upland Conditions

Bayissa Gedefa¹ Hisashi Urayama² Yoshie Yageta²
1.Bako Agricultural Research Center, P.O.Box, 03, Bako, Ethiopia
2.JICA Tsukuba International Center 3-6 Koyadai, Tsukuba-shi, Ibaraki-ken, Japan
*Corresponding author's email: bayissagedefa@gmail.com

Abstract

With the recent breakthrough in the development of the New Rice for Africa (NERICA) for the upland production system and the high demand for rice, farmers' interest in growing upland rice has increased. A field experiment was conducted at Tsukuba International center, Japan from May to October 2019 to evaluate the response of rice to seed rate and nitrogen fertilizer application and also determine economically optimum seed rate and nitrogen level for NERICA-4 upland variety. The experiment contained factorial combination of four levels of nitrogen (0, 30, 60, 90 kg ha⁻¹ and two seed rates (60& 80 Kg ha⁻¹). The experiment was laid out in randomized complete block design plot arrangement with three replications. The results from this study indicated that seed rate showed no significant effect on total dry matter weight, plant length, SPAD value, Leaf area index, panicle number, spikelet number, ripening ratio and grain yield while the total number of tillers was significant. However, nitrogen fertilizer rates significantly ($p < 0.05$) affected in plant length, tiller number, SPAD value, dry matter and Leaf area index. Except 1000grain, all the yield and yield component parameters have also shown a significant increase up to the rate of 90 kg N ha⁻¹. The highest grain yield (3340 kg·ha⁻¹) was recorded using 90 kg N ha⁻¹ followed by 60 kg N ha⁻¹ (3200 kg ha⁻¹) and the lowest yield (2100 kg ha⁻¹) was recorded from control. Marginal rate of return analysis is also showed that the treatment with N fertilizer rate of 90 kg N ha⁻¹ and seed rate 60 kg ha⁻¹ gave the highest Marginal rate of return 1217 % followed by the treatment with N fertilizer rate of 60 kg N ha⁻¹ and seed rate 60 kg ha⁻¹. The general trend of the data indicated that the fertilizer rate has a greatest impact on grain yield compared to the seed rate of rice crop. Based on this study using 60 kg/ha seed rate and application of 90 kg N ha⁻¹ gives high grain yield for rice crop production at experimental site.

Keywords: Economic profit, Seed rate, N fertilizer rate, Optimum grain yield, Marginal rate of return

DOI: 10.7176/JBAH/12-9-04

Publication date: May 31st 2022

Introduction

Rice is a staple food for more than half of the world population. It is not only a main source of calories but also an important source of income and employment (Zhao *et al.*, 2011). With the recent breakthrough in the development of the New Rice for Africa (NERICA) for upland production system and the high demand for rice, farmers' interest in growing upland rice has increased. In Ethiopia, rice has become one of the most important crops whereby its production and area coverage increase every year (Tesfaye *et al.*, 2005).

Despite its considerable importance, yield of the crop is very low especially in case of upland rice as farmers have limited access to production technologies. High yields are indeed possible under upland conditions. Under ideal conditions on experiment stations, yields of 7.0 and 7.2t/ha have been reported in Philippines and Peru respectively (IRRI, 1986). It is also indicted that a quantity as high as 5.4t/ha was reported in Nigeria (Abifarin *et al.*, 1972). Appropriate agronomic packages coupled with adequate nutrient supply at an appropriate time play are pivotal role in expression of yielding potential of crops. Among these, insufficient nitrogen fertilizers and inappropriate seed rats are important and research on these limiting factors will surely lead to high yields. Therefore, to increase the yield and yield components of the crop, it is necessary to apply Optimum seed rate and nitrogen rate which are economically feasible to the farmers.

Nitrogen is the most essential element that is applied most frequently and with high amount in rice production. High-yielding rice cultivars needs large amount of nitrogen to achieve acceptable grain yields. But application of nitrogen fertilizer either in excess or less than optimum rate affects both yield and quality of rice to remarkable extent (Shukla *et al.*, 2015).

Seed rate is among the crucial factors that limit yield of rice and the most important agronomic aspect which need due attention. Profitability and productivity of rice can be improved through use of optimum seeding rate at economically and agronomic ally viable levels. When the plant density exceeds an optimum level competition among plants for light above ground and nutrients below ground becomes sever (Bloch et al, 2002). Consequently, plant growth slows down and the grain yield decreases. However, very low plant density may not

enable to attain the yield plateau (Hay and Wolker, 1989).

However, in Ethiopia and particularly with farmers in western Oromia region, there still remains an information gap with regard to knowledge on Optimum seed and nitrogen rate to be used in upland rice cultivation. Therefore, the objectives of this experiment are to determine appropriate seed rate and nitrogen level and evaluate the economics of various seed rate and nitrogen level for optimum grain yield of rice in upland condition.

Material and Methods

Soil property of experimental site

For soil analysis, before planting soil samples were randomly taken from the experimental site at a depth of 15cm using an auger and the samples were mixed thoroughly to produce one representative composite sample. The soil samples were air-dried and ground to pass 2 and 0.5 mm (for total N) sieves. Then it was analyzed at the soil laboratory for pH, total carbon, nitrogen, available P and exchangeable K by following standard procedure. The soil chemical properties shown in table1, has slightly acidic nature with optimum total nitrogen content.

Table1. Analytical result of Soil properties of the experimental field

No	Properties	Values
1.	pH (H ₂ O)	6.2
2.	Total carbon (g kg ⁻¹)	35.5
3.	Total Nitrogen (g kg ⁻¹)	3.22
4.	Available P(Bray II, mg kg ⁻¹)	68.7
5.	Exchangeable K (cmolc kg ⁻¹)	2.09

Climatic conditions of experimental site

Climatic data was collected throughout the growing season using digital meteorological instruments at Tsukuba International center weather station. Monthly rainfall in (mm), temperature in (deg.C) and solar radiation (MJ/m²) are presented in (Fig1.).




Fig.1. Temperature, precipitation and Solar Radiation of the experimental site

Treatments were comprised of two factors: namely two seed rate levels (60, and 80, kg ha⁻¹), and four levels of nitrogen (0, 30, 60, and 90 kg N ha⁻¹). The experiment were arranged in a randomized completed block design with eight treatment and three replications. The plot size was 3mx3m and space between plots 0.5m whereas 0.5m space between blocks was left. The rice variety called NERICA4 (which have high yielding and medium maturity) was used. Seed were selected using salt solution of 1.06specific gravity. 100 seeds of this variety were counted, soaked in water for 24 hours and then incubated at 30 °C for 72 hours to test the viability. To prevent fungal diseases the seed were disinfected with Tough block (Talaromyces flavns powder hydrating agent) at

the rate of 200 time solution and after short drying the seeds were coated with (Thiram) as bird repellent (1% of seed weight). Planting was done manually by drilling in to furrows of 2-3cm depth the seeds at row spacing of 45 cm. 2-3 days after sowing Pre emergency herbicide pendemetheline (GO-GO-SAN) was applied for early weed control, second and third weeding were done manually 45 days after emergency.

Table2. Treatments combination

Seeding rate (kg ha ⁻¹)	Factors	Treatments combination
	Nitrogen amount ((kg ha ⁻¹)	
60 (60S)	0 (N0)	S60N0
	30 (N30)	S60N30
	60 (N60)	S60N60
	90 (N90)	S60N90
80(80S)	0 (N0)	S80N0
	30 (N30)	S80N30
	60 (N60)	S80N60
	90 (N90)	S80N90

Fertilizer application rate and timing

For all the plots, 100 of P₂O₅ and 80 kg ha⁻¹ of K₂O were applied as P and K sources respectively. All phosphorous and 75 % of the potassium were applied at basal whereas applied the rest of the potassium fertilizer was applied at 50 days before heading. Nitrogen was applied two times as basal and top dressing by using Ammonium Sulphate. As basal fertilizer (2/3 of total amount) and top dressing applies 50 days after sowing (DAS) 1/3 of total amount.

Table 3.Application time and amount of chemical fertilizer

Nutrient	Basal kg ha ⁻¹ (or gm ⁻²)	Top dressing* (kg ha ⁻¹ or gm ⁻²)	Total (kg ha ⁻¹ or gm ⁻²)
N30	20 (114)	10 (57)	30(171)
N60	40 (228)	20(114)	60(342)
N90	60 (342)	30(171)	90(513)
P ₂ O ₅	100 (686)	0(0)	100(686)
K ₂ O	60 (118)	20(79)	80(197)

*Top dressing: 50DAS (days after sowing).

Measurement of growth parameters

Investigations of growth parameters were recorded five times during the growth period within time interval of fifteen days (15 days) apart. The first observation was carried out at early tillering stage, at active tillering, maximum tillering, and panicle initiation and at heading stages. Five sampling areas of 50cm each were randomly selected in each plot from which growth parameters data were observed and were excluded in border sampling to avoid border effect. The growth parameters were investigated on plant height, tiller number, SPAD value, Leaf area index (LAI) at 35 Days After Sowing(DAS), 50 DAS, 65DAS, 80 DAS and dry matter weight at panicle initiation and at heading time.

Yield and yield components

An area of 1m² was used for partial harvesting to determine yield and yield components. Harvesting was done 30 days after heading (DAH) and number of panicles were counted and recorded. After drying the panicles for one week in green house bringing the moisture content to 14 %, threshing of grains had followed and % of ripened grain was determined by separating filled grain from empty grains by using water of specific gravity 1.0. Then is followed determination of 1000 grain weight. Lastly spikelet number per panicle was also determined by calculating the total numbers of grains both filled and empty divide grains which were harvested per plot. All these yield components were used to calculate for the final yield per square meter which was subsequently converted to corresponding yield in kilogram per hectare.

Data analysis

The data collected on different parameters were statistically analyzed using **Stat Cel 3** program. After analysis ANOVA the differences between the treatment means were compared by using the result of multiple comparisons method of Tukey Kramer test at significance (P5% level of significance. To identify the economic optimum rate of nitrogen fertilizer and seeding rate, economic analysis was also done using the CIMMYT partial budget analysis methodology (CIMMYT, 1988).

Result and Discussion

Plant Length

Data on plant length was collected at 35 days, 50Days, 65 days, 80days after sowing and heading time was recorded. Fig 2 showed plant length increased gradually in all treatments up to the final observation on heading time. The analysis of variance at $p < 0.05$ showed that there were significant difference among the four nitrogen amount applied at all sampling dates. The highest plant height was observed in the higher nitrogen level (N90 kg ha⁻¹ and N60 kg ha⁻¹) respectively. The lowest plant height was recorded from control treatment (zero nitrogen level) at all sampling dates. However, there were non-significant differences in plant height between seed rate (S60 and S80 kg ha⁻¹) at all growth stage. The general trend of the data indicated that the fertilizer rate has a greatest impact on plant length compared to the seed rate of rice crop. There was also no interaction effect observed between seed rate and nitrogen level. Two seed rate showed almost similar length plant of at 35 DAS, 50 DA 65DAS, 80DAS and heading time.




Fig.2. Effect of nitrogen level and Seed rate on plant length.

Number of tiller

Number of tillers per m² of rice increased over time by gradual elevation of nitrogen fertilizer up to 80 days of after planting afterwards showed a falling trend. Data presented in Fig.3. Indicated that seed rate had significant ($P < 0.05$) effect on tillers m⁻². It is evident from the data that maximum tillers (69.85m⁻²) were noted when plots were seeded with 80 kg ha⁻¹ (S80), while minimum tillers (59.50 m⁻²) were recorded when 60 kg ha⁻¹ seed rate was used (S60). The tillering potential was much higher at higher seed rate, when compared to lower seed rate. It might be due to that the number of tillers m⁻² increased with increase in seeding rate,

The data further showed that different nitrogen levels had significantly ($P < 0.01$) affected number of tillers m⁻². The highest number of tiller was recorded in the higher nitrogen level N90 (90 kg N ha⁻¹) while the lowest number of tiller per m² was recorded with no nitrogen fertilizer (N0). This might be due to increased nitrogen rate; therefore it is thought it could be led to greater stimulation of vegetative growth. All fertilizer levels significantly affected at vegetative and reproductive growth of the plant depending upon the availability of needed nutrition which leads to proportional increase in tillers m⁻². These results are also in conformity with Ayoub *et al.*, (1994), who reported that 120 kg N ha⁻¹ gave significantly the highest tillers m⁻². However the interaction effect of the seed rate and amount of nitrogen was not significant.




Fig.3. Effect of nitrogen level and Seed rate on tiller number.

SPAD value

SPAD value m⁻² was significantly ($P < 0.01$) affected by different nitrogen levels. It is clear from the data that maximum SPAD value (50.1 m⁻²) was recorded for plots treated with 90 kg N ha⁻¹ (N90), followed by plots in which 60 kg N ha⁻¹ (N60) was applied which was statistically similar to N30 (45.1 SPAD value m⁻²), while minimum (43.1) SPAD value m⁻² were recorded from control plots (N0). Generally, SPAD value increased with growth of young leaves and chlorophyll formation at early growth stage. However, analysis of ANOVA data presented in (Fig4.) indicated that SPAD value m⁻² were not significantly ($P < 0.05$) affected by different seed rate. The data also indicated that no interaction between seed rate and nitrogen levels at ($P < 0.05$). SPAD value indicted chlorophyll content of NERICA 4 was as shown below.




Fig.4. Effect of nitrogen level and Seed rate on Leaf SPAD Value.

Leaf area index

Leaf area index was investigated at panicle differentiation and heading stage. At both panicle differentiation

stage and heading stage there were no significant difference on seed rate and interaction effect of seed rate and application of nitrogen level. However there was significant difference on different level of nitrogen application at $P < 0.05$ (Table4). Nitrogen fertilizer (90 kg N ha^{-1}) was gave the highest leaf area index. These may attributed with number of tiller in a given area. Similar trends were shown with nitrogen level (90 kg N ha^{-1}).

Table4. Leaf Area Index at Panicle differentiation and heading stage

Treatments		Rice Growth stage	
		Panicle initiation	Heading time
Seed rate	S60	2.6c	2.98c
	S80	2.5c	2.97c
	Tukey(HSD)-5% Significance	ns	ns
Nitrogen _ Level	N0	2.38b	2.77b
	N30	2.5a	2.96ab
	N60	2.65a	3.06ab
	N90	2.65a	3.1a
	Tukey (HSD) - (5%)	**	*
S60	N0	2.36c	2.7ab
	N30	2.50ab	3.00ab
	N60	2.67b	3.10b
	N90	2.70a	3.10a
	S80	2.40c	2.83ab
S80	N0	2.50c	2.93ab
	N30	2.63c	3.03ab
	N60	2.60c	3.1ab
	N90	2.60c	3.1ab
Interaction	Seed X Nitrogen	ns	ns
CV (%)		5.5	6.2

In each column, common letters indicates non-significant difference by Turkeys' Honestly Significant difference (HSD) at 5% level; ns non-significant, * significant at the 0.05 level; ** significant at the 0.01 level

Dry weight

Dry weight production by rice plants increased progressively with the advancement of growth stages and reached rice peak at maturity (Table5). Dry weight was affected by the different level of nitrogen application at both panicle initiation stage and heading stage. 90 kg ha^{-1} of nitrogen level has produced significantly higher dry weight than control (0 kg ha^{-1}). However, dry weight was not significantly influenced by the seed rate at panicle and heading stage. Different seed rate application on (other hand) showed similar stem and leaf dry weight (table5). Interaction effect of the seed rate and nitrogen level was not significant.

Table5. Dry weight at Panicle differentiation and heading stage

Treatments	Rice growth stage		
		total dry weight (g m ⁻²) at Panicle initiation	Dry weight(gm ⁻²) at Heading stage
Seed rate	S60	133.09d	224.09c
	S80	135.86d	224.23c
	Tukey(HSD)-5% Significance	ns	ns
Nitrogen _ Level	N0	125.11b	215.28ab
	N30	131.80c	224.8b
	N60	136.28a	227.45b
	N90	144.75a	229.13a
	Tukey (HSD) - (5%)	**	*
S60	N0	124.06bc	216.06c
	N30	131.03c	223.7b
	N60	133.26ab	229.93a
	N90	144.03b	226.7bc
S80	N0	126.17	214.5bc
	N30	132.57c	225.9bc
	N60	139.3b	224.967bc
	N90	145.45a	231.57bc
Interaction	Seed X Nitrogen	ns	ns
CV (%)		6.05	2.9

In each column, common letters indicates non-significant difference by Turkeys' Honestly Significant difference (HSD) at 5% level; ns non-significant, * significant at the 0.05 level; ** significant at the 0.01 level

Yield and Yield components

Number of Panicle m⁻²

Nitrogen levels had significant effect on the number of panicles per meter square of rice (Table 6). The number of panicle per m² increased with increased level of nitrogen. The highest number of panicle per m² (248.67) was recorded when 90 N kg ha⁻¹ was applied while the lowest (157.50) from 0Nkg ha⁻¹ (control treatment). The number of effective tillers rather than total number of tillers contributes more to enhance productivity of rice plant. The lower of tiller number in present study was attributed to the failure in competition for lower level of nitrogen and aggravated death of the tillers due to mutual shading. (Fageria,, *et al*, 1997). Another explanation is that, competition for assimilates exists between developing panicles and young tillers during the beginning of panicle development causing suppression of growth of many young tillers. However number of panicles per m² was not affected by seeding rate.

Spikelet number Panicle⁻¹

Number of spikelet per panicle varied significantly due to different levels of applied nitrogen. Number of spikelet per panicle increased with the increase of nitrogen levels. The highest number of spikelet per panicle was recorded (86.89) at 90N kg ha⁻¹ While, the lowest number of spikelet (71.20) per panicle of rice was observed at 0 N kg ha⁻¹. Application of nitrogen fertilizer improved grain number of rice. Higher number of grains per panicle at higher nitrogen rate could be due to higher nitrogen absorption which favored formation of higher number of branches per panicle. However, number of spikelet per m² was not affected by seeding rate. The data also indicated that no interaction between seed rate and nitrogen levels at (P < 0.05).

1000 grain Weight

Amount of nitrogen levels and seed rate had no significant effect on the 1000 grain weight per meter square of rice (Table 6.). However, the maximum thousand grain weights were observed at 90N kg ha⁻¹(26.43g) which was statistically similar with 60N kg ha⁻¹. The lowest thousand grain weights was produced with 0 kg N ha⁻¹ (23.7g) which was similar to that of 0 and 30 N kg ha⁻¹. In case of thousand grain weight, the variation is very low among the treatments as it known to be by the genetically controlled character. Similar results were found by other scientists (Maske, N.S., 1997) with nitrogen fertilizer management and concluded that there was little opportunity to improve grain size through agronomic management.

Paddy Yield in kg ha⁻¹

Data indicated that significant difference of (P< 0.05) grain yield was recorded only nitrogen fertilizer rate

however, the seed rate and the interaction of nitrogen fertilizer and seed rate were not significant to alter grain yield (Table6). The highest grain yield (3340 kg ha⁻¹) was recorded using 90N kg ha⁻¹ followed by 60 N kg ha⁻¹ (3200 kg ha⁻¹) and the lowest yield (2100 kg ha⁻¹) was recorded from control (Table 6). Overall paddy yields obtained in this experiment were generally lower than the potential of the variety. A.E, *Et al* (2008) states the potential yields of NERICA4 was 5 t/ha under upland conditions. This low yields obtained might have been caused by unfavorable weather conditions (drought and relatively high temperature, typhoon) experienced during booting, heading and ripening stage. Yoshida (1981) reported that unfavorable weather conditions during heading and ripening may hampered continued growth and development of unfilled spikelet. Grain yield of rice plant is highly relying on the number of spike-bearing tillers produced by each plant, filled grains and grains weigh (Huang,*et al* 2011) .In this experiment, it is considered that at higher nitrogen level showed efficient of nitrogen for good assimilation for translocation and dry matter production of sink.

Table6. Effects of nitrogen level and seeding rate on yield and yield components

Treatments		Panicle number m ⁻²	Spikelet number Panicle ⁻¹	Percentage of ripened grains (%)	1000 grain weight (g)	Paddy Yield (kg ha ⁻¹)
Nitrogen-Level	N0	157.50ab	71.20ab	68.8b	24.58a	2150b
	N30	164.04b	71.73b	66.4b	24.91a	2340ab
	N60	244.33a	80.45a	73.7a	25.00a	3210a
	N90	248.67a	86.89a	75.6a	25.00a	3341a
Tukey Significance	(HSD)-5%	**	**	*	ns	*
Seed rate	S60	205.58c	76.744c	70.90ab	24.90b	2772c
	S80	201.83c	78.32c	71.30ab	24.80b	2750c
Tukey Significance	(HSD)-5%	ns	ns	ns	ns	ns
S60	N0	161.3d	77.17c	68.6c	24.67a	2105d
	N30	171.7d	85.01b	68c	25.00a	2479d
	N60	248c	72.95a	71.7ab	25.00a	3231bc
	N90	241.3c	71.85c	75.5ab	25.00a	3265c
S80	N0	153.7ab	84.6c	69b	24.50a	2201ab
	N30	157b	87.62c	647a	24.83a	2205b
	N60	240.7a	70.51c	75.7a	25.00a	3203a
	N90	256a	70.55c	0.758	25.00a	3420a
Interaction effect	Seed X nitrogen level	ns	ns	ns	ns	ns
CV (%)		4.2	4.8	11.2	12.2	5.8

In each column, common letters indicates non-significant difference by Turkeys' Honestly Significant difference (HSD) at 5% level; ns non-significant, * significant at the 0.05 level; ** significant at the 0.01 level.

Economic evaluation

Table 7 showed a partial budget for this seed rate and nitrogen level experiment. There were eight columns, representing the eight treatments (factorial combination of seed rate and nitrogen level). The first line of the budget presents the average yield from pooled blocks in the recommendation domain for each of the eight treatments. The final line of the partial budget shows the net benefits. This was calculated by subtracting the total costs that vary from the gross field benefits. All costs and benefits were calculated on hectare basis.

The gross benefit was calculated from price (seasonal average) of brown rice grain in the experiment area (4000Yen per 60kg). Variable cost was calculated from the costs involved for purchase of fertilizer and seed. Ammonium Sulphate, which was used as the source of nitrogen, was bought for (1500 Yen per 20kg). For purchase of seed at planting 610 Yen per 4kg was calculated and converted into hectare.

The partial budget analysis of fertilizer level and seed rates revealed that the maximum net benefit was attained from application of 90 N kg ha⁻¹ and 60 kg ha⁻¹ seed while the least net benefit was obtained from the unfertilized treatment (Table 4). The dominance analysis showed that the treatment with the highest fertilizer rate (90 N kg ha⁻¹) and seed rate (80kg ha⁻¹) were cost dominated; i.e., it provided gross margin that was less than that of the preceding treatment. Therefore, it was omitted from the analysis of marginal rate of return (MRR). The MRR analysis showed that the treatment with N fertilizer at the rate of 90 N kg ha⁻¹ and seed rate

60 kg ha⁻¹ gave the highest marginal rate of return 1217.01 % followed by the treatment with N fertilizer rate of 60 kg N ha⁻¹ and seed rates 60kg/ha (Table 4). The treatment with fertilizer rate of 30 N kg ha⁻¹ and seed rate 80kg ha⁻¹ gave marginal rate of return below 100 %, which indicated that this rate is not economically optimum level.

Table 7. Shows a partial budget for this seed rate and nitrogen level experiment.

Item & its price	Treatments							
	S60,N0	S60,N30	S60,N60	S60,N90	S80,N0	S80,N30	S80,N60	S80,N90
Nitrogen rate (Kg ha ⁻¹)	0	30	60	90	0	30	60	90
Seed rate(Kg ha ⁻¹)	60	60	60	60	80	80	80	80
Paddy Yield (kg ha ⁻¹)	2105	2479	3231	3265	2201	2205	3203	3420
Brown rice (Kg ha ⁻¹)	1684	1983.2	2584.8	2612	1760.8	1764	2562.4	2736
Gross field Benefits	112,266.7	132,213.30	172320	174133.3	117386.7	117600	170826	182400
Cost of fertilizer (yen ha ⁻¹)	0	2250	4500	6750	0	2250	4500	6750
Cost of seed (Yen ha ⁻¹)	9150	9150	9150	9150	12200	12200	12200	12200
Total cost that vary among treatment (Yen ha ⁻¹)	9150	11400	13650	15900	12200	14450	16700	18950
Net Benefit	103116	120813	158670	185233	105186	103150	154126	163450
Marginal rate of Return (MRR %)	0	786.01	1234.00	1217.00	67.12	0.6	676.00	615.65

The marginal rate of return (MRR) indicates what farmers can expect to gain on the average, in return for their investment when they decide to change from one practice (or set of practices) to another.

Conclusion

From the results it may be concluded that Investigated growth and yield component variables (plant height, SPAD value, tiller number, leaf area index, spike per panicle, number of fertile tiller per plant, ripening percentage, grain yield and dry matter) had significantly influenced by different levels of nitrogen fertilizer but seed rate significantly alters only of the tasted variable (number of fertile tiller per plant). The highest grain yield (3340 kg ha⁻¹) was recorded using 90 N kg ha⁻¹ followed by 60 kg N ha⁻¹ (3200 kg ha⁻¹) and the lowest yield (2100 kg ha⁻¹) was recorded from control (Table 6). The general trend of the data indicated that the fertilizer rate has a greatest impact on grain yield compared to the seed rate of rice crop. In other word NERICA 4 is more response to different nitrogen level than seed rate Based on this experiment using 60 kg ha⁻¹ seed rate and application of 90N kg ha⁻¹ gave high grain yield for rice crop production at experimental site.

Acknowledgement

First, I would like to thank the almighty God for he gave me health, strength and patience to withstand the inconveniences, which I came across through all the lines of life. I would like to express my sincere gratitude to the Japanese government through Japan International Cooperation Agency (JICA) for given me the opportunity to participate in this training program and support rendered to this training course. I would like also to extend my profound gratitude to JICA Tsukuba (TBIC) administration and all the staffs for their constant assistance provided throughout the entire period of our training in Japan. Special thanks go to program officer Ms. Miki Motumora for all her tireless administrative work. I would like to express my heart felt gratitude to my Instructor Dr. Hisashi Urayama and Ms. Yoshie Yageta for diligent effort & valuable guidance throughout this training course.

Last but not the least I would like to express my sincere appreciation and gratitude to my technical advisors for their giving me advice guidance, academic insights and shared experiences that made our course successful. My special recognition goes to my family, relatives and fellow participants for being with me all the time

providing me a moral support and encouragement.

References

- Abifarin. A.O.R. Chabron, M. Jacquot, R. Marie, and J.C. Moomaw. 1972. Upland improvement in West. P 625-635 in International Rice Research Institute, Rice Breeding. Los Baiios, Philippines.
- Ayoub, M., S. Quertin, S. Lussier and D.L. Smith. 1994. Timing and levels of nitrogen fertility effects on wheat yield in Eastern Canada. *Crop Sci.* 34: 748-756.
- Baloch, A.W., Soomro, A.M., Javed, M.A. and Ahmed, M. 2002. Optimum plant density for high yield in rice. *Asian Journal of Plant Science*, 1: 25–27.
- CIMMYT (1988). Agronomic data to farmers' recommendations: an economics training manual. Completely revised Edition. Mexico. 27-18-6.
- Fageria NK, Baligar VC, Jones CA 199, Growth and mineral nutrition of field crops. Marcel Dekker Inc., New York.
- Huang, M., Zou, Y.B., Jiang, P., Bing, X.I.A., Md, I. and Ao, H.J. (2011) Relationship between Grain Yield and Yield Components in Super Hybrid Rice. *Agricultural Sciences in China*, **10**, 1537-1544. [http://dx.doi.org/10.1016/S1671-2927\(11\)60149-1](http://dx.doi.org/10.1016/S1671-2927(11)60149-1)
- IRRI (International Rice Research Institute). 1986. Upland Rice: A Global Perspective. (Eds. Gupta and Toole). Los Banos, Laguna, Philippines.
- Maske, N.S., Borkar, S.L. and Rajgire, H.J. Effects of nitrogen levels on growth, yield and grain quality of rice. *J. Soils and Crops*. 1997, 7(1): 83-86.
- Shukla, V.K., Tiwar, R.K., Malviya, D.K., Singh, S.K. and Ram, U.S. 2015. Performance of rice varieties in relation to nitrogen levels under irrigation condition. *African Journal of Agricultural Research*, 10(12): 1517-1520.
- Tesfaye Z, Befekadu A, Aklilu A ,2005. Rice production, consumption and marketing: The case of Fogera, Dera and Libokemkem Districts of Amhara Region. Paper presented on the National workshop on the status of rice research and promotion in Ethiopia. 3-4 June 2005, Bahir Dar, Ethiopia.
- Yoshida, S. 1981. Fundamentals of Rice Crop Science. The International Rice Research Institute. P69-146.
- Zhao, L., Wu, L., Wu, M. and Li, Y. 2011. Nutrient uptake and water use efficiency as affected by Modified rice cultivation methods with irrigation. *Paddy Water Environment*, 9: 25-32.