

In-vitro Competition Bio-assay Experiment on the Effect of *Trichoderma* Species and Some Crop Pathogenic Fungi

Timothy Ipoola Olabiyi^{1*} and Michelina Ruocco²

1. Department of Crop and Environmental Protection, Ladoké Akintola University of Technology, P.M.B 4000, Ogbomosho, Nigeria.
2. Istituto per la Protezione delle Piante, CNR, Via Università 130, Portici (NA), 80055, Italy.

*E-mail of the corresponding author: t.olabiyi@yahoo.co.uk; tiolabiyi@lautech.edu.ng

The lead author is very grateful to Institute of Plant Protection of the Italian National Research Council, Italy that funded this research through Year 2009 Endure Crop Protection award.

Abstract

Fungi of the genus *Trichoderma* have a track record of being antagonist to quite of a number of agricultural important pathogens. *Trichoderma* have some unique characteristics that make it scientifically proven and suitable bio-control agents against varieties of pathogenic organism infecting economic food crops. *Trichoderma* has the advantage of being environment friendly and not hazardous to the health of human beings, livestock, soil and environment. Competitive bio-assay experiment was carried out in the laboratory on the effects of *Trichoderma* species (*T. atroviride* P1 isolates, *T. harzianum* T22 isolates, *T. viride*) on some crop pathogens (*Phytophthora cinnanerium*, *Botrytis cinaria* and *Rhizoctonia solani*). Pure culture of *Trichoderma* and pathogenic fungi were replicated four times and arranged in a complete block design. The result of the experiment shows that *Trichoderma* species are strong competitor of *P. cinnanerium*, *B. cinaria* and *R. solani*. Within 72 hours, the *Trichoderma* species were able to grow and completely overlap the *P. cinnanerium*, *B. cinaria* and *R. solani*. This strong competitiveness indicated that *Trichoderma* species would effectively inhibit the growth of *P. cinnanerium*, *B. cinaria* and *R. solani* on the infected crop; thus the application of *Trichoderma* species in the control of *P. cinnanerium*, *B. cinaria* and *R. solani* infected crops.

Keywords: *Trichoderma*, bio-control, *Phytophthora cinnanerium*, *Botrytis cinaria*, *Rhizoctonia solani*, pathogens, fungi.

1. Introduction

Biological control of disease/ pathogen is the application of natural enemies in the control/ eradication of the pathogen population. Biological control is an environmentally friendly, scientifically proven and effective means of mitigating pathogens or pests through the use of natural enemies. A world estimated loss due to crop diseases was up to 12%, while a loss due to post-harvest food spoilage was between 10 and 50%. Effective control of crop losses due to pests (micro-organism, insect and weed) therefore holds the keys for steady and stable food supply of the world. Amongst all effective and recommended controls of the crop pests, biological control holds a great promise for the future. Basically, biological control has the advantages of being environmentally friendly and not hazardous to the health of human beings, livestock and wildlife; especially now that the whole world is clamoring for IPM methods of pest control (Lorito *et al*, 2006; Woo *et al*, 2006; Olabiyi, 2009).

Fungi of the genus *Trichoderma* have a track record of being antagonist to quite a number of agriculturally important pests. It had been most effective bio-pesticides applied for crop protection since the era of traditional farming and nascent organic agriculture. *Trichoderma* have some unique characteristics that make it scientifically proven and suitable bio-control agents against varieties of pathogenic organisms infecting economic food crops. These are: non-toxic to human beings, livestock and wildlife; non-pathogenic organism on crops; compatible with other control methods (physical, chemical, cultural, planting of resistance variety); effective at low concentrations; easy and cheap to culture or produce; could be bottled or prepared in another easily distributable pack; *Trichoderma* is ubiquitous (Lorito, 1998; Olabiyi, 2009). *Trichoderma* is capable of producing secondary metabolites with antibiotic activity. Of recent, *Trichoderma* composted hardwood bark isolates, was reported to

produce a metabolite (Harzianic acid) with antifungal and plant growth promoting activity. *Trichoderma* species have been formulated and used as bio-pesticides, bio-protectants, bio-stimulants and bio-fertilizer on a large variety of crops (Reino *et al*, 2008; Vinale *et al*, 2009).

Objective of this study is to determine in-vitro competition bio-assay between *Trichoderma* species (*Trichoderma harzianum* P 1 isolate, *Trichoderma harzianum* T 22 isolate, *Trichoderma viride*) and pathogenic fungi (*Phytophthora cinnanerium*, *Botrytis cinaria* and *Rhizoctonia solani*).

2. Materials and Methods

2.1 Preparation of Potato Dextrose Agar (PDA)

Dissolve 27g of Potato Dextrose Broth (PDB) and 15g Micro Agar in 1 litre of deionised water in an Erlenmeyer conical flask (2 litre capacity). Sealed properly with cork, autoclaved at 121°C, and 15psi for 20 minutes. Allow the autoclaved media to cool and thereafter pour small quantity (20-25mls) into Petri dish inside the Lamina flow (sterilized condition and working tools). Cover up the Petri dish after solidification process. The media is ready for use or otherwise keep inside the refrigerator for subsequent use.

2.2 Source of *Trichoderma* and pathogenic fungi

Pure culture of *Trichoderma* species and pathogenic fungi used for the study were obtained from Istituto per la Protezione delle Piante, CNR, Via Università 130, Portici (NA), 80055, Italy. The *Trichoderma* species were *T. harzianum* P 1 isolate, *T. harzianum* T 22 isolate and *T. viride*; while the pathogenic fungi were *Phytophthora cinnanerium*, *Botrytis cinarea* and *Rhizoctonia solani*

2.3 Introduction of bio-control agents and pathogens to PDA

Bio-control agents and pathogenic fungi were carefully introduced onto the PDA. There were 15 treatments, replicated 4 times fitted into randomized complete block design. The treatments were *Trichoderma harzianum* P 1 isolate; *Trichoderma harzianum* T 22 ATCC isolate; *Trichoderma viride*; *Phytophthora cinnanerium*; *Botrytis cinarea*; *Rhizoctonia solani*; *Trichoderma harzianum* P 1 isolate and *Phytophthora cinnanerium*; *Trichoderma harzianum* P 1 isolate and *Botrytis cinarea*; *Trichoderma harzianum* P 1 isolate and *Rhizoctonia solani*; *Trichoderma harzianum* T 22 ATCC isolate and *Phytophthora cinnanerium*; *Trichoderma harzianum* T 22 ATCC isolate and *Botrytis cinarea*; *Trichoderma harzianum* T 22 ATCC isolate and *Rhizoctonia solani*; *Trichoderma viride* and *Phytophthora cinnanerium*; *Trichoderma viride* and *Botrytis cinarea*; *Trichoderma viride* and *Rhizoctonia solani*. The experiment was carried out under Lamina flow and immediately after the setting up of the competition bio-assay; they were arranged in the incubator at 25°C for 72 hours. Records of growth of each bio-control and pathogenic organisms were taken at every 24 hours. Picture of each treatment and treatment combinations were also taken at the 72nd hour.

3. Results

The results presented revealed the competitiveness of *Trichoderma* species and pathogenic fungi. Plate 1 elicits the competition between *Trichoderma harzianum* (T 22 and P1 isolates), *T. viride* and *R. solani*. It was evident that *Trichoderma* species inhibit the growth of *R. solani*. During the competition between *Trichoderma* species and *R. solani*, *Trichoderma* species proved to be aggressive competitor over *Rhizoctonia*. *Trichoderma* species grew faster and overlay on the pathogenic fungi (*R. solani*). Similar trend was observed in bio-assay competition between *Trichoderma harzianum* (T 22 and P1 isolates), *T. viride* and *Botrytis cinarea* (Plate 2). *Trichoderma* species grew very fast and then hindered further growth of the pathogenic fungi (*Botrytis cinarea*).




Plate 1: Competition assay between *Trichoderma* and pathogenic fungi (*R. solani*)

Plate 2: Competition assay between *Trichoderma* and pathogenic fungi (*B. cinerea*)

Plate 3 shows the bio-assay competition between *Trichoderma harzianum* (T 22 and P1 isolates), *T. viride* and *P. cinnanerium* in the laboratory. It was evident that *Trichoderma* species grew faster, overlay *P. cinnanerium* and prevented its further growth and development. *Trichoderma* species proved to be aggressive competitor over *P. cinnanerium*. Plate 4 shows the competitive bio-assay between *Trichoderma viride* and pathogenic fungi (*R. solani*, *Botrytis cinarea* and *P. cinnanerium*). It was evident that *T. viride* grew faster to inhibit further growth of the pathogenic fungi (*R. solani*, *Botrytis cinarea* and *P. cinnanerium*)



Plate 3: Competition assay between *Trichoderma* and pathogenic fungi (*P. cinnanerium*)

Plate 4: Competition assay between *T. viride* and pathogenic fungus

Bio-assay competition was observed between *T. harzianum* (P1 isolate) and pathogenic fungi - *R. solani*, *Botrytis cinarea* and *P. cinnanerium* (Plate 5). It was evident that *T. harzianum* (P1 isolate) grew very faster to suppress the growth of the pathogenic fungi. Similar trend was observed when *T. harzianum* (T22 isolate) was introduced to pathogenic fungi - *R. solani*, *Botrytis cinarea* and *P. cinnanerium* (Plate 6). *T. harzianum* (T22 isolate) proved to be an aggressive competitor over *R. solani*, *Botrytis cinarea* and *P. cinnanerium*



Plate 5: Competition assay between *T.harzianum* P1 isolate and pathogenic fungus
 Plate 6: Competition assay between *T. harzianum*T22 ATCC isolate and pathogenic fungus

Table 1 shows the time interval at which *Trichoderma harzianum* (T22 isolate) grew over pathogenic fungi - *R. solani*, *Botrytis cinarea* and *P. cinnanerium*. It was evident that within 72 hours (3 days), *Trichoderma harzianum* (T22 isolate) hindered the growth of *R. solani*, *Botrytis cinarea* and *P. cinnanerium*. Similar trend was observed when *Trichoderma harzianum* (P1 isolate) competed with *R. solani*, *Botrytis cinarea* and *P. cinnanerium* (Table 2).

Table 1: Competition assay between *Trichoderma harzianum* T22 isolate and pathogenic fungi (Figures are in cm)

Time (in hours)	T22 alone	T22 versus <i>Rhizoctonia</i>	T22 versus <i>Botrytis</i>	T22 versus <i>Phytophthora</i>
24	1.7 x 2.0	1.5 x 1.0	1.2 x 1.4	1.0 x 1.5
48	5.5 x 5.5	4.5 x 5.5	4.0 x 5.0	4.0 x 5.0
72	5.5 x 5.5	5.5 x 5.5	5.5 x 5.5	5.5 x 5.5

Table 2: Competition assay between *Trichoderma harzianum* P1 isolate and pathogenic fungi (Figures are in cm)

Time (in hours)	P1 alone	P1 versus <i>Rhizoctonia</i>	P1 versus <i>Botrytis</i>	P1 versus <i>Phytophthora</i>
24	0.9 x 0.9	1.2 x 1.1	1.2 x 1.2	0.9 x 0.9
48	1.8 x 2.0	2.0 x 3.0	2.3 x 2.6	2.0 x 2.4
72	3.0 x 2.8	4.0 x 5.0	3.5 x 4.5	3.5 x 2.8

Table 3 shows the time interval at which *Trichoderma viride* grew over pathogenic fungi - *R. solani*, *Botrytis cinarea* and *P. cinnanerium*. It was evident that within 3 days, *Trichoderma viride* prevented the growth and development of *R. solani*, *Botrytis cinarea* and *P. cinnanerium*. Similar trend was observed when pathogenic fungi (*R. solani*, *Botrytis cinarea* and *P. cinnanerium*) competed with *Trichoderma viride* (Table 4). *Trichoderma viride* was an aggressive competitor on pathogenic fungi.

Table 3: Competition assay between *Trichoderma viride* and pathogenic fungi (Figures are in cm)

Time (in hours)	<i>T. viride</i> alone	<i>T. viride</i> versus <i>Rhizoctonia</i>	<i>T. viride</i> versus <i>Botrytis</i>	<i>T. viride</i> versus <i>Phytophthora</i>
24	2.0 x 1.5	1.6 x 1.5	2.0 x 2.0	2.0 x 2.0
48	3.5 x 5.0	3.5 x 4.7	3.6 x 5.5	4.0 x 5.0
72	5.5 x 5.5	5.5 x 5.5	5.5 x 5.5	5.5 x 5.5

Table 4: Competition assay between *Rhizoctonia* and *Trichoderma* species (Figures are in cm)

Time (in hours)	<i>Rhizoctonia</i>	<i>Rhizoctonia</i> versus T22 isolate	<i>Rhizoctonia</i> versus P1 isolate	<i>Rhizoctonia</i> versus <i>T. Viride</i>
24	0.9 x 0.9	0.9 x 0.9	0.9 x 0.9	0.9 x 0.9
48	0.9 x 0.9	0.9 x 0.9	0.9 x 0.9	0.9 x 0.9
72	1.3 x 1.2	1.0 x 1.0	1.0 x 1.0	1.0 x 1.0

Table 5 elicits the time interval at which *Trichoderma harzianum* T 22 isolate, *Trichoderma harzianum* P1 isolate and *Trichoderma viride* suppressed the growth of the *Botrytis cinarea*. It was evident that within 3 days, *Trichoderma* species prevented the growth and development of *Botrytis cinarea*. Similar trend was observed when *Trichoderma* species competed with pathogenic fungus - *P. cinnanerium*. (Table 6).

Table 5: Competition assay between *Botrytis* and *Trichoderma* species (Figures are in cm)

Time (in hours)	<i>Botrytis</i>	<i>Botrytis</i> versus T22 isolate	<i>Botrytis</i> versus P1 isolate	<i>Botrytis</i> versus <i>T. Viride</i>
24	1.0 x 1.0	0.9 x 0.9	0.9 x 0.9	1.0 x 1.0
48	1.5 x 1.3	1.3 x 1.2	1.5 x 1.6	1.8 x 1.7
72	4.0 x 4.0	3.0 x 2.5	2.5 x 2.8	2.0 x 1.8

Table 6: Competition assay between *Phytophthora* and *Trichoderma* species (Figures are in cm)

Time (in hours)	<i>Phytophthora</i>	<i>Phytophthora</i> versus T22 isolate	<i>Phytophthora</i> versus P1 isolate	<i>Phytophthora</i> versus <i>T. Viride</i>
24	1.2 x 1.0	1.1 x 1.1	1.0 x 1.0	1.0 x 1.0
48	1.8 x 2.0	2.0 x 2.0	2.0 x 2.0	1.8 x 2.0
72	2.4 x 2.2	2.4 x 2.2	2.5 x 2.4	2.2 x 2.1

4. Discussion

The application of bio-control agents and/ or their metabolites for plant diseases control is one of the promising ways to reduce the dependence on chemicals in agriculture, particularly in crop production/ crop protection. In particular, *Trichoderma* are among the most effective bio-control bio-pesticides recommended for plant disease protection against plant diseases under organic agriculture. *Trichoderma* is listed both in Europe and USA as a pesticide permitted for use in organic farming (Woo *et al*, 2006; Olabiyi, 2004).

In recent decades, many bio-control agents have been used in plant protection. However, *Trichoderma* species have been recognized for a long period of time as registered commercial products and biological control agents for the control of plant diseases. Coupled with this, is the potency of *Trichoderma* species to increase plant growth and development (Lorito *et al*, 2006; Woo *et al*, 2006). *Trichoderma* species are known to involve in complex interactions with host plants and soil microbes. The mechanisms involved in the antagonism of *Trichoderma* species on the pathogen were reported to be competition for nutrient, induction of systemic resistance to pathogen, cell wall-lytic enzyme activity, mycoparasitism and antibiosis (Marra *et al*, 2006; Vinale *et al*, 2008; 2004; Lorito, 1998). *Trichoderma* is capable of producing secondary metabolites with antibiotic activity. Of recent, *Trichoderma* composted hardwood bark isolates, was reported to produce a metabolite (Harzianic acid) with antifungal and plant growth promoting activity. *Trichoderma* species have been formulated and used as bio-pesticides, bio-protectants, bio-stimulants and bio-fertilizer on a large variety of crops (Reino *et al*, 2008; Vinale *et al*, 2009). As at today, *Trichoderma*-based bio-fungicides are realities in agriculture, with more than 50 registered formulations worldwide.

References

- Lorito, M; Woo, S; Iaccarino, M & Scala, F. (2006). "In Microrganismi Antagonisti": Iaccarino, M Ed.; Idelson-Gnocchi, s.r.l.: Naples, pp 146-175
- Woo, S.L.; Scala, F.; Ruocco, M. & Lorito, M. (2006). "Phytopathology", 96: 181-185
- Marra, R.; Ambrosino, P; Carbone, V.; Vinale, F.; Woo, S.L.; Ruocco, M.; Ciliento, R.; Lanzuise, S.; Ferraioli, S.; Soriente, I., Gigante, S.; Turra, D.; Fogliano, V.; Scala, F. & Lorito, M. (2006). "Curriculum Genetics" 50: 307-321.
- Vinale, F.; Flematti, G.; Sivasithamparam, K.; Lorito, M.; Marra, R.; Shelton, B.W & Ghisalberti, E.L. (2009). "Harzianic Acids, An antifungal and plant growth promoting metabolite from *Trichoderma harzianum*". *Journal of Natural Products* 72: 2032-2035.
- Vinale, F., Sivasithamparam, K.; Ghisalberti, E.L.; Marra, R.; Woo, S.L. & Lorito, M. (2008). "Soil Biol. Biochemistry" 40: 1-10
- Lorito, M. (1998). "In *Trichoderma* and *Gliocladium*": Harman, G.E., Kubicek, C.P., Eds.: Taylor & Francis Ltd.: London; Vol. 2, Chapter 4, pp 73-99.
- Reino, J.L; Guerriero, R.F.; Hernandez-Gala, R. & Collado, I.G. (2008). "Phytochem: Rev". 7: 89-123.
- Olabi, T.I. (2004). "Diseases of food crops and their control principles". Amograf Publishers, Nigeria: pp. i-iv + 118.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

