

The Methods for Studying the Structure of Water Clusters (H₂O)_n, where n = 3–20. Water Clusters as Nano-structures

Ignat Ignatov^{1*} Oleg Mosin²

1. Scientific Research Center of Medical Biophysics (SRCMB),

N. Kopernik Street, 32, Sofia 1111, Bulgaria

2. Biotechnology Department, Moscow State University of Applied Biotechnology,

Talalikhina Street, 33, Moscow 109316, Russian Federation

* E-mail of the corresponding author: mbioph@dir.bg

Abstract

In this review it is reported about the research on the structure of intermolecular water cyclic associates (clusters) with general formula (H₂O)_n and their charged ionic clusters [H⁺(H₂O)_n]⁺ and [OH⁻(H₂O)_n]⁻ by means of computer modeling and spectroscopy methods as ¹H-NMR, IR-spectroscopy, NES, DNES, EXAFS-spectroscopy, X-Ray and neutrons diffraction. The computer calculation of polyhedral nanoclusters (H₂O)_n, where n = 3–20 are carried out. Based on this data the main structural mathematical models describing water structure (quasicrystalline, continuous, fractal, fractal-clathrate) have been examined and some important physical characteristics were obtained. The average energy of hydrogen bonding between H₂O molecules in the process of cluster formation was measured by the DNES method compiles -0.1067 ± 0.0011 eV. It was also shown that water clusters formed from D₂O were more stable, than those ones from H₂O due to isotopic effects of deuterium.

Keywords: hydrogen bond, water, structure, clusters, IR-spectroscopy, NES, DNES.

1. Introduction

Water with its anomalous physical and chemical properties outranks all other natural substances on the Earth. The ancient philosophers considered water as the most important component of the matter. It performs a vital role in numerous biochemical and metabolic processes occurring in cells with participation of water, being a universal polar solvent for hydrophilic molecules having an affinity for water. Hydroxyl groups (-OH) in H₂O molecule, are polar and therefore hydrophilic. Moreover water act as a reagent for a big number of chemical reactions (hydrolysis, oxidation-reduction reactions). In chemical processes water due to its high ionizing ability possesses strong amphoteric properties, and can act both as an acid and a base in reactions of chemical exchange.

Modern science has confirmed the role of water as a universal life sustaining component, which defines the

structure and properties of inorganic and organic objects, consisted from water.. The recent development of molecular and structural-chemical concepts has enabled to clarify an explanation of ability of water molecules to form the short-lived hydrogen bonds with neighboring molecules and many other chemical substances to bond them into intermolecular associates. It has also become clear the role of bounded water in forming hydrated substances and their physicochemical conduct in aqueous solutions.

The great scientific and practical interest have the studies of a variety of specific supramolecular structures – cyclic water clusters described by general formula $(\text{H}_2\text{O})_n$, which may be calculated and studied with the help of modern numerical computing methods. The clusters are also important for studying the structure of water and hydration phenomena at molecular level since they form the basic building blocks of the hydrated substances. This paper deals with the mathematical modeling of the water structure and water associates.

2. Nature of Hydrogen Bond in Liquids and Crystals

The peculiarities of chemical structure of H_2O molecule and weak bonds caused by electrostatic forces and donor-acceptor interaction between hydrogen and oxygen atoms in H_2O molecules create favorable conditions for formation of directed intermolecular hydrogen bonds ($\text{O}-\text{H}\dots\text{O}$) with neighboring H_2O molecules, binding them into complex intermolecular associates which composition represented by general formula $(\text{H}_2\text{O})_n$, where n can vary from 3 to 50 (Keutsch & Saykally, 2011). The hydrogen bond – a form of association between the electronegative O oxygen atom and a H hydrogen atom, covalently bound to another electronegative O oxygen atom, is of vital importance in the chemistry of intermolecular interactions, based on weak electrostatic forces and donor-acceptor interactions with charge-transfer (Pauling, 1960). It results from interaction between electron-deficient H-atom of one H_2O molecule (hydrogen donor) and unshared electron pair of an electronegative O-atom (hydrogen acceptor) on the neighboring H_2O molecule; the structure of hydrogen bonding, therefore may be defined as $\text{O}\dots\text{H}^{\delta+} - \text{O}^{\delta-}$. As the result, the electron of the H-atom due to its relatively weak bond with the proton easily shifts to the electronegative O-atom. The O-atom with increased electron density becomes partly negatively charged – δ^- , while the H-atom on the opposite side of the molecule becomes positively charged – δ^+ that leads to the polarization of $\text{O}^{\delta-} - \text{H}^{\delta+}$ covalent bond. In this process the proton becomes almost bared, and due to the electrostatic attraction forces are provided good conditions for convergence of $\text{O}\dots\text{O}$ or $\text{O}\dots\text{H}$ atoms, leading to the chemical exchange of a proton in the reaction $\text{O}-\text{H}\dots\text{O} \leftrightarrow \text{O}\dots\text{H}-\text{O}$. Although this interaction is essentially compensated by mutual repulsion of the molecules' nuclei and electrons, the effect of the electrostatic forces and donor-acceptor interactions for H_2O molecule compiles 5–10 kcal per 1 mole of substance. It is explained by negligible small atomic radius of hydrogen and shortage of inner electron shells, which enables the neighboring H_2O molecule to approach the hydrogen atom of another molecule at very close distance without experiencing any strong electrostatic repulsion.

The H_2O molecule has four sites of hydrogen bonding – two uncompensated positive charges at hydrogen atoms and two negative charges at the oxygen atom. Their mutual disposition is characterized by direction from the centre of regular tetrahedron (nucleus of oxygen atom) towards its vertexes. This allows to one H_2O molecule in condensed state to form up to 4 classical hydrogen bonds, two of which are donor bonds

and the other two – acceptor ones (taking into consideration bifurcate (“two-forked”) hydrogen bond – 5) (Pasichnyk *et al.*, 2008).

A hydrogen bond according to Bernal–Fowler rules (Bernal & Fowler, 1933) is characterized by the following parameters:

a) an oxygen atom of each H₂O molecule is bound with four neighboring hydrogen atoms: by covalent bonding with two own hydrogen atoms, and by hydrogen bonding – with two neighboring hydrogen atoms (as in the crystalline structure of ice); each hydrogen atom in its turn is bound with oxygen atom of neighbour H₂O molecule.

b) on the line of oxygen atom – there can be disposed only one proton H⁺;


c) the proton, which takes part in hydrogen bonding situated between two oxygen atoms, therefore has two equilibrium positions: it can be located near its oxygen atom at approximate distance of 1 Å, and near the neighboring oxygen atom at the distance of 1.7 Å as well, hence both a usual dimer HO–H...OH₂ and an ion pair HO...H–OH₂ may be formed during hydrogen bonding, i.e. the hydrogen bond is part electrostatic (~90%) and part (~10%) covalent (Isaacs *et al.*, 2000). The state of “a proton near the neighboring oxygen” is typical for the interphase boundary, i.e. near water-solid body or water–gas surfaces.

d) the hydrogen bonding of a triad O–H...O possess direction of the shorter O–H (→) covalent bond; the donor hydrogen bond tends to point directly at the acceptor electron pair (this direction means that the hydrogen atom being donated to the oxygen atom acceptor on another H₂O molecule).

The most remarkable peculiarity of hydrogen bond consists in its relatively low strength; it is 5–10 times weaker than chemical covalent bond (Pimentel & McClellan, 1960). In respect of energy hydrogen bond has an intermediate position between covalent bonds and intermolecular van der Waals forces, based on dipole-dipole interactions, holding the neutral molecules together in gasses or liquefied or solidified gasses. Hydrogen bonding produces interatomic distances shorter than the sum of van der Waals radii, and usually involves a limited number of interaction partners. These characteristics become more substantial when acceptors bind H-atoms from more electronegative donors. Hydrogen bonds hold H₂O molecules on 15% closer than if water was a simple liquid with van der Waals interactions. The hydrogen bond energy compiles 5–10 kcal/mole, while the energy of O–H covalent bonds in H₂O molecule – 109 kcal/mole (Arunan *et al.*, 2011). The values of the average energy ($\Delta E_{H...O}$) of hydrogen H...O-bonds between H₂O molecules make up 0.1067 ± 0.0011 eV (Antonov & Galabova, 1992). With fluctuations of water temperature the average energy of hydrogen H...O-bonds in of water molecule associates changes. That is why hydrogen bonds in liquid state are relatively weak and unstable: it is thought that they can easily form and disappear as the result of temperature fluctuations (Ignatov & Mosin, 2013a).

Another key feature of hydrogen bond consists in its cooperativity coupling. Hydrogen bonding leads to the formation of the next hydrogen bond and redistribution of electrons, which in its turn promotes the formation of the following hydrogen bond, which length increasing with distance. Cooperative hydrogen bonding increases the O–H bond length, at the same time causing a reduction in the H...O and O...O distances (Goryainov, 2012). The protons held by individual H₂O molecules may switch partners in an ordered manner within hydrogen networks (Bartha *et al.*, 2003). As the result, aqueous solutions may undergo autoprotolysis, i.e. the H⁺ proton is released from H₂O molecule and then transferred and accepted by the neighboring H₂O molecule resulting in formation of hydronium ions as H₃O⁺, H₅O₂⁺, H₇O₃⁺, H₉O₄⁺,

etc. This leads to the fact, that water should be considered as associated liquid composed from a set of individual H₂O molecules, linked together by hydrogen bonds and weak intermolecular van der Waals forces (Liu *et al.*, 1996). The simplest example of such associate can be a dimer of water:



The energy of the hydrogen bonding in the water dimer is 0.2 eV (~5 kcal/mol), which is larger than the energy of thermal motion of the molecules at the temperature of 300 K. Hydrogen bonds are easily disintegrated and re-formed through an interval of time, which makes water structure quite unstable and changeable (George, 1997). This process leads to structural inhomogeneity of water characterizing it as an associated heterogeneous two-phase liquid with short-range ordering, i.e. with regularity in mutual positioning of atoms and molecules, which reoccurs only at distances comparable to distances between initial atoms, i.e. the first H₂O layer. As it is known, a liquid in contrast to a solid body, is a dynamic system: its atoms, ions or molecules, keeping short-range order in mutual disposition, participate in thermal motion, the character of which is much more complicated than that of crystals. For example H₂O molecules in liquid state under normal conditions (1 atm, 22 °C) are quiet mobile and can oscillate around their rotation axes, as well as to perform the random and directed shifts. This enabled for some individual molecules due to cooperative interactions to “jump up” from one place to another in an elementary volume of water. Random motion of molecules in liquids causes continuous changes in the distances between them. The statistical character of ordered arrangement of molecules in liquids results in fluctuations – continuously occurring deviations not only from average density, but from average orientation as well, because molecules in liquids are capable to form groups, in which a particular orientation prevails. Thus, the smaller these deviations are, the more frequently they occur in liquids.




Figure 1. Hydrogen bonding between four individual H₂O molecules. It is shown the value of the angle between the covalent H–O–H bond in H₂O molecule.

The further important feature is that the hydrogen bonds are spatially oriented. As each H₂O molecule has four sites of hydrogen bond formation (two non-shared electron pairs at an oxygen atom and two uncompensated positive charges at a hydrogen atom), one H₂O molecule in a condensed state is capable to form hydrogen bonds with four H₂O molecules (two donor and two acceptor) (Figure 1), which results in forming a tetrahedron crystal structure clearly observed in ice crystals.

At present time 14 crystalline modifications of ice are known, each of them has its own structure and a character of disposition of hydrogen atoms (Table). Crystals of all ice modifications are made up from H₂O molecules, linked by hydrogen bonds into a 3D carcass, consisting of individual tetrahedrons, formed by four H₂O molecules (Figure 2). In the crystalline structure of natural ice I_h hydrogen bonds are oriented towards the tetrahedron apexes at strictly defined angles equal to 109°5' (in liquid water this angle is 104°5') (Mosin & Ignatov, 2013a). In ice structures I_c, VII and VIII this tetrahedron is nearly the same as a regular 4 triangular tetrahedron. In ice structures II, III, V and VI the tetrahedrons are noticeably distorted. In ice structures VI, VII and VIII two intercrossing systems of hydrogen bonds are distinguished. In the centre of the tetrahedron is located an oxygen atom, at each of the two vertices – H-atom, which electron take part in formation of covalent bond with an electron pair of O-atom. The rest two vertices of the tetrahedron are occupied by two pairs of non -shared electrons of O-atom not participating in formation of molecular bonds.

Table. Ice Crystal modifications and their physical characteristics

Modification	Crystal structure	Hydrogen bond lengths, Å	Angles H–O–H in tetragonals, °
I _h	Hexagonal	2.76	109.5
I _c	Cubic	2.76	109.5
II	Trigonal	2.75–2.84	80–128
III	Tetragonal	2.76–2.8	87–141
IV	Rhombic	2.78–2.88	70.1–109
V	Monoclinic	2.76–2.87	84–135
VI	Tetragonal	2.79–2.82	76–128
VII	Cubic	2.86	109.5
VIII	Cubic	2.86	109.5
IX	Tetragonal	2.76–2.8	87–141
X	Cubic	2.78	109.5
XI	Hexagonal	4.50	90.0
XII	Tetragonal	4.01	90.0
XIII	Monoclinic	7.47	90–109.7
XIV	rhombic	4.08	90.0

Keys: I_h – natural hexagonal ice; I_c – cubic ice.

The carcasses of hydrogen bonds allocate H_2O molecules in form of a spatial hexagon network with internal hollow hexagonal channels. In the nodes of this network O-atoms are orderly organized (crystalline state), forming regular hexagons, while H-atoms have various positions along the bonds (amorphous state). When ice melts, its network structure is destroyed: H_2O molecules begin to fall down into the network hollows, resulting in a denser structure of the liquid – this explains why water is heavier than ice. The hydrogen bonding explains other anomalies of water (anomaly of temperature, pressure, density, viscosity, fluidity etc. According to theoretical calculations, at the melting of the ice breaks about 15% of all hydrogen bonds (Mosin & Ignatov, 2011); by further heating to $40^{\circ}C$ breaks down about half of hydrogen bonds in water associates. In the water vapor hydrogen bonds are absent.

The main difference between the structure of ice and water is more diffuse arrangement of the atoms in the lattice and disturbance of long-range order. Thermal oscillations (fluctuations) lead to bending and breaking of hydrogen bonds. Those out of the equilibrium positions H_2O molecules fall into the adjacent structural voids and for a time held up there, as cavities correspond to the relative minimum of potential energy. This leads to an increase in the coordination number, and the formation of lattice defects. The coordination number (the number of nearest neighbors) varies from 4.4 at $1.5^{\circ}C$ to 4.9 at $80^{\circ}C$.





Figure 2. Hydrogen bonding in crystalline structure of natural ice I_h under the temperature $-10^{\circ}C$ and pressure 1 atm. Grey balls – oxygen atoms, black balls – hydrogen atoms. On the right below shown the structural unit of crystalline ice structure.



a) b)
Figure 3. Structures of water crystals in carbon nanotubes according to computer simulations (Bai *et al.*, 2006) (diameter of nanotubes 1.35–1.90 nm; pressure 10–40000 atm; temperature -23 °C): a) – a general view of the crystal structure of water in nanotubes, b) – the inner wall of the structure of water.




Figure 4. Image of hexamer nanocrystal of ice water (average size ~1 nm) by a scanning tunneling microscope after freezing water to 17 K on the hydrophobic metallic plates of Cu and Ag (Michaelides & Morgenstern, 2007).

Reports about evidence of existence of crystal structures in water have become more frequent in scientific literature nowadays (Michaelides & Morgenstern, 2007). As computer modeling showed, H₂O molecules being placed in carbon nanotubes under high pressure and low temperatures formed crystalline nanostructures like DNA double helix (Bai *et al.*, 2006). In modeling experiment water got frozen inside carbon nanotubes with diameter 1.35–1.90 nm with pressure 10–40000 atm and temperature -23 °C. Hydrogen bonds in an ice crystal got distorted leading to the formation of a double-walled helix. The inner wall of this structure represents a four-fold twisted helix, while the outer consists of four double helices, resembling a helix of DNA molecule (Figure 3). While being frozen at 17 K on hydrophobic surfaces of Cu,

Ag and their salts, water became crystallized into two-dimensional ice hexamer nanocrystal, consisting of six attached together H₂O molecules (Figure 4).

3. Structural Mathematical Models of Water

There are several groups of models, describing structure of liquids – microcrystalline, quasicrystalline, continuous, fractal and fractal-clathrate models. The microcrystalline model of J. Bernal and P. Fowler suggests that water is a nonequilibrium 2-phase liquid containing groups of oriented molecules – microcrystals with several dozens or hundreds of molecules (Bernal & Fowler, 1933; Timothy & Zwier, 2004). Within each microcrystal in liquid a solid body ordering (long-range order) is strictly kept. As water is denser than ice, it is assumed that its molecules are allocated in a different way than in ice: like the atoms of silicon in mineral tridymite or in its more solid modification of quartz. The intermittent increase in water density from 0 to 3.98 °C and other anomalous properties of water was explained by the existence of tridymite component at low temperatures (Pople, 1951).

Quasicrystalline model by analogy with a quasicrystal, i.g. a structure that is ordered but not periodic (Nemukhin, 1996), suggests that relative disposition of particles in liquids is nearly the same as in crystals; deviation from regularity increases with the distance from the initial H₂O molecule; far apart at longer distances there is no regularity in the disposition of H₂O molecules. Each H₂O molecule is surrounded by the four neighbouring ones, which are arranged around it in quite the same way, as they are in an ice crystal. However, in the second layer there appear deviations from regularity rapidly increasing with distance from the initial H₂O molecule. Studies of X-ray scattering in liquids consisting of polyatomic molecules, revealed not only some regular arrangement of H₂O molecules, but consistent pattern in mutual orientation of the molecules (Petkov *et al.*, 2012). This orientation becomes even more expressed for the polar molecules because of the hydrogen bonding effect.

Kinetic theory of liquids proposed by Y.I. Frenkel, also called a “jump-wait” model, explains the structural properties of liquids by peculiarities of thermal motion of their molecules (Frenkel, 1975). The thermal motion of H₂O molecules is characterized by two parameters: a period of oscillation of H₂O molecule around an equilibrium position and a period of “settled life”, i.e. period of oscillation around one particular equilibrium position. Average time of “settled life” of a H₂O molecule, within which H₂O molecules keep unchanged equilibrium orientation is called relaxation time τ :

$$\tau = \tau_0 \times e^{W/RT}, \quad (1)$$

where τ_0 is an average period of oscillations of a H₂O molecule around an equilibrium position (sec), W – a value of potential energy barrier, separating two neighboring equilibrium positions from each other (J), R – Boltzmann constant (J/K), T – absolute temperature (K).

According to calculations the relaxation time at room temperature makes up $\sim 4.5 \cdot 10^{-12}$ sec, while the period of one oscillation of H₂O molecules – 10^{-12} – 10^{-13} sec. That is why each H₂O molecule performs approximately 100 oscillations relative to the same equilibrium position before changing its place. Due to

thermal fluctuations one H₂O molecule within its “settled life” period of time oscillates around an equilibrium position, after then jumps up to new location and it continues to fluctuate up to the next jump. Through these abrupt movements of molecules in liquids occurs diffusion, which, in contrast to the continuous diffusion in gases, called diffusion jump. With increase in temperature the period of “settled life” of H₂O molecules in a temporary state of equilibrium is reduced that brings the structure of water closer to a gas, in which translational and rotational motions of H₂O molecules prevail. Theoretical studies show that, along with the fluctuation of molecules surrounded by his neighbors and activation jumps in liquids occur flowing movement of molecules along with their immediate environment. In other words, being in oscillating state, the molecules in the liquid displaced each time by a certain distance (less than the interatomic distance), causing continuous diffusion. It is believed that in the liquefied inert gases and metals dominates continuous diffusion, while for associated liquids as water is more likely the jump diffusion mechanism. The thermal motion of H₂O molecules leads to continuous changes in distances between them, which cause fluctuations – continual deviations not only from the average density, but from the average orientation of the molecules as well, because H₂O molecules can form groups in which a certain orientation predominates. An attempt to change the water volume (even by small quantity) triggers the process of deformation of hydrogen bonds that fact may explain low water compressibility. Ice melting also causes weakening and deformation of hydrogen bonds, which makes water denser than ice. At the temperature 3.98 °C water acquires anomalous state, in which the quasicrystalline phase is maximally densified by filling up of ice carcass hollows with H₂O molecules. Further increase in temperature and energy of thermal motion of H₂O molecules leads to the gradual disintegration of associated water structures and to the partial rupture of hydrogen bonds with essential reduce of the “settled life” of each H₂O molecule in water associates.

The continuous and “fractal” models consider water as a complex dynamic system with a hydrogen network, forming the empty cavities and non-bonded H₂O molecules distributed within the network. The structure is based upon a dimensional carcasses of individual H₂O molecules, joined together into a multi-molecular associate similar to a clathrate having a configuration of a regular polyhedron (Samoilov, 1963). Stated geometrically, this model represents the packed spheres with varying degrees of packaging. Thus, the pentagonal packing of spheres, showed in Figure 5b, is denser, than the sphere packing model in Figure 5a (its density is 72%). Analogous structure has clathrate hydrates (gas clathrates) – crystalline water-based solids physically resembling ice, in which small non polar gas molecules are placed inside cavities of hydrogen bonded H₂O molecules.





Figure 5. The sphere (a) and the pentagonal (b) packing of balls representing H₂O molecules in clusters.

In 1957 S. Frank and W. Wen proposed a model, postulating arbitrary formation of cyclic associates in water, which presumably had random groups of water associates – the “flickering clusters” with general formula (H₂O)_n, which are in a dynamic equilibrium with free H₂O molecules (Henry *et al.*, 1957). The hydrogen bonds between H₂O molecules are in dynamic equilibrium; they are constantly broken and re-formed in new configurations within a certain time interval; these processes are occurred cooperatively within short-living associated groups of H₂O molecules (clusters), which life spans are estimated from 10⁻¹⁰ to 10⁻¹¹ sec.

Water associates evidently may have polymer structure, because hydrogen bond is on 10% partially covalent bond (Isaacs *et al.*, 2000). In 1990 G.A. Domrachev and D.A. Selivanovsky formulated a model of H₂O-polimers based on the existence of mechanochemical reactions of ionization and dissociation of water in aqueous solutions (Domrachev & Selivanovsky, 1990). Water was thought as dynamically unstable quazi-polymer, composed of (H₂O)_n blocks with partially covalent by 10% hydrogen bonds, permitting at least to 10% of H₂O molecules to comby in a sufficiently long-lived polymer associates. Similarly with mechanochemical reactions in polymers, in case of mechanical impact on water, the energy absorbed by water required for splitting up the H–OH bond, concentrates in a micro-scale area of the liquid water structure.

The splitting up reaction of the H–OH bond in water polymer associates is expressed by the following equation:



where E – the energy of the H–OH bond, 460 kJ/mole; the dot denotes an unpaired electron.

Splitting up of the H–OH bond is accompanied by formation of new disordered bonds between “fragments” of the initial molecules, leading to the formation of fluctuation areas with different density fluctuations that can be observed in aqueous solutions.

Another interesting physical phenomenon was discovered by A. Antonov in 2005 (Antonov, 2005). It was established experimentally that at evaporation of water droplet the contact angle θ decreases discretely to zero, whereas the diameter of the droplet changes insignificantly. By measuring this angle within a regular time intervals a functional dependence $f(\theta)$ can be determined, which is designated by the spectrum of the water state. For practical purposes by registering the spectrum of water state it is possible to obtain information about the averaged energy of hydrogen bonds in an aqueous sample. For this purpose the model of W. Luck is used, which consider water as an associated liquid, consisted of O–H...O–H groups (Luck *et al.*, 1980). The major part of these groups is designated by the energy of hydrogen bonds ($-E$), while the others are free ($E = 0$). The energy distribution function $f(E)$ is measured in electron-volts (eV^{-1}) and may be varied under the influence of various external factors on water as temperature and pressure. For calculation of the function $f(E)$ experimental dependence between the water surface tension (θ) and the energy of hydrogen bonds (E) is established:

$$f(E) = \frac{14,33f(\theta)}{[1-(1+bE)^2]^2}, \quad (2)$$

where $b = 14.33 \text{ eV}^{-1}$

The energy of hydrogen bonds (E) is measured in electron-volts (eV) and is designated by the spectrum of energy distribution. The water spectrum characterized by a non-equilibrium process of water droplets evaporation, thus the term “non-equilibrium energy spectrum of water” (NES) is applied.

The difference $\Delta f(E) = f(\text{samples of water}) - f(\text{control sample of water})$ is called the “differential non-equilibrium energy spectrum of water” (DNES).

DNES is a measure of changes in the structure of water as a result of external influences. The cumulative effect of all other factors is the same for the control sample of water and the water sample, which is under the influence of this impact.




Figure 6. The total number of hydrogen bonds depending on the number of water molecules in clusters.

In 2005 R. Saykally (University of California, USA) calculated the possible number of hydrogen bonds and the stability of water clusters depending on the number of H₂O molecules (Figure 6) (Saykally, 2005). It was also estimated the possible number of hydrogen bonds (100) depending on the number of H₂O molecules (250) in clusters (Sykes, 2007). O. Loboda and O.V. Goncharuk provided data about the existence of icosahedral water clusters consisting of 280 H₂O molecules with the average size up to 3 nm (Loboda & Goncharuk, 2010). The ordering of water molecules into associates corresponds to a decrease in the entropy (randomness), or decrease in the overall Gibbs energy ($G = \Delta H - T\Delta S$). This means that the change in enthalpy ΔH minus the change in entropy ΔS (multiplied by the absolute temperature T) is a negative value. These results are consistent with our data on research of NES and DNES spectrum of water on which it may make conclusion about the number of H₂O molecules in water clusters. NES spectrum of water has energy ranges from -0.08 to -0.14 eV (Figure 7). The spectral range lies in the middle infrared range from 8 to 14 μm ("window" of the atmosphere transparency to electromagnetic radiation). Under these conditions, the relative stability of water clusters depends on external factors, primarily on the temperature.




Figure 7. NES-spectrum of deionized water (chemical purity 99.99%, pH = 6.5–7.5, total mineralization 200 mg/l, electric conductivity 10 $\mu\text{S/cm}$). On the horizontal axis shows the energy of the H...O hydrogen bonds in the associates $-E$ (eV). The vertical axis – energy distribution function $-f$ (eV^{-1}). k – the vibration

frequency of the H–O–H atoms (cm^{-1}); λ – wavelength (mm).

It was shown that the H_2O molecules change their position in clusters depending on the energy of intermolecular H...O hydrogen bonds. The values of the average energy ($E_{\text{H...O}}$) of hydrogen bonds between the H_2O molecules in the formation of cluster associates with formula $(\text{H}_2\text{O})_n$ compile 0.1067 ± 0.0011 eV. As the energy of hydrogen bonds between H_2O molecules increases up to -0.14 eV, the cluster formation of water becomes “destructuring”. In this case, the energy redistribution between the individual H_2O molecules occurs (Figure 7).

All these data indicate that the water is a complex associated non-equilibrium liquid consisting of associative groups containing according to the present data, from 3 to 20 individual H_2O molecules (Tokmachev *et al.*, 2010). Associates can be perceived as unstable groups (dimers, trimers, tetramers, pentamers, hexamers etc.) in which H_2O molecules are linked by van der Waals forces, dipole-dipole and other charge-transfer interactions, including hydrogen bonding. At room temperature, the degree of association of H_2O molecules may vary from 2 to 6. In 1993 K. Jordan (USA) (Tsai & Jordan, 1993) calculated the possible structural modifications of small water clusters consisting of six H_2O molecules (Figure 8a–c). Subsequently, it was shown that H_2O molecules capable of hydrogen bonding by forming the structures representing topological 1D rings and 2D chains composed from numerous H_2O molecules. Interpreting the experimental data, they are considered as pretty stable elements of the structure. According to computer simulations, clusters are able to interact with each other through the exposed protons on the outer surfaces of hydrogen bonds to form new clusters of more complex composition.




Figure 8. Calculation of small water cluster structures (a–c) with general formula $(\text{H}_2\text{O})_n$, where $n = 6$ (Tsai & Jordan, 1993).

In 2000 it was deciphered the structure of the trimer water, and in 2003 – tetramer, pentamer and the water hexamer (Wang & Jordan, 2003). Structures of water clusters with formula $(\text{H}_2\text{O})_n$, where $n = 3–5$, similar to the cage structure. Hexagonal structure with $n = 6$, consisting of six H_2O molecules at the hexagon vertices, is less stable than the cage structure. In the hexagon structure four H_2O molecules can be cross-linked by hydrogen bonds (Figure 9).




Figure 9. Cluster structure of a trimer, tetramer, pentamer and hexamer of water.

Quantum-chemical calculations of middle size clusters with the general formula $(\text{H}_2\text{O})_n$, where $n = 6-20$, have shown that the most stable structures are formed by the interaction of tetrameric and pentameric structures (Maheshwary *et al.*, 2001; Choi & Jordan, 2010). Thus the structures of $(\text{H}_2\text{O})_n$, where $n = 8, 12, 16$, and 20 are cubic, and structures $(\text{H}_2\text{O})_n$ where $n = 10$ and 15 – pentagons. Other structures with $n = 9, 11, 13, 14, 17, 18$ and 19 evidently have a mixed composition. Large tetrahedron clusters as $(\text{H}_2\text{O})_{196}$, $(\text{H}_2\text{O})_{224}$, $(\text{H}_2\text{O})_{252}$ (Figure 10) composed from the smaller ones formed a vertex of a $(\text{H}_2\text{O})_{14}$ tetrahedron are also described (Chaplin, 2011).




Figure 10. Tetrahedron water clusters $(\text{H}_2\text{O})_n$ with different symmetry ($n = 196, 224, 252$).

It is reasonable that the structure of liquid water should be related to the structure of hexagonal ice, formed from H_2O tetrahedrons, which exist under atmospheric pressure. In the computer simulation H_2O tetrahedrons grouped together, to form a variety of 3D-spatial and 1D, 2D-planar structures, the most common of which is hexagonal structure where 6 H_2O molecules (tetrahedrons) are combined into a ring. A similar type of structure is typical for ice I_h crystals. When ice melts, its hexagonal structure is destroyed, and a mixture of clusters consisting of tri-, tetra-, penta-, and hexamers of water and free H_2O molecules is formed. Structural studies of these clusters are significantly impeded, since the water is perceived as a mixture of different clusters that are in dynamic equilibrium with each other.

S. Zenin (Russia) calculated a cluster model based on a minimum "quantum" of water (Zenin, 1999), which is a 4 triangular tetrahedron composed of four 12 pentagonal dodecahedrons (Figure 11). The "quantum" consisted of 57 H_2O molecules interacting with each other at the expense of free hydrogen bonds exposed on the surface. Of 57 H_2O molecules in the "quantum" 17 H_2O molecules compose tetrahedral completely hydrophobic, i.e. saturated with four hydrogen bonds in the central carcass, and four dodecahedra on the surface of each there are 10 centers for the formation of hydrogen bond (O-H or O). 16 "quanta" form a bigger cluster structure consisted from 912 H_2O molecules similar to a tetrahedron.




Figure 11. Model of water associates according to S.V. Zenin: *a)* – associate of 57 H₂O molecules, tetrahedron of four dodecahedron (“quantum”); *b)* – bulk packaging of "quanta" (Zenin, 1999).




Figure 12. Icosahedron cluster based on 100 H₂O molecules and the underlying structure according to M. Chaplin's model.

M. Chaplin (London South Bank University, UK) (Chaplin, 2011) calculated the water structure based on 20 triangular icosahedron (Figure 12). The structure is based on minimal tetrahedral water cluster, consisting of 14 H₂O molecules. Arrangement of 20 of these 14-molecule structures forms an icosahedral network, consisting from 280 H₂O molecules. Each 280-molecule in icosahedral contains several substructures with each H₂O molecule involved in four hydrogen bonds; two as donor and two as acceptor. 13 overlapping icosahedra may form a more large cluster (tricontahedron) consisting from 1820 H₂O molecules, which has twice more H₂O molecules than in the previous model.

The clusters, evidently, may be rather stable under a certain conditions, and can be obtained in the isolated

state within a very short interval of time. There is also a reason to believe that the charged ions stabilize the clusters. Therefore, the clusters can be divided into positively and negatively charged ionic clusters – $[(\text{H}_2\text{O})_n]^+$, $[(\text{H}_2\text{O})_n]^-$, and not having a charge – neutral clusters with general formula $(\text{H}_2\text{O})_n$. Clusters, containing 20 individual H_2O molecules and a proton in the form of hydronium ion H_3O^+ (“magic” number) form the most stable ionic clusters $(\text{H}_2\text{O})_{20}\text{H}_3\text{O}^+$ or $(\text{H}_2\text{O})_{21}\text{H}^+$ (Figure 13) (Cui *et al.*, 2006). It is assumed that the stability of ionic clusters is due to the special clathrate structure in which 20 H_2O molecules are formed 12 pentagonal dodecahedron, in which cavities is captured the H_3O^+ ion. It is occurred because of all the clusters only the dodecahedron has a large cavities enough to accommodate a bulky H_3O^+ ion. Subsequently, due to cooperative interactions H_3O^+ is further able to move to the surface of the cluster and lose a proton H^+ leading to the formation of hydronium ions like H_5O_2^+ , H_7O_3^+ , and H_9O_4^+ , fixed on the surface of the cluster. These data show the diversity of supposed water structures and the complexity of molecular interactions between the different water clusters, the nature of which we are only beginning to understand.




Figure 13. Formation of ionic clusters $(\text{H}_2\text{O})_{20}\text{H}_3\text{O}^+$ or $(\text{H}_2\text{O})_{21}\text{H}^+$ with captured hydronium ion H_3O^+ .

4. Methods of Study Water Clusters

Cluster structures in water calculated theoretically on computer were confirmed by ^1H -NMR, IR, Raman, Compton scattering, EXAFS-spectroscopy and X-ray diffraction (Ignatov, 2005; Mosin & Ignatov, 2011). Information obtained with using the modern detection methods corresponds to femtosecond time, i.e instantaneous dynamics of intermolecular interactions in the molecular scale. The presence of hydrogen bonding causes the noticeable effect on vibrational and ^1H -NMR spectra. In ^1H -NMR the chemical shift of the proton involved in the hydrogen bonding shifts about 0.01 ppm reducing strength of hydrogen bonding while the temperature is raised (Yamaguchi *et al.*, 2001). Increased extent of hydrogen bonding within clusters results in a similar effect; the higher chemical shifts with greater cooperativity, the shorter hydrogen bonded O-H...O distances. ^1H peaks shift to greater ppm with increasing hydrogen bonding strength.

In IR-spectroscopy the characteristic vibration frequency bands containing hydrogen are reduced in spectra if hydrogen atom is included in the hydrogen bonding. Infrared absorption bands, such as OH-groups are much expanded when the hydrogen bond is formed, and their intensity increases. The main stretching band

in liquid water is shifted to a lower frequency (ν_3 , 3480 cm^{-1} and ν_1 , 3270 cm^{-1}), while the bending frequency (ν_2 , 1640 cm^{-1}) increased by hydrogen bonding (Ohno *et al.*, 2005). Increased strength of hydrogen bonding shifts the stretch vibration to lower frequencies with greatly increased intensity in the infrared due to the increased dipoles. With raising the temperature the stretch vibrations shift to higher frequency, while the intramolecular vibrations shift to lower frequencies (Ignatov, 2012; Ignatov & Mosin, 2013b). Hydrogen bond energy lies in the range of 2.3 kcal/mole for the N–H...O-bonds up to 7.0 kcal/mol for the bonds with hydrogen fluoride F–H...F. The strength of the hydrogen bonding depends on the cooperative/anticooperative character of the surrounding hydrogen bonds with the strongest hydrogen bonds giving the lowest vibrational frequencies.

In X-ray absorption spectroscopy (EXAFS-spectroscopy) X-ray radiation excites the electrons of the inner shell of an oxygen atom in H_2O molecules and stipulates their transition onto unoccupied upper electronic levels in the molecules (Wernet *et al.*, 2004). Probability of electron transitions and characteristics of the absorption contour depends mostly on the molecular environment. This allows by varying the energy of the X-rays to study the distribution of associations for HOH...OH₂ bond judging on lengths and angles of the covalently bonded hydrogen atom in the molecule. EXAFS spectrum near the oxygen atom is also sensitive to hydrogen bonding. This method is used to obtain information about the molecular structure of water in the first coordination sphere. Since the time of excitation of electrons is much smaller than the vibrational motions in liquids, X-ray probe of the electronic structure provides information about the instantaneous changes of configurations of the water structure.

Diffraction techniques (X-ray and neutron diffraction) on liquid water allow to calculate the function of density radial distribution (O or H) and the probability of detection of H_2O molecules at a certain distance from a randomly chosen individual H_2O molecule (Tokushima *et al.*, 2008). This allows to detect the irregularities in water by constructing the radial distribution function, i.e. the distance between the atoms of O, H, and O–H in H_2O molecule and its nearest neighbors. Thus, the distribution of the distances between the oxygen atoms at a room temperature, gives three major peaks measured at 2.8, 4.5 and 6,7 Å. The first maximum corresponds to the distance to the nearest neighbor and its value is approximately equal to the length of the hydrogen bond. The second maximum is close to the average edge length of a tetrahedron, as H_2O molecules in the crystalline structure of ice I_h arranged at the vertices of the tetrahedron allocated around the center of the molecule. The third peak, expressed very weakly, corresponds to the distance to the third and more distant neighboring H_2O molecules on the hydrogen network. In 1970 I.S. Andrianov and I.Z. Fisher calculated the distance up to the eighth of the neighboring H_2O molecule; the distance to the fifth the neighboring H_2O molecule turned out to be 3 Å, and to the sixth molecule – 3,1 Å. This allowed to draw conclusions about the geometry of hydrogen bonds and farther surroundings of H_2O molecules.

Another method for structural studies – neutron diffraction is similar to X-ray diffraction. However, because the neutron scattering lengths vary slightly among different atoms, this method is limited in the case of the isomorphous substitution of hydrogen atoms in H_2O molecule by deuterium (D). In practice generally operate with a crystal whose molecular structure is approximately defined by other methods. Then, it is measured the intensity of the neutron diffraction of this crystal. From these results, the Fourier transform is carried out; the measured neutron intensity and phase are using for calculation. Then, on the resultant Fourier map hydrogen and deuterium atoms are represented with much greater atomic weight than

on the electron density map, as the contribution of these atoms in neutron scattering is essentially big. On the resulting density map is determined the arrangement of hydrogen ^1H (negative density) and deuterium D (positive density) atoms. A variation of the method consists in that the crystal of an ordinary protonated water (H_2O) before measurements kept in 99.9 at.% of heavy water (D_2O). In this case the neutron diffraction can not only establish the localization of the hydrogen atoms, as well as to identify those protons that can be exchanged by deuterium, that is particularly important for the study of isotopic (H–D) exchange. Such information in some cases may help confirm the correctness of the water structure established by other methods.

Clusters formed of D_2O are some more stable and resistant than those ones from H_2O due to isotopic effects of deuterium caused by 2-fold increasing nuclear mass of deuterium (molecular mass of D_2O is more by 11% than that of H_2O). The structure of D_2O molecule is the same, as that of H_2O , with small distinction in values of lengths of covalent bonds. D_2O crystals have the same structure as a conventional ice I_h , the difference in unit cell size is very insignificant (0.1%). But they are heavy (0.982 g/cm^3 at 0°C over 0.917 g/cm^3 for conventional ice). D_2O boils at 101.44°C , freezes at 3.82°C , has density at 20°C 1.105 g/cm^3 , and the maximum density occurs not at the 3.89°C , as for H_2O , but at 11.2°C (1.106 g/cm^3). The mobility of D_3O^+ ion is 28.5% lower than that of H_3O^+ ion and OD^- ion – 39.8% lower than that of OH^- ion, the constant of ionization of D_2O is less than the constant of ionization of H_2O , which means that D_2O has a bit more hydrophobic properties than H_2O . All these effects lead that the hydrogen bonds formed by deuterium atoms differ in strength and energy from ordinary hydrogen bonds (O–H length 1.01 \AA , O–D length 0.98 \AA , D–O–D angle 106°). Commonly used molecular models use O–H lengths $\sim 0.955 \text{ \AA}$ and 1.00 \AA and H–O–H angles from $\sim 105.5^\circ$ to $\sim 109.4^\circ$. The substitution of H with D atom affects the stability and geometry of hydrogen bonds in apparently rather complex way and may, through the changes in the hydrogen bond zero-point vibrational energies, alter the conformational dynamics of hydrogen (deuterium)-bonded structures of associates. In general, isotopic effects stabilize hydrogen bond with participation of deuterium, resulting in somewhat greater stability of associates formed from D_2O molecules (Mosin & Ignatov, 2013b).

As a result of experiments on quasi-elastic neutron scattering was measured the most important parameter – the coefficient of self-diffusion of water at different temperatures and pressures. To analyse the self-diffusion coefficient on quasi-elastic neutron scattering, it is necessary to know the character of moving of molecules. If they move in accordance with the “jump-wait” model, then the “settled” life time (the time between jumps) of H_2O molecule compiles 3.2 ps. The newest methods of femtosecond laser spectroscopy allow to estimate the lifetime of the broken hydrogen bonds: a proton needs 200 fs to find a partner.

The studying the full details of the structure of associative elements in water can be made, considering all the parameters by computer simulation or numerical experiment. For this in a given space is chosen random ensemble of n H_2O molecules and are optimal parameters – the energy of interatomic interactions, bond length, the arrangement of atoms and molecules, the most consistent with the diffraction data. Data thus obtained are then extrapolated to the actual water structure, and further used to calculate thermodynamic parameters. Data obtained by computer experiments, show that the nature of the thermal motion of the molecules in the liquids corresponds on the whole to a vibration of the individual H_2O molecules near the equilibrium centers, with occasional jumps up to the new position.

The comparative analysis of IR-spectra of H₂O solutions and its deuterated analogues (D₂O, HDO) is of considerable interest for biophysical studies, because at changing of the atomic mass of hydrogen by deuterium atoms in H₂O molecule their interaction will also change, although the electronic structure of the molecule and its ability to form H-bonds, however, remains the same. The IR spectra of water usually contain three absorption bands, which can be identified as 1 – absorption band of the stretching vibration of OH⁻ group; 2 – absorption band of the first overtone of the bending vibration of the molecule HDO; 3 – absorption band of stretching vibration of OD⁻ group. OH⁻ group is able to absorb much infrared radiation in the infrared region of the IR-spectrum. Because of its polarity, these groups typically react with each other or with other polar groups to form intra-and intermolecular hydrogen bonds. The hydroxyl groups not involved in formation of hydrogen bonds are usually given the narrow bands in IR spectrum and the associated groups – broad intense absorption bands at lower frequencies. The magnitude of the frequency shift is determined by the strength of the hydrogen bond. Complication of the IR spectrum in the area of OH⁻ stretching vibrations can be explained by the existence of different types of associations, a manifestation of overtones and combination frequencies of OH⁻ groups in hydrogen bonding, as well as the proton tunneling effect (on the relay mechanism. Such complexity makes it difficult to interpret the IR spectrum and partly explains the discrepancy in the literature available on this subject.

The local maximums in IR-spectra reflect vibrational-rotational transitions in the ground electronic state; the substitution with deuterium changes the vibrational-rotational transitions in H₂O molecule, that is why it appear other local maximums in IR-spectra. In the water vapor state, the vibrations involve combinations of symmetric stretch (ν_1), asymmetric stretch (ν_3) and bending (ν_2) of the covalent bonds with absorption intensity (H₂O) $\nu_1; \nu_2; \nu_3 = 2671; 1178.4; 2787.7 \text{ cm}^{-1}$. For liquid water absorption bands are observed in other regions of the IR-spectrum, the most intense of which are located at 2100, cm^{-1} and 710–645 cm^{-1} . For D₂O molecule these ratio compiles 2723.7, 1403.5 and 3707.5 cm^{-1} , while for HDO molecule – 2671.6, 1178.4 and 2787.7 cm^{-1} . HDO (50 mole% H₂O + 50 mole% ²H₂O; ~50 % HDO, ~25 % H₂O, ~25 % D₂O) has local maxima in IR-spectra at 3415 cm^{-1} , 2495 cm^{-1} 1850 cm^{-1} and 1450 cm^{-1} assigned to OH⁻-stretch, OD⁻-stretch, as well as combination of bending and libration and HDO bending respectively.

In the IR-spectrum of liquid water absorbance band considerably broadened and shifted relative to the corresponding bands in the spectrum of water vapor. Their position depends on the temperature (Ignatov & Mosin, 2013b). The temperature dependence of individual spectral bands of liquid water is very complex (Zelmann, 1995). Furthermore, the complexity of the IR-spectrum in the area of OH⁻ stretching vibration can be explained by the existence of different types of H₂O associations, manifestation of overtones and composite frequencies of OH⁻ groups in the hydrogen bonds, and the tunneling effect of the proton (for relay mechanism) (Yukhnevitch, 1973). Such complexity makes it difficult to interpret the spectrum and partly explains the discrepancy in the literature available on this subject.

In liquid water and ice the IR-spectra are far more complex than those ones of the vapor due to vibrational overtones and combinations with librations (restricted rotations, i.g. rocking motions). These librations are due to the restrictions imposed by hydrogen bonding (minor L₁ band at 395.5 cm^{-1} ; major L₂ band at 686.3 cm^{-1} ; for liquid water at 0 °C, the absorbance of L₁ increasing with increasing temperature, while L₂ absorbance decreases but broadens with reduced wave number with increasing temperature (Brubach et al.,

2005). The IR spectra of liquid water usually contain three absorbance bands, which can be identified on absorption band of the stretching vibration of OH⁻ group; absorption band of the first overtone of the bending vibration of the molecule H²O and absorption band of stretching vibration of OD⁻ group (Max & Chapados, 2009). Hydroxyl group OH⁻ is able to absorb much infrared radiation in the infrared region of the IR-spectrum. Because of its polarity, these groups typically react with each other or with other polar groups to form intra-and intermolecular hydrogen bonds. The hydroxyl groups, which are not involved in formation of hydrogen bonds usually produce the narrow bands in IR spectrum, while the associated groups – broad intense absorbance bands at lower frequencies. The magnitude of the frequency shift is determined by the strength of the hydrogen bond. Complication of the IR spectrum in the area of OH⁻ stretching vibrations can be explained by the existence of different types of associations of H₂O molecules, a manifestation of overtones and combination frequencies of OH⁻ groups in hydrogen bonding, as well as the proton tunneling effect (on the relay mechanism).

Assignment of main absorption bands in the IR-spectrum of liquid water is given in Table 4. The IR spectrum of H₂O molecule was examined in detail from the microwave till the middle (4–17500 cm⁻¹) visible region and the ultraviolet region – from 200 nm⁻¹ to ionization limit at 98 nm⁻¹ (Walrafen, 1972). In the middle visible region at 4–7500 cm⁻¹ are located rotational spectrum and the bands corresponding to the vibrational-rotational transitions in the ground electronic state. In the ultraviolet region (200 to 98 nm⁻¹) are located bands corresponding to transitions from the excited electronic states close to the ionization limit in the electronic ground state. The intermediate region of the IR-spectrum – from 570 nm to 200 nm corresponds to transitions to higher vibrational levels of the ground electronic state.

Results of IR-spectroscopy with device Infra Spec VFA-IR show that at 4.1 μm, even at low concentrations of deuterium of 0.35 and 0.71%, there is observed a decline in the local maximums relative to the local maximum of 100% pure water (the local maximums in IR-spectra reflect vibrational-rotational transitions in the ground electronic state because at changing the atomic mass of hydrogen and deuterium atoms in the water molecule their interaction will also change, although the electronic structure of the molecule and its ability to form H-bonds, however, remains the same; with the substitution with deuterium the vibrational-rotational transitions are changed, that is why it appears other local maximums in IR-spectra. These results are shown in Figure 14. The result is reliable regarding the content of deuterium in natural waters from 0.015–0.03%.

ANALYSIS




Figure 14. The typical IR-spectra of water with varying content of deuterium

At further transition from H_2O monomers to H_4O_2 dimer and H_6O_3 trimer absorption maximum of valent stretching vibrations of the O-H bond is shifted toward lower frequencies ($\nu_3 = 3490 \text{ cm}^{-1}$ and $\nu_1 = 3280 \text{ cm}^{-1}$) (Eisenberg & Kauzmann, 1969) and the bending frequency increased ($\nu_2 = 1644 \text{ cm}^{-1}$) because of hydrogen bonding. The increased strength of hydrogen bonding typically shifts the stretch vibration to lower frequencies (red-shift) with greatly increased intensity in the infrared due to the increased dipoles. In contrast, for the deformation vibrations of the H—O—H, it is observed a shift towards higher frequencies. Absorption bands at 3546 and 3691 cm^{-1} were attributed to the stretching modes of the dimer $[(\text{H}_2\text{O})_2]$. These frequencies are significantly lower than the valence modes of ν_1 and ν_3 vibrations of isolated H_2O molecules at 3657 and 3756 cm^{-1} respectively). The absorption band at 3250 cm^{-1} represents overtones of deformation vibrations. Among frequencies between 3250 and 3420 cm^{-1} is possible Fermi resonance (this resonance is a single substitution of intensity of one fluctuations by another fluctuation when they accidentally overlap each other). The absorption band at 1620 cm^{-1} is attributed to the deformation mode of the dimer. This frequency is slightly higher than the deformation mode of the isolated H_2O molecule (1596 cm^{-1}). A shift of the band of deformation vibration of water in the direction of high frequencies at the transition from a liquid to a solid state is attributed by the appearance of additional force, preventing O-H bond bending. Deformation absorption band in IR-spectrum of water has a frequency at 1645 cm^{-1} and a very weak temperature dependence. It changes little in the transition to the individual H_2O molecule at a frequency of 1595 cm^{-1} . This frequency is found to be sufficiently stable, while all other frequencies are greatly affected by temperature changes, the dissolution of the salts and phase transitions. It is believed that the persistence of deformation oscillations is stipulated by processes of intermolecular interactions, i.g. by the change in bond angle as a result of interaction of H_2O molecules with each other, as well as with cations and anions.

Table 2. The assignment of main frequencies in IR-spectra of liquid water H₂O and D₂O

Main vibrations of liquid H ₂ O and ² H ₂ O				
Vibration(s)	H ₂ O (t = 25 °C)		D ₂ O (t = 25 °C)	
	ν , cm ⁻¹	E ₀ , M ⁻¹ cm ⁻¹	ν , cm ⁻¹	E ₀ , M ⁻¹ cm ⁻¹
Spinning ν_1 + deformation ν_2	780-1645	21.65	1210	17.10
Composite $\nu_1 + \nu_2$	2150	3.46	1555	1.88
Valence symmetrical ν_1 , valence asymmetrical ν_3 , and overtone $2\nu_2$	3290-3450	100.65	2510	69.70

Thus the study of the characteristics of the IR spectrum of water allows to answer the question not only on the physical parameters of the molecule and the covalent bonds at isotopic substitution with deuterium, but also to make a certain conclusion on associative environment in water. The latter fact is important in the study of structural and functional properties of water associates and its isotopomers at the isotopic substitution with deuterium. The substitution of H with D affects the stability and geometry of hydrogen bonds in an apparently rather complex way and may, through the changes in the hydrogen bond zero-point vibration energies, alter the conformational dynamics of hydrogen (deuterium)-bonded structures of macromolecules as DNA and proteins in D₂O (Cleland, 1976).

5. Conclusion

The experimental data obtained during the last years suggest that water is a complex dynamic associative system, consisting of tens and possibly hundreds individual H₂O molecules binding by multiple intermolecular hydrogen bonds, being in a state of dynamic equilibrium. Up till now is scientifically proven the existence of associative water clusters with general formula (H₂O)_n, where n = 3–20. Although calculated structural models explain pretty well many anomalous properties of water and being in a good agreement with the experimental data on the diffraction of X-rays and neutrons, Raman, Compton scattering and EXAFS-spectroscopy, they are the most difficult to agree with the dynamic properties of water – flow, viscosity and short relaxation times, which are measured by picoseconds.

References

- Antonov A. & Galabova T. (1992) Reports from the 6th Nat. Conference of Biomedical Physics and Engineering. Sofia.
- Antonov A. (2005) *Research of the Nonequilibrium Processes in the Area in Allocated Systems*. Diss. Thesis Doctor of Physical Sciences, Sofia: Blagoevgrad.
- Arunan E., Desiraju G.R., Klein R.A. et al. (2011) Definition of the hydrogen bond. *Pure Appl. Chem.*, 83(8), 1637–1641.

- Bai J., Wang J. & Zeng C. (2006) Multiwalled Ice Helices and Ice Nanotubes. *Proc. Natl. Acad. Sci.*, 3, 104-117.
- Atanasov, A., Karadzhov, S., Ivanova, E., Mosin, O. V. & Ignatov, I. (2014) Study of the Effects of Electrochemical Aqueous Sodium Chloride Solution (Anolite) on the Virus of Classical Swine Fever Virus. Mathematical Models of Anolyte and Catolyte as Types of Water, *Journal of Medicine, Physiology and Biophysics*, 4: 1-26.
- Bartha F., Kapuy O., Kozmutza C & Van Alsenoy C. (2003) Analysis of weakly bound structures: hydrogen bond and the electron density in a water dimer. *J. Mol. Struct. (Theochem)* 666, 117–122.
- Bernal J.D. & Fowler R.H. (1933) A Theory of Water and Ionic Solutions, with Particular Reference to Hydrogen and Hydroxyl Ions. *J. Chem. Phys.*, 1, 515–548.
- Brubach, J.B., Mermet, A., Filabozzi, A., Gerschel, A. & Roy, P. (2005) Signatures of the hydrogen bonding in the infrared bands of water. *J. Chem. Phys.*, 122: 184509.
- Chaplin M. (2011) The Water Molecule, Liquid Water, Hydrogen Bonds and Water Networks / in: *Water The Forgotten Biological Molecule*, D.Le Bihan & H.Fukuyama (eds.), Singapore: Pan Stanford Publishing Pte. Ltd.
- Choi T.N. & Jordan K.D. (2010) Application of the SCC-DFTB Method to $H^+(H_2O)_6$, $H^+(H_2O)_{21}$, and $H^+(H_2O)_{22}$. *J. Phys. Chem. B*, 114, 6932–6936.
- Christie R.A. & Jordan K.D. (2005) Monte Carlo Simulations of the Finite Temperature Properties of $(H_2O)_6$ / in: *Theory and Applications of Computational Chemistry: The First 40 Years, A Volume of Technical and Historical Perspectives*, Ed. C.E. Dykstra, G.Frenking, K.S. Kim & G. Scuseria. New York: Theochem.
- Cleland W.N. (1976) Isotope effects on enzyme-catalyzed reactions. Ed. W.N. Cleland, M.N. O'Leary, D.D. Northrop. Baltimore, London, Tokyo, University Park Press, 303 p.
- Cui J., Liu H. & Jordan K.D. (2006) Theoretical Characterization of the $(H_2O)_{21}$ Cluster: Application of an n-body Decomposition Procedure. *J. Phys. Chem.*, 110, 18872–18878.
- Domrachev G.A. & Selivanovsky D.A. (1990) *The Role of Sound and Liquid Water as Dynamically Unstable Polymer System in Abiogenous Production of Oxygen and the Origin of Life on the Earth*, Preprint N 1'90. Gorky: Inst. of Organometallic Chem. of the USSR Acad. Sci. [in Russian].
- Eisenberg, D. & Kauzmann, W. (1969) *The Structure and Properties of Water*. Oxford University Press, London.
- Frenkel Y.I. (1975) *Kinetic theory of liquids*. Leningrag: Nauka [in Russian].
- George A.J. (1997) *An Introduction to Hydrogen Bonding (Topics in Physical Chemistry)*. London: Oxford University Press.
- Goryainov S.V. (2012) A model of phase transitions in double-well Morse potential: Application to hydrogen bond. *Physica B*, 407, 4233–4237.
- Henry S., Frank S. & Wen-Yang Wen. (1957) Ion-solvent interaction. Structural Aspects of Ion-solvent Interaction in Aqueous Solutions: a Suggested Picture of Water Structure. *Discuss. Faraday Soc.*, 24, 133–140.
- Ignatov I. & Mosin O.V. (2013a) Isotopic Composition of Water and its Temperature in Modeling of Primordial Hydrosphere Experiments. *Nauchnoe Obozrenie*, 1, 17–27 [in Russian].

- Ignatov I. & Mosin O.V. (2013b) Possible Processes for Origin of Life and Living Matter with modeling of Physiological Processes of Bacterium *Bacillus Subtilis* in Heavy Water us Model System. *Journal of Natural Sciences Research*, 83(8), 65–76.
- Ignatov, I., Antonov, A. & Galabova, T. (1998) Medical Biophysics – Biophysical Fields of Man, *Gea Libris*, Sofia.
- Ignatov, I., Antonov, A. & Galabova, T., (2002) Scientific Research Studies with Christos Drossinakis (October 2001 – October 2002), *Int. Conference “Man and Nature”*, SRCMB, Sofia.
- Ignatov, I. (2005) Energy Biomedicine, *Gea-Libris*, Sofia: 1–88.
- Ignatov, I. (2007) There are not Reliable Results with Research with Infrared Spectroscopy of Homeopathic Solutions after Avogadro’s Number, *Ministry of Health*, 196-199. [in Russian]
- Marinov, M., Ignatov, I. (2008) Color Kirlian Spectral Analysis. Color Observation with Visual Analyzer, *Euromedica*, Hanover, 57-59
- Ignatov, I. (2010) Which water is optimal for the origin (generation) of life? *Euromedica*, Hanover: 34-35.
- Ignatov, I. (2011) Entropy and time in living matter, *Euromedica*: 74.
- Ignatov, I., Tsvetkova, V. (2011) Water for the Origin of Life and “Informationability” of Water, Kirlian (Electric Images) of Different Types of Water, *Euromedica*, Hanover: 62-65.
- Ignatov, I. (2012) Origin of Life and Living Matter in Hot Mineral Water, Conference on the Physics, Chemistry and Biology of Water, *Vermont Photonics*, USA.
- Ignatov, I., & Mosin, O.V. (2012) Isotopic Composition of Water and its Temperature in Modeling of Primordial Hydrosphere Experiments, VIII Intern. Conference Perspectives of the Development of Science and Technique, *Biochemistry and Biophysics*, **15**: 41–49.
- Ignatov, I. & Mosin, O. V. & Naneva, K. (2012) Water in the Human Body is Information Bearer about Longevity, *Euromedica*, Hanover: 110-111.
- Ignatov, I. & Mosin, O. V. (2013) Method for Color Coronal (Kirlian) Spectral Analysis, *Biomedical Radio electronics, Biomedical Technologies and Radio electronics*, **3**: 38-47. [in Russian]
- Ignatov, I. & Mosin, O. V. (2013) Modeling of Possible Processes for Origin of Life and Living Matter in Hot Mineral and Seawater with Deuterium, *Journal of Environment and Earth Science*, **3**(14): 103-118.
- Ignatov, I., Mosin, O. V. (2013) Structural Mathematical Models Describing Water Clusters, *Journal of Mathematical Theory and Modeling*, **3** (11): 72-87.
- Ignatov, I. & Mosin, O.V. (2014) Photoreceptors in Visual Perception and Additive Color Mixing. Bacteriorhodopsin in Nano- and Biotechnologies, *Advances in Physics Theories and Applications*, **27**: 20-37.
- Ignatov, I. & Mosin, O. V. (2014) The Structure and Composition of Carbonaceous Fullerene Containing Mineral Shungite and Microporous Crystalline Aluminosilicate Mineral Zeolite. Mathematical Model of Interaction of Shungite and Zeolite with Water Molecules *Advances in Physics Theories and Applications*, **28**: 10-21.
- Ignatov, I., Mosin, O.V., Velikov, B., Bauer, E. & Tyminski, G. (2014) Longevity Factors and Mountain Water as a Factor. Research in Mountain and Field Areas in Bulgaria, *Civil and Environmental Research*, **6** (4): 51-60.

- Ignatov, I., Mosin, O. V., Niggli, H.&Drossinakis, Ch. (2014) Evaluating Possible Methods and Approaches for Registering of Electromagnetic Waves Emitted from the Human Body, *Advances in Physics Theories and Applications*, **30**: 15-33.
- Ignatov, I., Mosin, O.V.&Drossinakis, Ch. (2014) Infrared Thermal Field Emitted from Human Body. Thermovision, *Journal of Medicine, Physiology, Biophysics*, **1**:1-12.
- Ignatov, I., Mosin, O.V.&Velikov, B. (2014) Longevity Factors and Mountain Water of Bulgaria in Factorial Research of Longevity, *Journal of Medicine, Physiology, Biophysics*, **1**:13-33.
- Ignatov, I.&Mosin,O.V. (2014) Visual Perception. Electromagnetic Conception for the Eyesight. Rhodopsin and Bacteriodopsin, *Journal of Medicine, Physiology and Biophysics*, **2**:1-19.
- Ignatov, I.&Mosin,O.V. (2014) The Structure and Composition of Shungite and Zeolite. Mathematical Model of Distribution of Hydrogen Bonds of Water Molecules in Solution of Shungite and Zeolite, *Journal of Medicine, Physiology and Biophysics*, **2**: 20-36.
- Ignatov, I., Mosin,O.V., Velikov, B., Bauer, E.&Tyminski, G. (2014) Research of Longevity Factors and Mountain Water as a Factor in Teteven Municipality, Bulgaria, *Journal of Medicine, Physiology and Biophysics*, **2**: 37-52.
- Ignatov, I.&Mosin,O.V. (2014) Modeling of Possible Processes for Origin of Life and Living Matter in Hot Mineral Water. Research of Physiological Processes of Bacterium Bacillus Subtilis in Hot Heavy Water, *Journal of Medicine, Physiology and Biophysics*, **2**: 53-70.
- Ignatov, I.&Mosin,O.V. (2014) Mathematical Models of Distribution of Water Molecules Regarding Energies of Hydrogen Bonds, *Journal of Medicine, Physiology and Biophysics*, **2**: 71-94.
- Ignatov, I.&Mosin,O.V. (2014) Modeling of Possible Conditions For Origin of First Organic Forms in Hot Mineral Water, *Journal of Medicine, Physiology and Biophysics*, **3**: 1-14.
- Ignatov, I.&Mosin,O.V. (2014) Mathematical Model of Interaction of Carbonaceous Fullerene Containing Mineral Shungite and Aluminosilicate Mineral Zeolite with Water, *Journal of Medicine, Physiology and Biophysics*, **3**: 15-29.
- Ignatov, I. &Mosin,O.V. (2014) Studying of Phototransformans of Light Signal by Photoreceptor Pigments – Rhodopsin, Iodopsin and Bacteriorhopsin and Additive Mixing of Colors, *Journal of Medicine, Physiology and Biophysics*, **3**: 30-47.
- Ignatov, I. &Mosin,O.V. (2014) Mathematical Models Describing Water Clusters as Interaction among Water Molecules. Distributions of Energies of Hydrogen Bonds, *Journal of Medicine, Physiology and Biophysics*, **3**: 48-70.
- Ignatov, I., Mosin, O. V.&Bauer, E. (2014) Carbonaceous Fullerene Mineral Shungite and Aluminosilicate Mineral Zeolite. Mathematical Model and Practical Application of Water Solution of Water Shungite and Zeolite, *Journal of Medicine, Physiology and Biophysics*, **4**: 27-44.
- Ignatov, I. &Mosin, O. V. (2014) Nature of Hydrogen Bonds in Liquids and Crystals. Ice Crystal Modifications and Their Physical Characteristics, *Journal of Medicine, Physiology and Biophysics*, **4**: 58-80.
- Ignatov, I.& Mosin, O. V. (2014) Research of Isotopic Effects of Deuterium in Cells of Microorganisms in the Presence of D₂O in IR Spectra in Hot Mineral Water for Origin of Life, *Journal of Medicine, Physiology and Biophysics*, **4**: 45-57.

- Ignatov, I. & Mosin, O. V. (2014) Visual Perception and Electromagnetic Conception for the Eyesight. Rhodopsin and Bacteriorhodopsin in Nano- and Biotechnologies, *Journal of Health, Medicine and Nursing*, **4**: 1-20.
- Ignatov, I., Mosin, O.V., Velikov, B., Bauer, E. & Tyminski, G. (2014) Research of Longevity Factors and Mountain Water as a Factor in Teteven, Yablanitsa and Ugarchin Municipalities, Lovech Region, Bulgaria, *Journal of Health, Medicine and Nursing*, **4**: 21-36.
- Ignatov, I. Mosin, O.V., Velikov, B. Bauer, E. & Tyminski, G. (2014) Mountain Water as Main Longevity Factor in Research of Phenomenon of Longevity in Mountain Areas in Bulgaria, *European Journal of Molecular Biotechnology*, **4** (2): 52-71.
- Ignatov, I., Mosin, O. V., Niggli, H., Drossinakis, Ch. & Stoyanov, Ch. (2014) Registration of Electromagnetic Waves Emitted the Human Body, *Journal of Medicine, Physiology and Biophysics*, **5**: 1-22.
- Ignatov, I. & Mosin, O. V. (2014) Modeling of Possible Processes for Origin of Life and Living Matter in Sea and Hot Mineral Water. Process of Formation of Stromatolites, *Journal of Medicine, Physiology and Biophysics*, **5**: 23-46.
- Ignatov, I., Mosin, O. V. (2014) Coronal Gas Discharge Effect in Modeling of Non-Equilibrium Conditions with Gas Electric Discharge Simulating Primary Atmosphere and Hydrosphere for Origin of Life and Living Matter, *Journal of Medicine, Physiology and Biophysics*, **5**: 47-70.
- Ignatov, I., Mosin, O. V. & Bauer, E. (2014) Mathematical Model and Practical Application of Water Solution of Shungite and Zeolite. Composition of Aluminosilicate Mineral Zeolite and Carbonaceous Fullerene Containing Mineral Shungite, *Journal of Health, Medicine and Nursing*, **5**: 12-29.
- Ignatov, I. & Mosin, O. V. (2014) Hydrogen Bonds among Molecules in Liquid and Solid State of Water. Modifications of Ice Crystals, *Journal of Health, Medicine and Nursing*, **5**: 56-79.
- Ignatov, I. & Mosin, O. V. (2014) Microorganisms in the Presence of D2O and IR Spectra in Hot Mineral Water with More Deuterium Atoms for Origin of Life and Living Matter, *Journal of Health, Medicine and Nursing*, **5**: 80-93.
- Ignatov, I. & Mosin, O.V. (2014) Color Coronal (Kirlian) Spectral Analysis in Modeling of Nonequilibrium Conditions with the Gas Electric Discharges, Simulating Primary Atmosphere, *Biomedical Radioelectronics*, **2**: 42-51. [in Russian]
- Ignatov, I., Mosin, O.V. & Stoyanov, Ch. (2014) Biophysical Fields. Color Coronal Spectral Analysis. Registration with Water Spectral Analysis. Biophoton Emission, *Journal of Medicine, Physiology and Biophysics*, **6**: 1-22.
- Ignatov, I. & Mosin, O.V. (2014) Hot Mineral Water with More Deuterium for Origin of Live and Living Matter. Process of Formation of Stromatolites, *Journal of Health, Medicine and Nursing*, **6**: 1-24.
- Ignatov, I. & Mosin, O.V. (2014) Origin of Life and Living Matter in Primary Atmosphere and Hydrosphere. Modeling of Non-equilibrium Electric Gas Discharge Conditions, *Journal of Health, Medicine and Nursing* **6**: 25-49
- Ignatov, I. & Mosin, O.V. (2014) Methods for Measurements of Water Spectrum. Differential Non-equilibrium Energy Spectrum Method (DNES), *Journal of Health, Medicine and Nursing* **6**: 50-72.

- Isaacs E.D., Shukla A., Platzman P.M, et all. (2000) Compton scattering evidence for covalency of the hydrogen bond in ice, *J. Phys. Chem. Solids*, 61, 403–406.
- Karadzov, S., Atanasov, A., Ivanova, E., Mosin, O. V. & Ignatov, I. (2014) Mathematical Models of Electrochemical Aqueous Sodium Chloride Solutions (Anolyte and Catolyte) as Types of Water. Study of the Effects of Anolyte on the Virus of Classical Swine Fever Virus, *Journal of Health, Medicine and Nursing*, 5: 30-55.
- Keutsch F. & Saykally R. (2011) Water Clusters: Untangling the Mysteries of the Liquid, One Molecule at a Time. *PNAS*, 98(19), 105330–10540.
- Liu K., Cruzan J.D. & Saykally R.J. (1996) Water clusters. *Science Magazine*, 271(5251), 929–933.
- Loboda O. & Goncharuk V. (2010) Theoretical study on icosahedral water clusters. *Chemical Physics Letters*, 484(4–6), 144–147.
- Luck W., Schiöberg D. & Ulrich S. (1980) Infrared Investigation of Water Structure in Desalination Membranes. *J. Chem. Soc. Faraday Trans.*, 2(76), 136–147.
- Maheshwary S., Patel N., Sathyamurthy N., Kulkarni A.D. & Gadre S.R. (2001) Structure and Stability of Water Clusters (H₂O)_n, n = 8-20: An Ab Initio Investigation. *J. Phys. Chem.*, 105, 10525–10537.
- Max, J.J. & Chapados, C. (2009) Isotope effects in liquid water by infrared spectroscopy. III. H₂O and D₂O spectra from 6000 to 0 cm⁻¹. *J. Chem. Phys.*, 131: 184505.
- Michaelides A. & Morgenstern K. (2007) Ice Nanoclusters at Hydrophobic Metal Surfaces. *Nat. Mat.*, 6, 597–599.
- Mosin O.V. & Ignatov I. (2011) Structure of water and physical reality. *Soznanije i physicheskaja realnost*, 10(1), 32–48 [in Russian].
- Mosin O.V. & Ignatov I. (2013a) The ice crystal riddles. *Soznanije i physicheskaja realnost*, 17(5), 21–31 [in Russian].
- Mosin O.V. & Ignatov I. (2013b) Structure of water. *Chimija*, 1, 12–32 [in Russian].
- Mosin, O.V. & Ignatov, I. (2014) Improved of Method for Isolation of Photochrome Transmembrane Protein Bacteriorhodopsin from Purple Membranes of Halobacterium Halobacterium Halobium, *Journal of Medicine, Physiology and Biophysics*, 3: 71-86.
- Mosin, O.V. & Ignatov, I. (2014) Photochrome Transmembrane Protein Bacteriorhodopsin from Purple Membranes of Halobacterium Halobium in Nano- and Biotechnologies, *Journal of Medicine, Physiology and Biophysics*, 4: 81-99.
- Mosin, O.V. & Ignatov, I. (2014) Facultative Methylophilic Bacterium Brevibacterium Methylicum in the Process of Preparation of Highly Deuterated Phenylalanine, Alanine, Valine and Leucine/Isoleucine, *Journal of Health, Medicine and Nursing*, 4: 37-56.
- Mosin, O.V. & Ignatov, I. (2014) Isolation of Photochrome Transmembrane Protein Bacteriorhodopsin from Purple Membranes of Halobacterium Halobacterium Halobium, New Method for Isolation, *Journal of Health, Medicine and Nursing*, 4: 57-73.
- Mosin, O.V. & Ignatov, I. (2014) Research from a Bacterium Bacillus Subtilis B-3157 by FAB Method of Biosynthetic Pathways of 2H-Labeled Purine Ribonucleoside Inosine, *Journal of Health, Medicine and Nursing*, 4: 74-93.

- Mosin, O. V. & Ignatov, I. (2014) Phenomenon of Biological Adaptation to Deuterium Oxide, *Journal of Medicine, Physiology and Biophysics*, **5**: 71-107.
- Mosin, O. V. & Ignatov, I. (2014) The Strain of Facultative Methylophilic Bacterium *Brevibacterium Methylicum* 5662 with RuMP Cycle of Carbon Assimilation for Biosynthesis of Deuterium Labeled Phenylalanine, *Journal of Medicine, Physiology and Biophysics*, **5**: 108-128.
- Mosin, O. V. & Ignatov, I. (2014) Photochrome Transmembrane Protein Bacteriorhodopsin from Purple Membrane of *Halobacterium Halobium*. Applications in Bio- and Nanotechnologies, *Journal of Health, Medicine and Nursing*, **5**: 94-112.
- Mosin, O.V. & Ignatov, I. (2014) Biosynthesis of Deuterium-labeled Transmembrane Protein Bacteriorhodopsin using a *Halobacterium Halobium*, *Journal of Medicine, Physiology and Biophysics*, **6**: 23-41.
- Mosin, O.V. & Ignatov, I. (2014) Photochrome Protein Bacteriorhodopsin Protein from Purple Membranes of *Halobacterium Halobium*. Application Applications in Bio- and Nanotechnologies, *Journal of Medicine, Physiology and Biophysics*, **6**: 42-60.
- Mosin, O.V. & Ignatov, I. (2014) Metabolism and Physiology of Methylophilic Microorganisms, *Journal of Medicine, Physiology and Biophysics*, **6**: 61-84.
- Mosin, O.V. & Ignatov, I. (2014) On the Nature of Biological Adaptation on Deuterium Oxide, *Journal of Medicine, Physiology and Biophysics*, **6**: 85-121.
- Mosin, O.V. & Ignatov, I. (2014) Phenomenon of Biological Adaptation to Heavy Water, *Journal of Health, Medicine and Nursing*, **6**: 73-110.
- Mosin, O.V. & Ignatov, I. (2014) Biosynthesis of Deuterium Labeled Phenylalanine. Facultative Methylophilic Bacterium *Brevibacterium Methylicum* #5662 with RuMP Cycle of Carbon Assimilation, *Journal of Health, Medicine and Nursing*, **6**: 111-132.
- Mosin, O.V. & Ignatov, I. (2014) Basic Concept of Magnetic Water Treatment, *European Journal of Molecular Biotechnology*, **4** (2): 72-85.
- Nemukhin A.B. (1996) The variety of clusters. *Russian Chemical Journal*, 40(2), 48–56 [in Russian].
- Ohno K., Okimura M., Akai N. & Katsumoto Y. (2005) The effect of cooperative hydrogen bonding on the OH stretching-band shift for water clusters studied by matrix-isolation infrared spectroscopy and density functional theory. *Phys. Chem. Chem. Phys.*, **7**, 3005–3014.
- Pasichnyk I. Everaers R. & Maggs A.C. (2008) Simulating van der Waals-interactions in water/hydrocarbon-based complex fluids. *J. Phys. Chem. B*, 112(6), 1761–1764.
- Pauling L. (ed.) (1960) *The Nature of the Chemical Bond and the Structure of Molecules and Crystals*. New York: Cornell University Press.
- Petkov V., Ren Y. & Suchomel M. (2012) Molecular arrangement in water: random but not quite. *J. Phys. Condens. Matter*, **24**, 155102.
- Pimentel G.C. & McClellan A.L. (1960) The hydrogen bond (L. Pauling, Ed). New York: Reinhold Publishing Co.
- Pople J.A. (1951) Molecular Association in Liquids: A Theory of the Structure of Water. *Proceedings of the Royal Society*, **205**, 1081–1089.
- Samoilov O.Ya. (1963) Studies of the structure of water. *Journal of Structural Chemistry*, **4**(4), 459–461.

- Sykes M. (2007) Simulations of RNA Base Pairs in a Nanodroplet Reveal Solvation-Dependent Stability. *PNAS*, 104(30), 12336–12340.
- Saykally R. (2005) Unified Description of Temperature-Dependent Hydrogen Bond Rearrangements in Liquid Water. *PNAS*, 102(40), 14171–14174.
- Timothy S. & Zwier S. (2004) Chemistry: the Structure of Protonated Water Clusters, *Science*. 304(5674), 1119–1120.
- Tokmachev A.M., Tchougreeff A.L. & Dronskowski R. (2010) Hydrogen-Bond Networks in Water Clusters (H₂O)₂₀: An Exhaustive Quantum-Chemical. *European Journal of Chemical Physics And Physical Chemistry*, 11(2), 384–388.
- Tokushima T., Harada Y., Takahashi O., et al. (2008) High resolution X-ray emission spectroscopy of liquid water: The observation of two structural motifs. *Chem. Phys. Lett.*, 460, 387–400.
- Tsai C.J. & Jordan K.D. (1993) Theoretical Study of the (H₂O)₆ Cluster. *Chem. Phys. Letters*, 213, 181–188.
- Wang F & Jordan K.D. (2003) A Parallel Tempering Monte Carlo Investigation of (H₂O)₆⁻. *J. Chem. Phys.*, 119, 11645–11653.
- Wernet Ph., Nordlund D., Bergmann U., et al. (2004) The structure of the first coordination shell in liquid water. *Science*, 304, 995–999.
- Walrafen, G.E. (1972) Raman and infrared spectral investigations of water structure. In *Water a Comprehensive Treatise*, F. Franks, Ed., Vol. 1, Plenum Press, New York, pp. 151–214.
- Zenin S.V. (1999) *Structured water condition as a basis for controlling the behavior of living systems and safety*. Diss. Doctor Biological Sciences. Moscow: Institute of Medical Biological Problems [in Russian].
- Yamaguchi Y, Yasutake N. & Nagaoka M. (2001) Theoretical prediction of proton chemical shift in supercritical water using gas-phase approximation. *Chem. Phys. Lett.*, 340, 129–136.
- Yukhnevitch, G.B. (1973) *Infrared spectroscopy of water*. Moscow, Nauka, 207 p. [in Russian].
- Zelmann, H.R. (1995) Temperature dependence of the optical constants for liquid H₂O and D₂O in the far IR region. *J. Mol. Struct.*, 350: 95–114.