

Length-Weight Relationship and Condition Factor of the West African Fiddler Crab (*Ucatangeri*) in MboRiver, Akwalbom State, Nigeria

*Okon, E. A. and Sikoki, F. D.

Department of Animal and Environmental Biology, Faculty of Biological Sciences, College of Natural and Applied Sciences, University of Port Harcourt, Nigeria

*Corresponding Author: nuelating@yahoo.com

Abstract

The length- weight relationship and condition factor of the West African fiddler crab (*Ucatangeri*) in Mbo river of Akwalbom State, Nigeria, was conducted for 12 consecutive months (April, 2012 – March, 2013). *Ucatangeri* exhibits sexual dimorphism with sex ratio of 1.2:1.0 which was significantly biased in favour of males. The length-weight relationship showed negative allometric growth, $b=1.6431$. With an average of 9.61, *Ucatangeri* had a very high condition factor which is an indication of the well-being of the species.

Keywords: Growth rate, Crab, Mbo.

1.0 Introduction

Crabs, together with other macro-benthic invertebrates, constitute the link between the unavailable nutrients in detritus and useful protein materials in fish and shellfish. Most benthic organisms feed on debris that settle on the bottom of the water and in turn serve as food for a wide range of fishes (Adebisi, 1989; Ajao and Fagade, 1990; Oke, 1990; Idowu and Ugwumba, 2005). They also accelerate the breakdown of decaying organic matter into simpler inorganic forms such as phosphates and nitrates (Gallep et. al., 1978). All forms of aquatic plants, which are the first link of several food chains existing in aquatic environment, can utilize the nutrients. These organisms therefore form a major link in the food chain as most fishes, birds and mammals depend directly or indirectly on them for their food supply (Barnes and Hughes, 1988).

The West African fiddler crab (*Ucatangeri*) is a species of fiddler crab that lives in the eastern Atlantic Ocean. It is the largest species in the genus *Uca*, with a carapace up to 50 millimetres (2.0 in) wide, and up to 25 mm (1.0 in) long. The males have one claw much larger than the other, which they use for communication. The carapace is violet to black, or sometimes yellowish in females, while the appendages are yellowish brown. The range of *Ucatangeri* extends from southern Portugal southwards to Angola; it is the most abundant crab in the Gambia (Hartnoll 1988).

Length-weight relationship is regarded as more suitable for assessing not only fish, but also crustacean (Sukumaran&Neelakantan, 1997; Tabash, 2001). The relationships between carapace length and weight of the crabs have many uses. They are often used to calculate the standing stock biomass, condition indices, analysis of ontogenetic changes and several other aspects of crustacean population dynamics (Atar&Seçer, 2003). In addition they are

used for the management of population. According to Lagler (1968), the relationship can be used to estimate the recovery of edible meat from crabs of various sizes. Body weight, total length and carapace length are the most frequently used dimensions in the study of crustaceans (Sukumaran&Neelakantan, 1997).

The earliest published estimate of crab population densities was that of Turnbull-Kemp (1960), who calculated the density of *Potamonautesperlatus* individuals greater than 17 mm CW at Inyanga (Zimbabwe) as 0.8–1.3 m⁻², with a combined wet weight of 5.5–13.6 g m⁻². This is low in comparison with an estimated density of the same species in the Buffalo River (Cape Province, South Africa) at 1.7–5.2 crabs m⁻² (based only on specimens >25 mm CW) and a (dry) biomass of 54–136 g m⁻² (Hill & O'Keefe 1992). In the work of Bello-Olusoji, et. al., (2009), there was a strong relationship ($r^2 = 0.81$) between length and weight of males and females and the entire crab population. The LWR was allometric for all crabs. Results also showed that there was a weak correlation ($r^2 = 0.36$) between fecundity and total body weight of the crabs. The length-weight distribution pattern did not show remarkable differences between species, sexes, and populations. Abowei and George (2009) examined the length - weight relationship and condition factor of *Callinectes amnicola* from Okpoka Creek in the Niger Delta area of Nigeria for a period of one year. Growth generally exhibited a negative allometry in all the sexes. Generally, the regression equations revealed high correlation in all the sexes. The monthly K values of the males ranged from 0.0984 and 0.1503 (mean = 0.1214). The estimated K value for both sexes combined was 0.1260. Similarly, estimated K values for the females ranged from 0.0889 – 0.1524. The overall mean K value for the females was 0.1229. The minimum K value (0.0984) for the males was observed in February. The female *C. amnicola* minimum K value (0.0889) and K_{max} (0.1580) values were obtained in November.

2.0 Materials and Method

2.1 **The Study Area:** Mbo River (Fig. 1) is a tributary of the Cross River. It lies between longitude 8°:00' – 8°:30'E and latitude 4°:30'–4°:45'N, and falls within the tropical rainforest belt with equatorial climate regime. The location of the study area is just north of the Equator and within the humid tropics and its proximity to the sea makes the area generally humid. The crabs were collected with the assistance of fishers using artisanal baited pots, traps and hands (manual) to pick and dislodge them from their nest. They were placed in clean polythene bags and taken to the laboratory for analysis.

2.2 **Determination of Length-Weight and Condition Factor:** The total body weight (BW) of the crab was measured to the nearest 0.1 g using a weighing balance (HANNA), while the carapace length (CL) was measured with a laboratory measuring board to the nearest 0.1 cm. The length-weight relationship was estimated using the equation:

$$W = aL^b$$

where W is the weight, *a* is the intercept, L is carapace length and *b* is the slope. The value of *a* and *b* were computed from the log transformed values of length and weight, i.e.

$$\text{Log } W = \log a + b \text{ Log } L,$$

where W = weight (g) of the crabs, L = horizontal carapace length (using the linear regression routine of Microsoft Office Excel in PC windows (2003). The parameters *a* (intercept) and *b* (slope) were estimated by linear regression based on logarithms;

$\log(W) = \log(a) + b \log(L)$. The significance of regression was assessed by analysis of variance (ANOVA). Equations expressing the length-weight relationships of *Ucatangeri* were calculated in relation to sex. For testing possible significant ($P > 0.01$) differences between the sexes Student's test was used for comparison of the two slopes. FORTRAN program Mature II computer technique was used to estimate the size at sexual maturity.

The Fulton's condition factor (cf) was calculated according to Bagenal (1978) with the formula;

$$K = 100W/L^3.$$

where K is the condition factor (cf), W is the total body weight (BW), L is the carapace length (CL) and 3 is a constant.

3.0 Results and Discussion

The length-weight relationship of *Ucatangeri* in Mbo river showed negative allometric growth, $b=1.6431$ (fig. 2). The variation between the carapace length and the mean weight was statistically highly significant ($p < 0.05$).

The growth coefficient (*b*) values have some implications and significant impacts on the well-being of fishes (including shell fish) and fishery. The negative allometric growth ($b < 3$) means the crabs were lighter than their body weights. Fish with high *b* ($b > 3$) values are heavy for their lengths, while those with low *b* are lighter (Wootton, 1998). Positive allometric growth is an indication of a crab's heaviness and by implication the crabs are heavier than their lengths. The change of *b* values depends primarily on the shape and fatness of the species, seasons or time of the years, temperature, salinity, food (quantity, quality and size), sex and stage of maturity (Sparre, 1992). The parameter *b*, unlike the parameter *a*, may vary seasonally, and even daily, and between habitats. The observed values of the regression coefficient (*b*) for *Ucatangeri* showed negative allometric growth, $b=1.6431$ which is less than 3. This is in agreement with the work of Abowei and George (2009). However, these results are contrary to what was observed by Akin-Oriola et al., (2005) who reported $b > 3$ (positive allometric growth) for *Callinectes pallidus* from Ojo Creek in Badagry, Lagos State.

The monthly variation in condition factor of *Ucatangeri* is shown in Table 1. The condition factor was nearly uniform between August 2012 and March 2013, and higher in the month of July 2012. The values of the condition factor ranged between 8.43 and 10.79 with a mean of 9.61 for combined sexes. This high condition factor indicates the well-being of the species.

The correlation factor (*r*) showed that carapace length is positively correlated with the weight of the species. Available literature lack information on correlation factor (*r*). Length –Weight relationship of *Ucatangeri* showed allometric growth rate of the species in Mboriver.

Like growth coefficients, condition factor (K) has impacts on the well-being of some aquatic organisms. It varies from species to species, and changes according to morphology, sex, age, reproductive state associated with gonadic maturity stages variations (Frederick & Thomas, 1987; Wootton, 1999). Variations in K may also be indicative of food abundance, adaptation to environment and gonadal development of fish (King, 1995). Low K means the fish are light for their lengths, an indication of low feeding intensity and spawning activity. High K value is an assumption of high feeding intensity and gradual increase in accumulated fat that also suggests preparation for a new reproductive period (Braga and Gennari-Filho, 1990). With an average of 9.61, *Ucatangeri* had a very high condition factor which is an indication of the well-being of the species. This is in agreement with reports of Lawson and Oloko (2013) as well as that of Arimoro and Idoro (2007).

4.0 Conclusion

The length-weight relationship and condition factor are comparable to works done by earlier researchers in other water bodies, implying that Mbo River was suitable for commercial crab production. A very high condition factor is an indication of the well-being of the species. There is a great need for a conscientious effort in the development of commercial crab culture in Nigeria.

REFERENCES

- Abolude, D.S., Davies, O.A., and Avong, D.W. (2009). Level of Heavy Metals in Freshwater Crab (*Cardisoma guanhumi*) Obtained from Ahmadu Bello University Reservoir, Zaria Nigeria. *International Journal of Animal and Veterinary Advances* 1(2): 54-58.
- Abowei, J.F.K. and George, A.D.I. (2009). A Study of the Length -Weight Relationship and Condition Factor of *Callinectes amnicola* (De Rochebrune, 1883) from Okpoka Creek, Niger Delta, Nigeria. *International Journal of Animal and Veterinary Advances* 1(2): 66-72, 2009
- Adamczewska, A. M., van Aardt, W. J. & Morris, S. (1997). Role of lungs and gills in an African fresh-water crab, *Potamonautes warreni* (Decapoda: Potamoidea), in gas exchange with water, with air, and during exercise. *Journal of Crustacean Biology* 17, 596-608.
- Adebisi, A.A., (1988). The physico-chemical hydrology of a tropical seasonal river Upper Ogun River. *Hydrologia*, 79: 159-165.
- Adegboye, D. (1983). On the non-existence of an indigenous species of crayfish on the continent of Africa. *Freshwater Crayfish* 5, 464-569.
- Ajana, A.M. (1996). Survey of Coastal and Brackishwater Shellfish Fisheries of Delta State. A Refils Report for the National Agricultural Research Project (NARP). Nigerian Institute for Oceanography and Marine Research (NIOMR), Lagos.
- Ajao, E.A. and S.O. Fagade, (1990). A study of sediment communities in Lagos Lagoon, Nigeria. *J. Oil Chem. Pollut.*, 7: 85-105.
- Ajayi, T.O. (1997). The status of the marine fisheries resources of the Gulf of Guinea. In: C.A. Ibe and S.G. Zabi (Eds.) *State of the Coastal and Marine Environment of the Gulf of Guinea*: 131-157
- Akin-Oriola, G., Anetekhai, M and Olowonirejuaro, K. (2005). Morphometric and Meristic Studies in Two Crabs: *Cardisoma armatum* and *Callinectes pallidus* from the Ojo creek in Badagry, Lagos. *Turkish Journal of Fisheries and Aquatic Sciences* 5: 85-89.
- Amadi, A.A. (1990). A comparative ecology of estuaries in Nigeria. *Hydrobiol.*, 208: 27-38
- APHA (1998): Standard methods for the examination of water and wastewater, 20th edition. American public health association, Washington D.C.
- Arimoro, F.O., and Idoro, B.O. (2007): Ecological Studies and Biology of *Callinectes amnicola* (Family: Portunidae) in the Lower Reaches of Warri River, Delta State, Nigeria. *World Journal of Zoology* 2 (2): 57-66,
- Avenant-Oldewage, A. (1993). Occurrence of *Temnocephalachaeropsis* on *Cherax tenuimanus* imported into South Africa, and notes on its infestation of an indigenous crab. *South African Journal of Science* 89, 427-428.
- Awosika, L., Osuntogun, N., Oyewo, E. and Awobamise, A. (2002). Nigeria National Report Phase 1: Integrated Problem Analysis. "Development and Protection of the Coastal and Marine Environment in Sub-Saharan Africa" GEF MSP Sub-Saharan Africa Project (GF/6010-0016): 10 – 11.
- Bagenal, T. (1978). *Method for Assessment of Fish Production in Freshwaters*. Blackwell Scientific Publication, Oxford. Pp 21-75.
- Barnes, R.D. and S. Hughes, (1988): *An introduction to marine ecology*. 2nd Edn., Blackwell Scientific Publications, UK., pp: 351.
- Bayliss, J. (2002). The East Usambara tree – hole crab (Brachyura: Potamoidea: Potamonautidae) – a striking example of crustacean adaptation in closed canopy forest, Tanzania. *African Journal of Ecology* 40, 26-34.
- Bello-Olusoji, O.A., Anifowose, O. J and Sodamola, M. Y. (2009) Length-Weight Relationships, Condition Factor and Fecundity of the West Africa Freshwater Crab, *Sudanonautes africanus* (Milne-Edwards 1883), in Western Nigeria. *West African Journal of Applied Ecology*, vol. 16, 65-75.
- Bertrand, J.-Y. (1979). Observations sur les crabes *Sudanonautes africanus* (Milne-Edwards 1869) et leurs terriers d'une savane de Lamto (Côte d'Ivoire). *Bulletin de la Société Zoologique de France* 104, 27-36.
- Beverton, R. J. H. and Holt, S. J. (1966). *Manual of methods for fish stock assessment*. Part II Tables of yield function. *FAO Fishery Biology Technical Paper*. (38): 67pp (ver. 1)
- Brandis, D., Storch, V. & Turkey, M. (2000). Taxonomy and zoogeography of the freshwater crabs of Europe, North Africa, and the Middle East. *Senckenbergiana Biologica* 80, 5-56.
- Butler, J. R. A. & Marshall, B. E. (1996). Resource use within the crab-eating guild of the upper Kairezi River,

- Zimbabwe. *Journal of Tropical Ecology* 12, 475-490.
- Cantrell, M. A. (1980). On the terrestrial behaviour of the freshwater crab *Potamonautescholoensis* (Chace). *Nyala* 6, 95-107.
- Carriol, R.-P. & Secretan, S. (1992). Présence dans le Bassin du Lac Albert (Ouganda), dès le Miocène supérieur, de *Potamonautes (Acanthothelphusa) niloticus* (Crustacea, Brachyura). *Comptes Rendus de l'Académie des Sciences, Paris, Série II* 314, 411-417.
- Connell, A. D. & Robertson, W. (1986). Recent records of megalopae of the crab *Varunalitterata* (Fabr.) entering Natal estuaries. *South African Journal of Zoology* 21, 184-185.
- Corace, R. G. III, Cumberlidge, N. & Garms, R. (2001). A new species of freshwater crab (Brachyura: Potamoidea: Potamonautidae) from the Ruwenzori region of western Uganda, East Africa. *Proceedings of the Biological Society of Washington* 114, 178-187.
- Cumberlidge, N. (1986). Ventilation of the branchial chambers in the amphibious West African freshwater crab, *Sudonautes (Convexonautes) aubryimonodi* (Bals, 1929) (Brachyura, Potamonautidae). *Hydrobiologia* 134, 53-65.
- Cumberlidge, N. (1991). The respiratory system of *Globonautes macropus* (Rathbun, 1898), a terrestrial freshwater crab from Liberia (Gecarcinucoidea, Gecarcinucidae). *Crustaceana* 61, 69-80.
- Cumberlidge, N. & Sachs, R. (1991). Ecology, distribution, and growth in *Globonautes macropus* (Rathbun, 1898), a tree-living freshwater crab from the rainforests of Liberia (Gecarcinucoidea, Gecarcinucidae). *Crustaceana* 61, 55-68.
- Cumberlidge, N. (1994). *Louisea*, a new genus of freshwater crab (Brachyura: Potamoidea: Potamonautidae) for *Globonautes macropus edaensis* Bott, 1969 from Cameroon. *Proceedings of the Biological Society of Washington* 107, 12-131.
- Cumberlidge, N. (1996). A taxonomic revision of fresh-water crabs (Brachyura, Potamoidea, Gecarcinucidae) from the upper Guinea forest of West Africa. *Crustaceana* 69, 681-695.
- Cumberlidge, N. (1999). *The freshwater crabs of West Africa. Family Potamonautidae*. Paris, IRD. Collection Faune et Flore Tropicales No. 236.
- Cumberlidge, N., Sternberg, R. V., Bill, I. R. & Martin, H. (1999). A revision of the genus *Platythelphusa* A. Milne-Edwards, 1887 from Lake Tanganyika, East Africa (Decapoda: Potamoidea: Platythelphusidae). *Journal of Natural History* 33, 1487-1512.
- Cumberlidge, N. & Sternberg, R. V. (2002). The freshwater crabs of Madagascar (Crustacea, Decapoda, Potamoidea). *Zoosystema* 24, 41-79.
- Cumberlidge, N. & Wranik, W. (2002). A new genus and new species of freshwater crab (Potamoidea, Potamidae) from Socotra Island, Yemen. *Journal of Natural History* 36, 51-64.
- Cumberlidge, N., Boyko, C. B. & Harvey, A. W. (2002). A new genus and species of freshwater crab (Crustacea, Decapoda, Potamoidea) from northern Madagascar, and a second new species associated with *Pandanus* leaf axils. *Journal of Natural History* 36, 65-77.
- Cumberlidge, N. & Reed, S. K. (2004). *Erimetopus vandenbrandeni* (Bals, 1936) n. comb., with notes on the taxonomy of the genus *Erimetopus* Rathbun, 1894 (Brachyura: Potamoidea: Potamonautidae) from Central Africa. *Zootaxa*.
- Cumberlidge, N. & Vannini, M. (2004). A new species of tree-hole living freshwater crab (Brachyura: Potamoidea: Potamonautidae) from coastal East Africa. *Journal of Natural History*.
- Daniels, S. R. (2001). Allometric growth, handedness, and morphological variation in *Potamonautes warreni* (Calman, 1918) (Decapoda, Brachyura, Potamonautidae) with a redescription of the species. *Crustaceana* 74, 237-253.
- Daniels, S. R., Stewart, B. A. & Cook, P. A. (2002). Congruent patterns of genetic variation in a burrowing freshwater crab revealed by allozymes and mt DNA sequence analysis. *Hydrobiologia* 468, 171-179.
- Daniels, S. R., Stewart, B. A., Gouws, G., Cunningham, M. & Mathee, C. A. (2002b). Phylogenetic relationships of the southern African freshwater crab fauna (Decapoda: Potamonautidae: *Potamonautes*) derived from multiple data sets reveal biogeographic patterning. *Molecular Phylogenetics and Evolution* 25, 511-523.
- Disney, R. H. L. (1971). Notes on *Simulium ovazzae* Grenier & Mouchet (Diptera: Simuliidae) and river crabs (Malacostraca: Potamidae) and their association. *Journal of Natural History* 5, 677-689.
- Dobson, M., Magana, A., Mathooko, J. M. & Ndegwa, F. K. (2002). Detritivores in Kenyan highland streams: more evidence for the paucity of shredders in the tropics? *Freshwater Biology* 47, 909-919.
- Dobson, M. (2004). Freshwater Crabs in Africa. *Freshwater Forum*, 21: 3-26.
- Dominey, W. J. & Snyder, A. M. (1988). Kleptoparasitism of freshwater crabs by cichlid fishes endemic to Lake Barombi Mbo, Cameroon, West Africa. *Environmental Biology of Fishes* 22, 155-160.
- Ekwu, A. O. and Sikoki, F. D. (2006). Phytoplankton diversity in the Cross River estuary of Nigeria. *J. Appl. Sci. Environ. Mgt.* Vol 10(1): 89-95

- Ewa-oboho, I. and Abby-kalio, N. J. (1993). Theoretical Basis of Allometric Relationships in Juvenile Brachyura: Data from a West African Mangrove Swamp Crab Population
- Flower, S. S. (1931). Notes on freshwater crabs in Egypt, Sinai, and the Sudan. *Proceedings of the Zoological Society of London* 1931, 729-735.
- Francis, A. (2003). Studies on the Ichthyofauna of the Andoni River System in the Niger Delta of Nigeria. A PhD Thesis submitted to the Department of Animal and Environmental Biology, University of Port Harcourt, Pp13-39.
- Francis, A., Sikoki, F. D. and Ansa E. J. (2007). physico-chemical parameters of the Andoni river system-Niger Delta, Nigeria. *Journal of fisheries international* 2(1) pp.27-31.
- Gallep, G.W., J.F. Kitchell and S.M. Bartell, (1978): Phosphorus release from lake sediments as affected by chironomid. *Ver intervere for Limnologic*, 20:458-465
- Gayanilo Jr., F. C., Sparre, P., and Pauly, D. (1996). FAO-ICLARM Stock Assessment Tools (FISAT), FAO, Rome. 126pp (+3 diskettes).
- Gayanilo Jr., F. C. and Pauly, D. (1997). FAO-ICLARM Stock Assessment Tools (FISAT). FAO Computerised Information Series (fisheries) No. 8, pp262
- George, A.D.I., J.F.N. Abowei, and J. F. Alfred-Ockiya (2010): The Distribution, Abundance and Seasonality of Benthic Macro Invertebrate in Okpoka Creek Sediments, Niger Delta, Nigeria. *Res. J. Appl. Sci. Eng. Technol.*, 2(1): 11-18, 2010
- Gouws, G. & Stewart, B. A. (2001). Potamonautidriver crabs (Decapoda, Brachyura, Potamonautidae) of KwaZulu-Natal, South Africa. *Water SA* 27, 85-98.
- Harper, D. M., Mavuti, K. M. & Muchiri Mucui, S. (1990). Ecology and management of Lake Naivasha, Kenya, in relation to climatic change, alien species introductions and agricultural development. *Environmental Conservation* 17, 328-336.
- Hill, M. P. & O'Keefe, J. H. (1992). Some aspects of the ecology of the freshwater crab (*Potamonautesperlatus* Milne Edwards) in the upper reaches of the Buffalo River, eastern Cape Province, South Africa. *South African Journal of Aquatic Sciences* 18, 42-50.
- Holly Dublin, Jean-Christophe Collen, Craig Hilton-Taylor, and Simon N. Stuart (2009) Wildlife in a Changing World: An Analysis of the 2008 IUCN Red List of Threatened Species. *IUCN*. pp. vii–viii. ISBN 978-2-8317-1063-1
- Hussein, M. A. & Obuid, A. H. (1993). Freshwater crabs in Egypt and a redescription of *Potamonautes niloticus*. *Zoology in the Middle East* 8, 113-119.
- Hynes, H. B. N., Williams, T. R. & Kershaw, W. E. (1961). Freshwater crabs and *Simulium neavei* in East Africa I. – Preliminary observations made on the slopes of Mount Elgon in December, 1960, and January, 1961. *Annals of Tropical Medicine and Parasitology* 55, 197-201.
- Idowu, E.O. and A.A.A. Ugwumba, (2005): Physical, chemical and benthic faunal characteristics of a Southern Nigeria Reservoir. *The Zoologist*, 3: 15-25.
- Ikomi, R.B., F.O. Arimoro and O.K. Odihirin, (2005): Composition, distribution and abundance of macroinvertebrates of the upper reaches of River Ethiopia Delta State, Nigeria. *The Zoologist*, 3: 68-81
- King, J. M. (1983). Abundance, biomass and density of benthic macroinvertebrates in a western Cape River, south Africa. *Transactions of the Royal Society of South Africa* 45, 11-34.
- King, R.P. (1995). Fisheries Biology, Assessment and Management. *Fishing News Books, Surrey*, pp: 336.
- Lawal-Are, A. O. (2010). Reproductive Biology of the Blue Crab, *Callinectes amnicola* (De Rocheburne) in the Lagos Lagoon, Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences* 10: 1-7
- Lawson, E. O., and Oloko, R.T. (2013). Growth patterns, Sex ratios and Fecundity estimates in Blue Crab (*Callinectes amnicola*) from Yewa River, Southwest Nigeria.
- Le Cren, E. D. (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the Perch (*Perca fluviatilis*) *J. Ani. Ecol.*, 20:201-219.
- Litulo, C. (2004). Fecundity of the Pantropical Fiddler Crab *Uca annulipes* (H. Milne Edwards, 1837) (Brachyura: Ocypodidae) at Costa do Sol Mangrove, Maputo Bay, Southern Mozambique. *Western Indian Ocean J. Mar. Sci. Vol. 3, No. 1*, pp. 87–91.
- McMahon, J. P., Highton, R. B. & Goiny, H. (1958). The eradication of *Simulium neavei* from Kenya. *Bulletin of the World Health Organisation* 19, 75-107
- Mikkola, H. (1996). Alien freshwater crustacean and indigenous mollusc species with aquaculture potential in eastern and southern Africa. *Southern African Journal of Aquatic Sciences* 22, 90-99.
- Moore, J. E. S. (1903). *The Tanganyika problem. An account of the researches concerning the existence of marine animals in Central Africa*. Hurst & Blackett Ltd, London
- Morris S. F. (1976). A new fossil freshwater crab from the Ngorora Formation (Miocene) of Kenya. *Bulletin of the British Museum (Natural History) Geology* 27, 295-300.
- Morris, S. & van Aardt, W. J. (1998). Salt and water relations, and nitrogen excretion, in the amphibious African

- freshwater crab *Potamonautesarreni* in water and in air. *Journal of Experimental Biology* 201, 883-893.
- Moses, B. S. (1987): the influence of flood regime on fish catch and fish communities in the Cross River floodplain ecosystem, Nigeria. *Environ. Biol. Fish* 18:51-65.
- Odiete, W.O., (1999): Environmental physiology of animals and Pollution. Diversified Resources, Lagos, Nigeria, pp: 220-246.
- Oduro, W., Ellis, W.O., Oduro, I. and Tetteh, D.(2001). Nutritional quality of selected Ghanaian crab species. *J. Ghana Sci. Ass.*, 3: 37-40.
- Oesterling, M. J.(1990). Shellfish – Crustaceans. In: Martin, R. E. and Flick, G. J. (Eds) The Seafood Industry. Van Nostrand Reinhold Publ., New York: 88-102
- Oke, O.O., (1990): Limnology and macrobenthos of owenareservoir. M.Phil. Thesis, University of Ibadan, Ibadan, Nigeria. pp: 96.
- Okon, E.A., Sikoki, F.D. and Komi, G.W. (2013). Physico-Chemical parameters of the Mbo River, Niger Delta, Nigeria. Proceeding of International Conference on Sustainable Development. Vol.12 No.3, pp 138-144 July 15-18, 2013, Chinua Achebe Arts Theatre, University of Calabar, Nigeria. International Research and Development Institute. www.irdi.org.ng
- Owhonda, N. K., Akinrotimi, O. A., Ansa E. J., Edun, O. M., Anyanwu, P. E., Opara, J. Y., and Onunkwo, D. O. (2007). Wet season variation in some physiochemical parameters of brackish water fish ponds and main channels in Buguma, Rivers State, Nigeria. *Journal of Fisheries International* 2(3)255-259.
- Ng, P. K. L., Števíć, Z. & Pretzmann, G. (1995). A revision of the family Deckeniidae Ortmann, 1897 (Crustacea: Decapoda: Brachyura: Potamoidea), with description of a new genus (Gecarcinucidae: Gecarcinucoidea) from the Seychelles, Indian Ocean. *Journal of Natural History* 29, 581-600.
- Okafor, F. C. (1988). The ecology of *Sudanonautes (Sudanonautes) africanus* (H. Milne-Edwards) (Crustacea: Decapoda) in southeastern Nigeria. *Tropical Ecology* 29, 89-97.
- Ollivier, G., Boussinesq, M., Albaret, J. L., Cumberlidge, N., Farhati, K., Chipaux, J.P. & Baysadde-Dufour, C. (1995). Etude épidémiologique d'une distomatose à *Paragonimus* sp. au sud- Cameroun. *Bulletin de la Société de Pathologie Exotique* 88, 164- 169.
- Pauly, D., (1980). A selection of simple methods for the assessment of tropical fish stocks. *FAO Fisheries Circular*, 729, pp: 54.
- Pauly, D., (1983). Some simple methods for the assessment of tropical stocks. *FAO Fish. Tech. Pap.* 234, pp: 52.
- Pauly, D. (1986). On improving the operation and use of the ELEFAN programs. Part III: Correcting length - frequency data for effects of gear selection and /or incomplete re-enactment. *ICLARM Fishbyte*, 4(2):11-13.
- Pauly, D. (1990). Length-converted catch curves and the seasonal growth of fishes. *Fishbyte* 8(3): 33-38.
- Pauly, D. (1993). Editorial: Fishbyte section, NAGA, *The ICLARM Quarterly* 16(2-3) :26.
- Pauly D. and Munro J.L. (1984). Once more on the comparison of growth in fish and invertebrate. *ICLARM, Fishbyte*, 2(1): 21
- Pinheiro', M. A. A.; Baveloni, M. D. and Terceiro, O. L. (2003). Fecundity of the mangrove crab *Ucidescordatus* (Linnaeus, 1763) (Brachyura, Ocypodidae). *Invertebrate Reproduction and Development*, 43:1 (2003) 19-26
- Powell, D. G. (1979). Estimation of mortality and growth parameters from the length-frequency in the catch. *Rapp. P. -v. Reun CIEM* 175: 167-169.
- Purves, M. G., Kruuk, H. & Nel, J. A. J. (1994). Crabs *Potamonautes perlatus* in the diet of otter *Aonyx capensis* and water mongoose *Atilax paludinosus* in a freshwater habitat in South Africa. *Zeitschrift für Säugetierkunde* 59, 332-341.
- Rathbun, M. J. (1921). The brachyuran crabs collected by the American Museum Congo expedition 1909-1915. *Bulletin of the American Museum of Natural History* 43, 379-474.
- Ravera, O., (1998): Utility and limits of biological and chemical monitoring of the aquatic environment. *Annal. Dichim.*, 88: 909- 913.
- Ravera, O., (2000): Ecological monitoring for water body management. Proceedings of monitoring Tailormade III. International Workshop on Information for Sustainable Water Management, pp: 157-167.
- Ravi, R., Manisseri, M. K. and Kuriakose, S. (2008). Relationship between morphometric characteristics and fecundity of *Portunus pelagicus* (Linnaeus, 1758). *Morphometri. Jc. cMhaarr. acBtieorli. s tAicss. aInnddiafe, c5u0n d(i2ty) : 217 - 220, July - December 2008*
- Raybould, J. N. (1969). Studies on the immature stages of the *Simulium neavei* Roubaud complex and their associated crabs in the Eastern Usambara Mountains in Tanzania. I. Investigations in rivers and large streams. *Annals of Tropical Medicine and Parasitology* 63, 269-287.
- Raybould, J. N. & Mhiddin, H. K. (1978). Studies on the immature stages of the *Simulium neavei* Roubaud complex and their associated crabs in the Eastern Usambara Mountains in Tanzania. III. Investigations

- on development and survival and their relevance to control. *Annals of Tropical Medicine and Parasitology* 72, 177- 187.
- Raybould, J. N., Grunewald, J. & Mhiddin, H. K. (1978). Studies on the immature stages of the *Simulium neavei* Roubaud complex and their associated crabs in the Eastern Usambara Mountains in Tanzania. IV. Observations on the crabs and their attached larvae under exceptionally conditions. *Annals of Tropical Medicine and Parasitology* 72, 189- 194.
- Sachs, R. & Cumberlidge, N. (1991). The dwarf river crab *Liberonautes latidactylus nanoides* Cumberlidge & Sachs, 1989, from Liberia – a new second intermediate host of *Paragonimus uterobilateralis*. *Tropical Medicine and Parasitology* 42, 73-74.
- Sanders, M. J., Du Preez, H. H. & van Vuren, J. H. J. (1999). Monitoring cadmium and zinc contamination in freshwater systems with the use of the freshwater crab, *Potamonautes warreni*. *Water SA* 25, 91-98.
- Schuwert, P.-M.M., Lewis, J. W. & Jones, P. (2001). The potential use of the South African river crab, *Potamonautes warreni*, as a bioindicator species for heavy metal contamination. *Ecotoxicology* 10, 159-166.
- Shaw, J. (1959). Salt and water balance in the East African fresh-water crab, *Potamonniloticus* (M. Edw.). *Journal of Experimental Biology* 36, 157-176.
- Sikoki F.D. (2013). Fishes in Nigerian waters: No place to hide: Inaugural lecture series No. 100. University of Port Harcourt press Ltd. Pp 36-42.
- Somers, M. J. & Nel, J. A. J. (1998). Dominance and population structure of freshwater crabs (*Potamonautes perlatus* Milne Edwards). *South African Journal of Zoology* 33, 31-36.
- Sparre, P., E. Ursin and S.C. Venema, (1989). Introduction to tropical fish stock assessment, Part 1 Manual. *FAO Fisheries Technical Paper* No. 306. 1 FAO Rome, pp: 337.
- Turnbull-Kemp, P. St. J. (1960). Quantitative estimations of populations of the river crab *Potamon (Potamonautes) perlatus* (M. Edw.), in Rhodesian trout streams. *Nature* 185, 481.
- Udoh, J. P. and Nlewadim, A.A. (2011). Population characteristics of the swimming crab *Callinectes amnicola* De Rocheburne, 1883 (Crustacea, Brachyura, Portunidae) in the Qua Iboe River estuary, Nigeria. *AACL Bioflux*, 2011, Volume 4, Issue 3.
- Udoh, J. P., Nlewadim, A. A. and Ofor, C. (2009). Maturity estimation in male swimming brachyuran crab, *Callinectes amnicola* (De Rocheburne, 1883) (DECAPODA, PORTUNIDAE) in the Imo river estuary, Nigeria
- Udoiong, O. M., (2010): Threats to species of epibenthic Macro-fauna of some mangrove wetlands in Eastern Obolo, Nigeria. *World Journal of Applied Science and Technology* Vol.2. No. 2 (2010). 232 - 244
- Udonsi, J.K. (1987). Endemic *Paragonimus* infection in Upper Igwu Basin, Nigeria: a preliminary report on a renewed outbreak. *Annals of Tropical Medicine and Parasitology* 81, 57-62.
- van Eeden, P. H. & Schoonbee, H. J. (1991). Bio-accumulation of heavy metals by the fresh-water crab *Potamonautes warreni* from a polluted wetland. *South African Journal of Wildlife Research* 21, 103-108.
- van Someren, V. D. & McMahon, J. (1950). Phoretic association between *Afronurus* and *Simulium* species, and the discovery of the early stages of *Simulium neavei* on freshwater crabs. *Nature* 166, 350-351.
- Voelker J. & Sachs, R. (1977). Über die Verbreitung von Lungeneiern (*Paragonimus africanus* und *P. uterobilateralis*) in West-Kamerun und Ost-Nigeria auf Grund von Untersuchungen an Süßwasserkrabben auf Befehl mit Metazerkarien. *Tropenmedizin und Parasitologie* 28, 129- 133.
- Voelker, J., Sachs, R., Volkmer, K. L., & Braband, H. (1975). On the epidemiology of paragonimiasis in man and in animals in Nigeria, West Africa. *Veterinary Medical Review* 1-2, 161-175.
- West, K., Cohen, A. & Baron, M. (1991). Morphology and behaviour of crabs and gastropods from Lake Tanganyika, Africa: implications for lacustrine predator-prey coevolution. *Evolution* 45, 589-607.
- Wetherall, J. A. (1986). A new method for estimating growth and mortality parameters from length-frequency data. *Fishbite* 4(1): 12- 14
- Williams, T. R. (1961). The diet of freshwater crabs associated with *Simulium neavei* in East Africa. I. – Crabs from West and East Uganda collected by the Cambridge East African Expedition, 1959. *Annals of Tropical Medicine and Parasitology* 55, 128-131.
- Williams, T. R. (1962). The diet of freshwater crabs associated with *Simulium neavei* in East Africa. II. – The diet of *Potamonautes berardi* from Mount Elgon, Uganda. *Annals of Tropical Medicine and Parasitology* 56, 362-367.
- Williams, T. R., Hynes, H. B. N. & Kershaw, W. E. (1964). Freshwater crabs and *Simulium neavei* in East Africa II. – Further observations made during a second visit to East Africa in February–April 1962 (the dry season). *Annals of Tropical Medicine and Parasitology* 58, 159-167.
- Williams, T. R. (1965). The diet of freshwater crabs associated with *Simulium neavei* in East Africa. III. – The diet of *Potamonautes niloticus* and of an unidentified crab species from Mount Elgon, Uganda.

Annals of Tropical Medicine and Parasitology 59, 47-50.


Williams, T. R. (1968). The taxonomy of the East African river-crabs and their association with the *Simulium neavei* complex. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 62, 29-34.

Williams, T. R. (1991). Freshwater crabs and *Simulium neavei* in East Africa III. – Morphological variation in *Potamonautes sloveni* (Decapoda: Potamidae). *Annals of Tropical Medicine and Parasitology* 85, 181-188.

Yeo, D. C. J., Peter K. L., Neil Cumberlidge, Célio Magalhães, Savel

R. Daniels & Martha R. Campos (2008). *Global diversity of crabs (Crustacea: Decapoda: Brachyura) in freshwater*. In E. V. Balian, C. Lévêque, H. Segers & K. Martens. "Freshwater Animal Diversity Assessment". *Hydrobiologia* 595 (1): 275-286

Zabbey, N., (2002): An ecological survey of benthic Macro-invertebrates of Woji Creek, off the Bonny River System Rivers State. M.Sc. Thesis, University of Port Harcourt, pp: 102



**Fig. 1: Map of Mbo River showing the proposed sampling station
(Insert: Map of Nigeria showing the location of the study area)**




Fig. 2: Length-Weight Relationship of *Ucatangerin* Mbo River

Table 1 : Monthly Variation in Condition Factor of *Ucatangerin* Mbo River

Month	Mean Weight(g)	Mean length(cm)	Condition factor (K)
April 2012	2.6153	3.0633	9.0982
May 2012	3.4336	3.3733	8.9451
June 2012	3.0657	3.1867	9.4734
July 2012	3.5593	3.2067	10.7942
August 2012	2.9117	3.2500	8.4819
September 2012	2.9297	3.2117	8.5344
October 2012	2.9370	3.2300	8.7156
November 2012	2.8580	3.2367	8.4286
December 2012	3.0203	3.2667	8.6641
January 2013	3.1550	3.2800	8.9408
February 2013	3.1328	3.2767	8.9048
March 2013	3.2473	3.3333	8.7679

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:
<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

