

Non-Linear Effort dynamics for Harvesting in a Predator- Prey System

Reenu Rani* Sunita Gakkhar

Department of Mathematics, Indian Institute of Technology, Roorkee, Roorkee-247667, Uttarakhand, India

* E-mail of the corresponding author: reenuitr2012@gmail.com

Abstract

In this paper, a non-linear harvesting of prey is considered in a prey-predator system. The predator is considered to be of modified Leslie- Gower type. The effort is taken as dynamic variable. The steady states of the system are determined and the dynamical behavior of the system for its all steady states is discussed under certain conditions. Necessary condition for global stability of the system is analyzed at the positive interior equilibrium point. Numerical simulations are carried out to explore the dynamics of the system for the suitable choice of parameters.

Keywords: Modified Leslie-Gower predation, Nonlinear Harvesting, stability, Numerical Simulations.

1. Introduction

The ecological non-linear models of interacting populations have been studied extensively by many authors .In recent years, the need for more food/ resources is growing at large scale, which has resulted in over exploitation of several biological resources. Even there is a global concern to protect the ecosystem from exploitation at large scale. Therefore, there is a need for a sustainable development policy in various spheres of human activity to protect ecosystems. In particular, bio-economic modeling is concerned with scientific management of the exploitation of renewable resources like fisheries and forestry. Hence harvesting of ecosystems has been of interest to economists and ecologists. A harvesting policy refers to the management of biological resources by systematically controlling the period, intensity and type of harvesting. The primary objective here is to maximize productivity without depleting or driving the stocks to extinction. In recent years, many works on optimal management of renewable resources are done. An excellent introduction to optimal management of renewable resources is given by Clark [3] and also policies related to bionomic exploitations of renewable resources are discussed by him. Harvesting has a strong impact on the dynamics of biological resources. The severity of the impact depends on the nature of the implementation of harvesting strategy. Basically three types of harvesting strategies are reported in literature (i) constant harvesting (ii) proportional harvesting, $h(x) = qEx$ and (iii) nonlinear harvesting, $H(x) = qEx/(m1E + m2x)$ (Holling type-II. Many researchers have analyzed mathematical models using non-linear harvesting by considering different type of growth depending upon the species and their interactions [5, 6, 12] etc.

Zhu and Lan [16] studied Leslie- Gower model with constant harvesting in prey. Though there are numerous works on predator- prey system incorporating the harvesting of the species. Leslie Gower predator prey model and modified Leslie Gower predator-prey under the assumption of the nonlinear harvesting on prey species has been studied by Gupta and Banerjee et al. [5]. But in this paper, we consider a dynamical reaction model of modified Leslie Gower predator- prey incorporating the non-linear harvesting on prey where effort is taken as a dynamic variable. The present paper deals with a dynamic reaction model in the case of a prey-predator type fishery system, while the model we study here is especially based on a modified version of the Leslie Gower scheme, where only the prey species is subjected to non- linear harvesting. The main aim of this paper is to examine the effect of harvesting on such kind of prey - predator system with the effort dynamic which can give the best possible benefit through harvesting to the society while preventing extinction of the species.

2. The Mathematical Model

Let $x(t)$, $y(t)$ and $E(t)$ are the densities of prey, predator population and the harvesting effort at a time 't'. The Holling type- II functional response and non-linear harvesting is considered for a logistically growing prey species and predator is assumed to be modified Leslie- Gower type. The mathematical model for the dynamics of the system governed by the following system of differential equations:

$$(2.1) \quad \begin{aligned} \frac{dx}{dt} &= rx \left(1 - \frac{x}{k}\right) - \frac{\alpha xy}{\alpha + x} - \frac{qEx}{m_1 E + m_2 x} = xf(x, y, E), \\ \frac{dy}{dt} &= sy \left(1 - \frac{\beta y}{a + x}\right) = yg(x, y), \\ \frac{dE}{dt} &= \eta E \left(\frac{qpx}{m_1 E + m_2 x} - c\right) = Eh(x, E), \end{aligned}$$

$$x(0) > 0, y(0) > 0, E(0) > 0.$$

The constants $r, s, k, \alpha, \beta, p, c,$ and η are the parameters assuming only positive values. The parameters $r, s, k, \alpha, \beta, p, c,$ and η represents intrinsic growth rate of prey population, growth rate for the predator, environmental carrying capacity, en-counter rate of predator to prey, maximum rate of the reduction of predator population, price and cost per unit mass and stiffness parameter, respectively.

3. Uniform Boundedness

Theorem: All the solutions of the system (2.1) which start in the region R_3 are uniformly bounded.

Proof: let us consider a function $\phi(t)$ such that

$$\begin{aligned} \phi(t) &= x(t) + y(t) + \frac{1}{\eta p} E(t) \\ \frac{d\phi(t)}{dt} &= x'(t) + y'(t) + \frac{1}{\eta p} E'(t) \\ &= rx \left(1 - \frac{x}{k}\right) - \frac{\alpha xy}{\alpha + x} - \frac{qEx}{m_1 E + m_2 x} + sy - \frac{s\beta y^2}{a + x} + \frac{qpx}{m_1 E + m_2 x} - \frac{cE}{\eta p} \\ \frac{d\phi(t)}{dt} &\leq \left(rx - \frac{rx^2}{k} \right) + sy' - \frac{\beta sy^2}{a + k} - \frac{cE}{\eta p} \end{aligned}$$

Introducing, a positive constant N and rewrite the above equation, we get

$$\frac{d\phi(t)}{dt} + N\phi(t) \leq \left((r + N)x - \frac{rx^2}{k} \right) + \left((s + N)y - \frac{\beta sy^2}{a + k} \right) - \frac{(c - N)E}{\eta p}$$

Let us $c > N$, Then we have:

$$\frac{d\phi(t)}{dt} + N\phi(t) \leq -\frac{r}{k} \left(x - \frac{k(r + N)}{2r} \right)^2 - \frac{s\beta}{a + k} \left(y - \frac{(s + N)(k + a)}{2s\beta} \right)^2 + M$$

$$\frac{d\phi(t)}{dt} + N\phi(t) \leq M; \quad M = \frac{k^2(r + N)^2}{4r^2} + \frac{(s + N)^2(k + a)^2}{4s^2\beta^2}$$

Solving above inequality, we get

$$0 < \lim_{t \rightarrow \infty} \phi(t) \leq \frac{M}{N}$$

All the solutions of the system (2.1) which start in the region R_3 are confined in the region: This proves the result.

$$R = \left\{ (x, y, E) \in R; 0 < x(t) + y(t) + \frac{1}{\eta p} E(t) \leq M + \phi, \phi > 0 \right\}$$

4. Existence of Equilibrium Points

For the model (2.1), there exists six non-negative equilibrium points which are given below:

1. $P_0(0, 0, 0)$ is a trivial equilibrium point.
2. $P_1(k, 0, 0)$ is the axial equilibrium point on x-axis.
3. $P_2(0, a/\beta, 0)$ is the axial point on y-axis in the absent of prey and harvesting.
4. $P_3(\bar{x}, \bar{y}, \bar{E})$ boundary equilibrium point on xy-plane in the absence of harvesting effort. The

equilibrium level \bar{x} and \bar{y} are the solution of the following equations:

$$r\bar{x}\left(1 - \frac{\bar{x}}{k}\right) - \frac{\alpha\bar{x}\bar{y}}{\alpha + \bar{x}} = 0; \left(1 - \frac{\beta\bar{y}}{a + \bar{x}}\right) = 0$$

Solving, we get, $\bar{x} = r\left(1 - \frac{\alpha}{r\beta}\right)$ $\bar{y} = \frac{a + \bar{x}}{\beta}$

Point $P_3(\bar{x}, \bar{y}, 0)$ exists for $r\beta > \alpha$.

5. $P_4(\hat{x}, 0, \hat{E})$ is boundary equilibrium point on xy- plane in the absence of harvesting effort. These equilibrium levels are the solution of the following equations:

$$r\hat{x}\left(1 - \frac{\hat{x}}{k}\right) - \frac{\alpha\hat{x}\hat{y}}{\alpha + \hat{x}} - \frac{q\hat{E}\hat{x}}{m_1\hat{E} + m_2\hat{x}} = 0$$

$$\frac{qp\hat{x}}{m_1\hat{E} + m_2\hat{x}} - c = 0$$

Solving, we get the values as: $\hat{x} = k\left(1 - \frac{qL}{r(m_1L + m_2)}\right)$ and $\hat{E} = L\hat{x}$; where $L = \frac{pq - cm_2}{cm_1}$

$P_4(\hat{x}, 0, \hat{E})$ is positive if $r(m_1L + m_2) > qL$ and $c < \frac{pq}{m_2}$

6. $P_5(x^*, y^*, E^*)$ is the unique interior point of the following equations:

$$rx^*\left(1 - \frac{x^*}{k}\right) - \frac{\alpha x^* y^*}{\alpha + x^*} - \frac{qE^* x^*}{m_1 E^* + m_2 x^*} = 0; sy^*\left(1 - \frac{\beta y^*}{a + x^*}\right) = 0$$

$$\eta E^* \left(\frac{qp x^*}{m_1 E^* + m_2 x^*} - c\right) = 0$$

$$x^* = k\left(1 - \frac{\alpha}{r\beta} - \frac{qL}{r(m_1L + m_2)}\right) \quad y^* = \frac{a + x^*}{\beta} \quad \text{and} \quad E^* = Lx^*$$

x^* and y^* are positive for $\frac{\alpha}{r\beta} + \frac{qL}{r(m_1L + m_2)} < 1$ and E^* is positive for $c < \frac{pq}{m_2}$.

6. Local Stability of Various Equilibrium Points

Here we discuss the local stability conditions for feasible equilibrium points of the system (2.1) based upon the standard linearization technique and then using the well-known Routh- Hurwitz criterion to determine the nature of eigenvalues of the Jacobian matrix evaluated at the equilibrium point[10]. The Jacobian matrix of the system (2.1) at any point (x, y, E) is given by:

$$J(x, y, E) = \begin{bmatrix} x \frac{\partial f}{\partial x} + f & x \frac{\partial f}{\partial y} & x \frac{\partial f}{\partial E} \\ y \frac{\partial g}{\partial x} & y \frac{\partial g}{\partial y} + g & y \frac{\partial g}{\partial E} \\ E \frac{\partial h}{\partial x} & E \frac{\partial h}{\partial y} & E \frac{\partial h}{\partial E} + h \end{bmatrix}$$

$$J(x, y, E) = \begin{bmatrix} x \left(-\frac{r}{k} + \frac{\alpha y}{(a+x)^2} + \frac{qEm_2}{(m_1E + m_2x)^2} \right) + f & \frac{-\alpha x}{a+x} & m_2q \left(\frac{x}{m_1E + m_2x} \right)^2 \\ \frac{\beta sy^2}{(a+x)^2} & \frac{\beta sy}{a+x} + g & 0 \\ \frac{m_1\eta pqE^2}{(m_1E + m_2x)^2} & 0 & -\frac{m_1\eta pqEx}{(m_1E + m_2x)^2} + h \end{bmatrix}$$

6.1. Stability of point (0, 0, 0);

The Jacobian matrix evaluated at the equilibrium point $P_0(0, 0, 0)$ is given by :

$$J_0(0,0,0) = \begin{bmatrix} -r & 0 & 0 \\ 0 & s & 0 \\ 0 & 0 & -\eta c \end{bmatrix}$$

The eigen values of J_0 evaluated at equilibrium point (0, 0, 0) are $\lambda_1 = -r < 0$, $\lambda_2 = s > 0$ and $\lambda_3 = -\eta c < 0$. Hence the origin (0, 0, 0) is a saddle point with unstable manifold in y direction and stable manifold in x as well as in E-direction.

6.2. Stability of point (k, 0, 0);

The Jacobian matrix evaluated at the equilibrium point $P_1(k, 0, 0)$ is given by:

$$J_1(k,0,0) = \begin{bmatrix} -r & \frac{-\alpha k}{a+k} & \frac{-q}{m_2} \\ 0 & s & 0 \\ 0 & 0 & \eta \left(\frac{pq}{m_2} - c \right) \end{bmatrix}$$

The eigen values of J_1 evaluated at equilibrium point (k, 0, 0) are $\lambda_1 = -r < 0$, $\lambda_2 = s > 0$ and $\lambda_3 = \eta \left(\frac{pq}{m_2} - c \right)$. The equilibrium point (k, 0, 0) is saddle point as $\lambda_2 = s > 0$ with unstable manifold in y- direction and stable manifold in x-direction and also in E-direction if $\frac{pq}{m_2} < c$.

6.3. Stability of point (0, a/β, 0);

The Jacobian matrix evaluated at the equilibrium point $P_2(0, a/\beta, 0)$ is given by:

$$J_2(0, \frac{a}{\beta}, 0) = \begin{bmatrix} r - \frac{\alpha}{\beta} & 0 & 0 \\ \frac{-s}{\beta} & -s & 0 \\ 0 & 0 & -\eta c \end{bmatrix}$$

The eigen values of J_2 evaluated at equilibrium point (k, 0, 0) are $\lambda_1 = r - \frac{\alpha}{\beta}$, $\lambda_2 = -s < 0$, $\lambda_3 = -\eta c < 0$ and . If $\lambda_1 < 0$, then the equilibrium point $P_2(0, a/\beta, 0)$ is locally asymptotically stable.

Otherwise, it is a saddle point with an unstable manifold in x-direction and stable manifold in y as well as in E direction.

6.4. Stability of point $P_3(\bar{x}, \bar{y}, 0)$;

The Jacobian matrix evaluated at the equilibrium point $P_3(\bar{x}, \bar{y}, 0)$ is given by:

$$J_3(\bar{x}, \bar{y}, 0) = \begin{bmatrix} \bar{x} \left(-\frac{r}{k} + \frac{\alpha \bar{y}}{(a + \bar{x})^2} \right) & -\frac{\alpha \bar{x}}{a + \bar{x}} & -\frac{q}{m_2} \\ \frac{\beta s \bar{y}^2}{(a + \bar{x})^2} & \frac{\beta s \bar{y}}{a + \bar{x}} & 0 \\ 0 & 0 & \eta \left(\frac{pq}{m_2} - c \right) \end{bmatrix}$$

The characteristic equation associated to the matrix $J_3(\bar{x}, \bar{y}, 0)$ yields the eigen values as follows:

$$\lambda_{1,2} = \left(\frac{\alpha}{\beta} - r - s \right) + \frac{s\beta \bar{x} \bar{y}}{a + \bar{x}} \pm \sqrt{\left(\left(\frac{\alpha}{\beta} - r - s \right) + \frac{\alpha \bar{x}}{a + \bar{x}} \right)^2 - 4 \frac{s\beta \bar{x} \bar{y}}{a + \bar{x}}} \quad \text{and} \quad \lambda_3 = \eta \left(\frac{pq}{m_2} - c \right)$$

- (i) If $\frac{s\beta \bar{x} \bar{y}}{a + \bar{x}} < r + s - \frac{\alpha}{\beta}$, then the equilibrium point P_3 is locally asymptotically stable. It is a saddle point with an unstable manifold in E-direction for $\frac{pq}{m_2} > c$. For $\frac{pq}{m_2} = c$, there is a chance of bifurcation.

6.5. Stability of equilibrium point $P_4(\hat{x}, 0, \hat{E})$

The Jacobian matrix evaluated at the equilibrium point $P_4(\hat{x}, 0, \hat{E})$ is given by:

$$J_4(\hat{x}, 0, \hat{E}) = \begin{bmatrix} \hat{x} \left(-\frac{r}{k} + \frac{q \hat{E} m_2}{(m_1 \hat{E} + m_2 \hat{x})^2} \right) & -\frac{\alpha \hat{x}}{a + \hat{x}} & -m_2 q \left(\frac{\hat{x}}{m_1 \hat{E} + m_2 \hat{x}} \right)^2 \\ 0 & s & 0 \\ \frac{m_1 \eta p q \hat{E}}{(m_1 \hat{E} + m_2 \hat{x})^2} & 0 & -\frac{m_1 \eta p q \hat{E} \hat{x}}{(m_1 \hat{E} + m_2 \hat{x})^2} \end{bmatrix}$$

Corresponding to equilibrium point P_4 , one of the eigen value is $\lambda_2 = s > 0$, and the other two can be

$$J^*_4(\hat{x}, 0, \hat{E}) = \begin{bmatrix} \hat{x} \left(-\frac{r}{k} + \frac{q \hat{E} m_2}{(m_1 \hat{E} + m_2 \hat{x})^2} \right) & -m_2 q \left(\frac{\hat{x}}{m_1 \hat{E} + m_2 \hat{x}} \right)^2 \\ \frac{m_1 \eta p q \hat{E}}{(m_1 \hat{E} + m_2 \hat{x})^2} & -\frac{m_1 \eta p q \hat{E} \hat{x}}{(m_1 \hat{E} + m_2 \hat{x})^2} \end{bmatrix}$$

obtained as the eigen values of the following 2×2 matrix:

If (i) $\text{Tr}(J^*_4) < 0$, the the point P_4 is a saddle point. (ii) If $\text{Tr}(J^*_4) = 0$, the we get pair of imaginary roots which shows that there is a case of hopf bifurcation.

6.6. Stability of Interior Equilibrium Point $P_5(x^*, y^*, E^*)$

Theorem: The unique positive interior equilibrium (x^*, y^*, E^*) is asymptotically locally stable provided condition (6.1) is satisfied.

The Jacobian matrix evaluated at the equilibrium point $P_5(x^*, y^*, E^*)$ is given by:

$$J(x^*, y^*, E^*) = \begin{bmatrix} x \left(-\frac{r}{k} + \frac{\alpha y^*}{(a+x^*)^2} + \frac{qE^* m_2}{(m_1 E^* + m_2 x^*)^2} \right) & \frac{-\alpha x^*}{a+x^*} & m_2 q \left(\frac{x^*}{m_1 E^* + m_2 x^*} \right)^2 \\ \frac{\beta s y^{*2}}{(a+x^*)^2} & \frac{\beta s y^*}{a+x^*} & 0 \\ \frac{m_1 \eta p q E^{*2}}{(m_1 E^* + m_2 x^*)^2} & 0 & -\frac{m_1 \eta p q E^* x^*}{(m_1 E^* + m_2 x^*)^2} \end{bmatrix}$$

The characteristics equations of the above Jacobian matrix about the equilibrium point $P_5(x^*, y^*, E^*)$ is given by:

$$\lambda^3 + A_1 \lambda^2 + A_2 \lambda + A_3 = 0,$$

where

$$\begin{aligned} A_1 &= -(a_{11} + a_{22} + a_{33}) \\ A_2 &= a_{22} a_{33} + (a_{11} a_{33} - a_{13} a_{31}) + (a_{11} a_{22} - a_{12} a_{21}) \\ A_3 &= a_{11} a_{22} a_{33} - a_{12} a_{21} a_{33} - a_{13} a_{31} a_{22} \end{aligned}$$

Using Routh- Hurwitz Criteria, the conditions for local stability of point P_5 are $A_1 > 0$, $A_2 > 0$ and $A_1 A_2 - A_3 > 0$

Note that $A_1 > 0$ if
$$\frac{-r}{k} - \frac{\alpha y^*}{(a+x^*)^2} - \frac{qE^* m_2}{(m_1 E^* + m_2 x^*)^2} < 0 \tag{6.1}$$

Also, $A_2 > 0$ and $A_1 A_2 - A_3 > 0$ for the condition (6.1)

Thus, the interior equilibrium point P_5 is locally asymptotically stable for the sufficient condition (6.1).

7. Global Stability

Theorem: The interior equilibrium point (x^*, y^*, E^*) is globally asymptotically stable.

Proof: Consider a Lyapunov function $V(x, y, E)$ such that:

$$V(x, y, E) = d_0 \left[(x - x^*) - x^* \log \frac{x}{x^*} \right] + d_1 \left[(y - y^*) - y^* \log \frac{y}{y^*} \right] + d_2 \left[(E - E^*) - E^* \log \frac{E}{E^*} \right]$$

As $V(x, y, E)$ is zero at the equilibrium point (x^*, y^*, E^*) i.e., $V(x^*, y^*, E^*) = 0$

and positive for all other values of (x, y, E) .

$$\begin{aligned} \frac{dV}{dt} &= d_0 (x - x^*) \frac{\dot{x}}{x} + d_1 (y - y^*) \frac{\dot{y}}{y} + d_2 (E - E^*) \frac{\dot{E}}{E} \\ &= d_0 (x - x^*) \left(r \left(1 - \frac{x}{k} \right) - \frac{\alpha y}{a+x} - \frac{qE}{m_1 E + m_2 x} \right) + d_1 (y - y^*) \left(s - \frac{\beta s y}{a+x} \right) + d_2 (E - E^*) \left(\frac{q p x}{m_1 E + m_2 x} - c \right) \end{aligned}$$

$$\frac{dV}{dt} = d_0 (x - x^*)^2 \left[\frac{r}{k} - \frac{\alpha y^2}{(a+x)(a+x^*)} - \frac{qE^*}{(m_1 E + m_2 x)(m_1 E^* + m_2 x^*)} \right] + \frac{(x - x^*)(y - y^*)}{(a+x)} \left[-d_0 \alpha x^* + \frac{d_1 s \beta E^*}{a+x^*} \right]$$

$$+ \frac{-d_0 q x^* + d_2 q p E^*}{(m_1 E + m_2 x)(m_1 E^* + m_2 x^*)} (x - x^*)(E - E^*) - \frac{d_1 s \beta (y - y^*)^2}{a+x} - \frac{d_2 q p x^* (E - E^*)^2}{(m_1 E + m_2 x)(m_1 E^* + m_2 x^*)}$$

$$\frac{dV}{dt} = -d_0 (x - x^*)^2 \left[\frac{r}{k} - \frac{\alpha y^2}{(a+x)(a+x^*)} - \frac{qE^*}{(m_1 E + m_2 x)(m_1 E^* + m_2 x^*)} \right] + \frac{d_1 s \beta (y - y^*)^2}{a+x} - \frac{d_2 q p x^* (E - E^*)^2}{(m_1 E + m_2 x)(m_1 E^* + m_2 x^*)}$$

Let $d_0 = 1$ $d_1 = \frac{a+x}{s \beta E^*}$ and $d_2 = \frac{q x^*}{q p E^*}$

$$\frac{dV}{dt} < 0 \quad \text{if} \quad \frac{r}{k} - \frac{\alpha y^*}{(a+x)(a+x^*)} - \frac{qE^*}{(m_1 E + m_2 x)(m_1 E^* + m_2 x^*)} > 0 \tag{7.1}$$

Thus, interior equilibrium point (x^*, y^*, E^*) is globally stable for the above condition (7.1).

8. Numerical Simulations

In this section, numerical simulations are carried out for different choice of parameters to investigate the dynamic behavior of the system. We start the numerical investigation keeping all the parameters fixed except 'c'.

Let $r = 0.5$, $k = 100$, $\alpha = 0.005$, $m_1 = 0.5$, $m_2 = 0.5$, $q = 0.15$, $a = 3$, $s = 1$, $\beta = 0.15$, $\eta = 1$, $p = 5$;




Figure 1: Time series analysis of prey population, predator population and Effort 'E' for different value of cost 'c' with initial level (40, 350, 10).




Figure 2: Phase portrait of prey population, predator population and Effort 'E' for different initial level corresponding (A).c=1.2 and (B). c=1.8

In the figure-1, diagrams (A), (B) and (C) give long term behavior of trajectories of prey- predator populations and effort 'E' w.r.t time 't'. which shows that for the initial condition (40, 350, 10) in the interior, for the different value of cost 'c' and keeping other parameters fixed, all the trajectories converges to its interior equilibrium point which means that interior equilibrium is asymptotically locally stable provided $0 < c < 1.5$. Also, we see that as the value of cost 'c' will start increase then eventually harvesting effort will start decrease and in resulted prey population increases as well predator population also. But, for $c > 1.5$ effort 'E' will tends to zero.

Figure-2(A) represents phase plane trajectories of species x, y and effort E with the different initial levels which represents that the interior point $(x^*, y^* E^*) = (81.3332, 562.2056, 20.333)$ is asymptotically global stable for $c = 1.2$ as $0 < c < 1.5$ for the existence of interior equilibrium point. The figure-2(B) represents phase plane trajectories of different biomasses with the different initial levels at the interior which converge to the point $(93.3340, 642.2877, 0.0000)$ on the boundary plane i.e., x-y plane corresponding to $c = 1.8$ for the condition $c > 1.5$, keeping other parameters fixed.. Therefore, for the condition $c > 1.5$, it shows that for the every initial points on the x - y - E space converge to the point on x - y plane asymptotically which means that there is on harvesting of the prey population for this condition as it is not profitable to fishermen to do harvesting.

5. Conclusion

This paper is concerned with the study of nonlinear harvesting and the conservation of ecological resources for a two-dimensional prey predator type dynamical system. We have considered a prey predator model with Holling type-II functional response and nonlinear harvesting of prey. The harvesting effort is taken as a dynamic variable. The conditions for existence and local asymptotic stability of various equilibrium points have been examined. It is established that the coexistence of prey and predator population depends upon the proper harvesting strategies

and hence we can avoid the risk of extinction or over exploitation of the species. By analytical and numerical results, we have examined that for fixed value of price 'p' per unit mass and other parameters, as the value of cost 'c' is increasing, then harvesting level will start to decrease. after a time, we will find a level of cost 'c' where harvesting effort will tends to zero. So for the co-existence of prey- predator populations along with effort dynamics, there is a restriction on value of cost 'c' so that there can co-exist all the species with optimal level of harvesting effort.

References

- Aziz-Alaoui M.A., Daher Okiye M., Boundedness and global stability for a predator prey model with modified Leslie- Gower and Holling type-II schemes, *Appl. Math. Lett.* 16(7)(2003)1069 - 1075.
- Aziz-Alaoui M. A., Study of a Leslie- Gower- type tri-trophic population, *Chaos, Solitons and Fractals* 14(2002)1275 - 1293.
- Clark C.W., *Mathematical Bio-economics: the Optimal Management of Renewable Resources*, Wiley, New York, 1976.
- Gakkhar, S. Singh, B.P., The dynamics of a food web consisting of two preys and a harvesting predator, *chaos, Solitons and fractals* 34(2007)1346 -1356.
- Gupta RP., Chandra P., Bifurcation analysis of modified Leslie- Gower predator- prey model with Michaelis-Menten type prey harvesting. *J Math Anal Appl* 398(2013)278 - 95.
- Krishna S. V., Srinivasu P. D. N. and Kaymakcalan B., Conservation of an ecosystem through optimal taxation, *Bulletin of Mathematical Biology*, 60(1998)569 - 584.
- Lenzini P., Rebaza J., Non-constant predator harvesting on ratio- dependent predator- prey models, *Appl. Math. Sci.* 4(16)(2010)791 - 803.
- Li Y., Xiao D., Bifurcations of a predator prey system of Holling and Leslie types, *Chaos Solitons Fractals* 34(2007)606 - 620.
- Murray J. D., *Mathematical Biology*, vol. I, Springer- Verlag, Berlin, 2003.
- Perko L., *Differential Equations and Dynamical Systems*, Springer, New York, 1996.
- Pradhan T., Chaudhuri K. S., A dynamic reaction model of two- species fishery with taxation as a control instrument: A capital theoretic analysis, *Ecological Modeling* 121(1999)1 - 16.
- Srinivasu P.D.N., Bio-economics of a renewable resource in presence of a predator, *Nonlinear Anal. RWA* 2(2001)497 - 06.
- Xiao D., Jennings L., Bifurcations of a ratio- dependent predator-prey system with constant rate harvesting, *SIAM J. Appl. Math.* 65(2005)737 - 753.
- Zhang, N., Chen, F., Su, Q., Wu, T., Dynamic behaviors of a harvesting Leslie Gower predator prey model. *Discrete Dyn. Nat. Soc.*2011(2011).
- Zhu, C.R., Lan, K.Q., Phase portraits, Hopf bifurcations and limit cycles of Leslie Gower predator prey systems with harvesting rates. *Discrete Contin. Dyn. Syst. Ser. B* 14(2010)289 - 306.

Reenu Rani: She is pursuing her PhD at IIT Roorkee since July 2012. Her research area is Mathematical Modelling of Renewable Resources. She has completed her M.Sc. (Hns) from Panjab University, Chandigarh during the session 2008-2010. She has done her graduation from Mata Gujri College, Fatehgarh Sahib affiliated to Punjabi University, Patiala during 2004-2007.

Sunita Gakkhar: She is working as Professor at IIT Roorkee. Her main areas of interest are Bio- mathematics, computer Applications, Fluid Dynamics, Mathematical Modelling of Environmental and Ecological Systems. She has done her PhD at IIT Kanpur in 1982. She has completed her M.Sc. in Applied Mathematics from University of Roorkee in 1977 and B.Sc. in Maths, Physics and Chemistry from Meerut University in 1975. She had Certificate of Merit Institution of Engineers award: for the paper in *Journal of Institution of Engineers (India)* (1996-1997). She is awarded Chancellor's Gold medal: for securing highest marks in all postgraduate courses in the university (1977) and also A N Khosla Gold Medal: For securing Highest marks in M.Sc. (1977). She has many publications in different area

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

