

# Quantum Mechanical Behaviour, Quantum Tunneling, Higgs Boson , Distorted Space And Time, Schrödinger's Wave Function, Neuron DNA, Particles (Hypothetical signature Less Particles) And Consciousness

## A “Syncopated Syncretism And Atrophied Asseveration” Model

<sup>\*1</sup>Dr K N Prasanna Kumar, <sup>2</sup>Prof B S Kiranagi And <sup>3</sup>Prof C S Bagewadi

<sup>\*1</sup>Dr K N Prasanna Kumar, Post doctoral researcher, Dr KNP Kumar has three PhD's, one each in Mathematics, Economics and Political science and a D.Litt. in Political Science, Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India [Correspondence Mail id : drknpkumar@gmail.com](mailto:drknpkumar@gmail.com)

<sup>2</sup>Prof B S Kiranagi, UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

<sup>3</sup>Prof C S Bagewadi, Chairman , Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu university, Shankarghatta, Shimoga district, Karnataka, India

### Abstract:

We give a holistic model for the systems mentioned in the foregoing. Most important implication is that Higgs Boson is the one, which warps space and time. Concept of Neuron DNA and signature less particles are introduced.

### Key Words:

Quantum Information, Space warp, Quantum Tunneling, Environmental decoherence, Schrödinger's wave function, Gravitational lensing, Black holes, Higgs Boson, Consciousness

### Introduction:

We take in to consideration the following to build the 36 story model which consummates and consolidates the parameters and processes involved:

1. Quantum Information
2. Quantum Mechanical behaviour
3. Quantum Tunneling
4. Non adiabatic multi photon process in the strong vibronic coupling limit
5. Environmental Decoherence(Green House Effects for example)
6. Schrodinger's wave function
7. Gravitational lensing
8. Black holes
9. Faster than Light Particles (Neuron DNA- Mind, a signature less particles. How do you classify that? Total energy =Existing matter-Energy attributable to signature less particles. Einstein did not take in consideration psychic energy which is taken to be holistically conservational ,but individually and collectively non conservative)
10. Consciousness( Total awareness- use ASDCII and Information field capacity to find the total storage- Please refer Gesellschaft-Gememshaft paper on the subject matter)
11. Higgs Boson

12. Distorted Space and Time

**Notation :**

**Quantum Mechanical Behaviour And Quantum Information  
Module Numbered One**

$G_{13}$  : Category One Of Quantum Mechanical Behaviour

$G_{14}$  : Category Two Of Quantum Mechanical Behaviour

$G_{15}$  : Category Three Of Quantum Mechanical Behaviour

$T_{13}$  : Category One Of Quantum Information

$T_{14}$  : Category Two Of Quantum Information

$T_{15}$  :Category Three Of Quantum Information

**Non Adiabatic Multi Phonon Process In The Strong Vibronic Coupling And Quantum Tunneling**

**Module Numbered Two:**

$G_{16}$  : Category One Of Non Adiabatic Multi Phonon Process

$G_{17}$  : Category Two Of Non Adiabatic Multi phonon Process

$G_{18}$  : Category Three Of Non Adiabatic Multi Phonon Process

$T_{16}$  :Category One Of Quantum Tunneling(There Are Lot Of Tunnels)

$T_{17}$  : Category Two Of Quantum Tunneling

$T_{18}$  : Category Three Of Quantum Tunneling

**Environmental Decoherence (For Example Green House Effects) And Collapse of Schrodinger's Wave  
Function:**

**Module Numbered Three:**

$G_{20}$  : Category One Of Collapse Of Schrodinger's Wave Function(There Are Lot Of Potentialities)

$G_{21}$  :Category Two Of Collapse Of Schrodinger's Wave Function

$G_{22}$  : Category Three Of Collapse Of Schrodinger's Wave Function

$T_{20}$  : Category One Of Environmental Decoherence

$T_{21}$  :Category Two Of Environmental Decoherence

$T_{22}$  : Category Three Of Environmental Decoherence

**Gravitational Lensing And Black holes**

**Module Numbered Four:**

$G_{24}$  : Category One Of Black holes

$G_{25}$  : Category Two Of Black holes

$G_{26}$  : Category Three Ofblack Holes

$T_{24}$  :Category One Of gravitational Lensing

$T_{25}$  :Category Two Of Gravitational Lensing

$T_{26}$  : Category Three Of Gravitational Lensing

**Faster Than Light Particles(Hypothetical Particles Of Neuron DNA-Mind) And Consciousness(Total Awareness With Visual Images: Calculated Based On Ascii And Information Field Capacity)**

**Module Numbered Five:**

$G_{28}$  : Category One Of Faster Than Light Particles(Signatureless Neuron Dna)

$G_{29}$  : Category Two Of Faster Than Light(Signatureless neuron Dna)

$G_{30}$  :Category Three Of Faster Thank Light Neuron Dna Particles Without Signature

$T_{28}$  :Category One Of Consciousness(Just Total Knowledge That Is Stored Like In Computer-See Gratification Deprivation Model For Details)

$T_{29}$  :Category Two Of Consciousness

$T_{30}$  :Category Three Of Consciousness

**Distorted Space And Time (St Warp) And Higgs Boson**

**Module Numbered Six:**

$G_{32}$  : Category One Of Higgs Boson

$G_{33}$  : Category Two Of Higgs Boson

$G_{34}$  : Category Three Of Higgs Boson

$T_{32}$  : Category One Of Distorted Space And Time

$T_{33}$  : Category Two Of Distorted Space And Time

$T_{34}$  : Category Three Of Distorted Space And Time

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$   
 $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)} : (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$   
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$   
 $(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$   
*are Accentuation coefficients*

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$   
 $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$   
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$

$$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$$

are Dissipation coefficients

### Quantum Mechanical Behaviour And Quantum Information

#### Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15}$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15}$$

$$+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor}$$

$$-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}$$

### Non Adiabatic Multi Phonon Process In The Strong Vibronic Coupling And Quantum Tunneling

#### Module Numbered Two:

The differential system of this model is now ( Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16}$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17}$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18}$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17}$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18}$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor}$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor}$$

### Environmental Decoherence (For Example Green House Effects) And Collapse of Schrodinger's Wave Function:

#### Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22}$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21}$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22}$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \textit{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \textit{First detritions factor}$$

### Gravitational Lensing And Black holes

#### Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24}$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26}$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24}$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25}$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26}$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \textbf{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \textbf{First detritions factor}$$

### Faster Than Light Particles (Hypothetical Particles Of Neuron Dna-Mind) And Consciousness(Total Awareness With Visual Images: Calculated Based On Ascii And Information Field Capacity)

#### Module Numbered Five

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29}$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30}$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28}$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29}$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30}$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \textit{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \textit{First detritions factor}$$

### Distorted Space And Time(St Warp) And Higgs Boson:

#### Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34}$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32}$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33}$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34}$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \textit{First augmentation factor}$$

$$-(b''_{32})^{(6)}((G_{35}), t) = \textit{First detritions factor}$$

### Holistic Concatenated Equations Henceforth Referred To As “Global Equations”

1. Quantum Information
2. Quantum Mechanical behavior
3. Quantum Tunneling
4. Non adiabatic multi photon process in the strong vibronic coupling limit
5. Environmental Decoherence(Green House Effects for example)
6. Schrodinger's wave function
7. Gravitational Lensing
8. Black holes
9. Faster than Light Particles (Neuron DNA- Mind, a signature less particles How do you classify that? Total energy =Existing matter-Energy attributable to signature less particles. Einstein did not take in consideration psychic energy which is taken to be holistically conservational ,but individually and collectively non conservative)
10. Consciousness( Total awareness- use ASCII and Information field capacity to find the total storage- Please refer Gesellschaft- Gemeinschaft paper on the subject matter)
11. Higgs Boson
12. Distorted Space and Time

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[ \begin{array}{ccc} (a'_{13})^{(1)} \boxed{+(a''_{13})^{(1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[ \begin{array}{ccc} (a'_{14})^{(1)} \boxed{+(a''_{14})^{(1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[ \begin{array}{ccc} (a'_{15})^{(1)} \boxed{+(a''_{15})^{(1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)} \end{array} \right] G_{15}$$

Where  $\boxed{(a''_{13})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{14})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{15})^{(1)}(T_{14}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{ccc} (b'_{13})^{(1)} \boxed{-(b''_{13})^{(1)}(G, t)} & \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{ccc} (b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} & \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{ccc} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} & \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{15}$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[ \begin{array}{ccc} (a'_{16})^{(2)} \boxed{+(a''_{16})^{(2)}(T_{17}, t)} & \boxed{+(a''_{13})^{(1,1)}(T_{14}, t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \end{array} \right] G_{16}$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[ \begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{17}$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[ \begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{18}$$

Where  $+(a''_{16})^{(2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2)}(T_{17}, t)$  are first augmentation coefficients for category 1, 2 and 3  
 $+(a''_{13})^{(1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1)}(T_{14}, t)$  are second augmentation coefficient for category 1, 2 and 3  
 $+(a''_{20})^{(3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3  
 $+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3  
 $+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3  
 $+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[ \begin{array}{ccc} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{16}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[ \begin{array}{ccc} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{17}$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[ \begin{array}{ccc} (b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}, t) & - (b''_{15})^{(1,1)}(G, t) & - (b''_{22})^{(3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{18}$$

where  $-(b''_{16})^{(2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2)}(G_{19}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{13})^{(1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1)}(G, t)$  are second detrition coefficients for category 1,2 and 3  
 $-(b''_{20})^{(3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1,2 and 3  
 $-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1,2 and 3  
 $-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1,2 and 3  
 $-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[ \begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a''_{16})^{(2,2,2)}(T_{17}, t) & + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[ \begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a''_{17})^{(2,2,2)}(T_{17}, t) & + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{21}$$



$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[ \begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a''_{18})^{(2,2,2)}(T_{17}, t) & + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{22}$$

$+(a''_{20})^{(3)}(T_{21}, t), +(a''_{21})^{(3)}(T_{21}, t), +(a''_{22})^{(3)}(T_{21}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2)}(T_{17}, t)$  are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1)}(T_{14}, t)$  are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[ \begin{array}{ccc} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) & - (b''_{16})^{(2,2,2)}(G_{19}, t) & - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{20}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[ \begin{array}{ccc} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) & - (b''_{17})^{(2,2,2)}(G_{19}, t) & - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{21}$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{ccc} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) & - (b''_{18})^{(2,2,2)}(G_{19}, t) & - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{22}$$

$-(b''_{20})^{(3)}(G_{23}, t), -(b''_{21})^{(3)}(G_{23}, t), -(b''_{22})^{(3)}(G_{23}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t), -(b''_{14})^{(1,1,1)}(G, t), -(b''_{15})^{(1,1,1)}(G, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6,6,6)}(G_{35}, t), -(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \end{array} \right] G_{24}$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \end{array} \right] G_{25}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \end{array} \right] G_{26}$$

Where  $(a''_{24})^{(4)}(T_{25}, t), (a''_{25})^{(4)}(T_{25}, t), (a''_{26})^{(4)}(T_{25}, t)$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5)}(T_{29}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, and 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[ \begin{array}{ccc} \boxed{(b'_{24})^{(4)}} & \boxed{-(b''_{24})^{(4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)} & \end{array} \right] T_{24}$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[ \begin{array}{ccc} \boxed{(b'_{25})^{(4)}} & \boxed{-(b''_{25})^{(4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)} & \end{array} \right] T_{25}$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[ \begin{array}{ccc} \boxed{(b'_{26})^{(4)}} & \boxed{-(b''_{26})^{(4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)} & \end{array} \right] T_{26}$$

Where  $\boxed{-(b''_{24})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4)}(G_{27}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[ \begin{array}{ccc} \boxed{(a'_{28})^{(5)}} & \boxed{+(a''_{28})^{(5)}(T_{29}, t)} & \boxed{+(a''_{24})^{(4,4)}(T_{25}, t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} & \end{array} \right] G_{28}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[ \begin{array}{ccc} \boxed{(a'_{29})^{(5)}} & \boxed{+(a''_{29})^{(5)}(T_{29}, t)} & \boxed{+(a''_{25})^{(4,4)}(T_{25}, t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} & \end{array} \right] G_{29}$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[ \begin{array}{ccc} \boxed{(a'_{30})^{(5)}} & \boxed{+(a''_{30})^{(5)}(T_{29}, t)} & \boxed{+(a''_{26})^{(4,4)}(T_{25}, t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} & \end{array} \right] G_{30}$$

Where  $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$  are first augmentation coefficients for category 1, 2 and 3

And  $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[ \begin{array}{ccc} \boxed{(b'_{28})^{(5)}} & \boxed{-(b''_{28})^{(5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} & \end{array} \right] T_{28}$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[ \begin{array}{ccc} \boxed{(b'_{29})^{(5)}} & \boxed{-(b''_{29})^{(5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} & \end{array} \right] T_{29}$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[ \begin{array}{ccc} \boxed{(b'_{30})^{(5)}} & \boxed{-(b''_{30})^{(5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} & \end{array} \right] T_{30}$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[ \begin{array}{ccc} \boxed{(a'_{32})^{(6)}} & \boxed{+(a''_{32})^{(6)}(T_{33}, t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)} & \boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)} & \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{ccc} \boxed{(a'_{33})^{(6)}} & \boxed{+(a''_{33})^{(6)}(T_{33}, t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)} & \boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)} & \end{array} \right] G_{33}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{ccc} \boxed{(a'_{34})^{(6)}} & \boxed{+(a''_{34})^{(6)}(T_{33}, t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)} & \boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)} & \end{array} \right] G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6)}(T_{33}, t)}$  are first augmentation coefficients for category 1,2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)}$  are second augmentation coefficients for category 1,2 and 3

$\boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)}$  are third augmentation coefficients for category 1,2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)}$  - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)}$  - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)}$  sixth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{32})^{(6)} \boxed{-(b''_{32})^{(6)}(G_{35}, t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{32}$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{ccc} (b'_{33})^{(6)} \boxed{-(b''_{33})^{(6)}(G_{35}, t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{33}$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{34})^{(6)} \boxed{-(b''_{34})^{(6)}(G_{35}, t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6)}(G_{35}, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)}$  are third detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2, and 3  
 $\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2, and 3  
 $\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2, and 3

Where we suppose

- (A)  $(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0$ ,  
 $i, j = 13, 14, 15$
- (B) The functions  $(a''_i)^{(1)}, (b''_i)^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

(C)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)}$   
 $\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$  are positive constants and  $\boxed{i = 13, 14, 15}$

They satisfy Lipschitz condition:

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t) \cdot (T_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the first augmentation coefficient WOULD be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  :

(D)  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  :

(E) There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together with  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ ,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

(F)  $(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, i, j = 16, 17, 18$

(G) The functions  $(a_i'')^{(2)}, (b_i'')^{(2)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ :

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)}$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}$$

(H)  $\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)}$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)}$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$  :

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}, t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\hat{M}_{16})^{(2)}t}$$

$$|(b_i'')^{(2)}(G_{19}, t) - (b_i'')^{(2)}(G_{19}, t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(2)}(T_{17}, t)$  and  $(a_i'')^{(2)}(T_{17}, t) \cdot (T_{17}, t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a_i'')^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a_i'')^{(2)}(T_{17}, t)$ , the SECOND augmentation coefficient would be absolutely continuous.

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$  :

(I)  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$  :

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18$ ,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [ (a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)} ] < 1$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [ (b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)} ] < 1$$

Where we suppose

(J)  $(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20,21,22$

The functions  $(a''_i)^{(3)}, (b''_i)^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$  :

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20,21,22$

They satisfy Lipschitz condition:

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T'_{21} - T_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G'_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(3)}(T'_{21}, t)$  and  $(a''_i)^{(3)}(T_{21}, t) \cdot (T'_{21}, t)$  And  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a''_i)^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a''_i)^{(3)}(T_{21}, t)$ , the THIRD augmentation coefficient, would be absolutely continuous.

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$  :

(K)  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22,$  satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26$$

(M) The functions  $(a''_i)^{(4)}, (b''_i)^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$(N) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$  :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

They satisfy Lipschitz condition:

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(4)}(T'_{25}, t)$  and  $(a''_i)^{(4)}(T_{25}, t)$ .  $(T'_{25}, t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a''_i)^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 4$  then the function  $(a''_i)^{(4)}(T_{25}, t)$ , the **FOURTH augmentation coefficient WOULD** be absolutely continuous.

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$  :

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$  :

(Q) There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants

$(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26,$   
 satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30$$

(S) The functions  $(a''_i)^{(5)}, (b''_i)^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$ :

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$(T) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$ :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$

They satisfy Lipschitz condition:

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31})' - (G_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(5)}(T'_{29}, t)$  and  $(a''_i)^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to be noted that  $(a''_i)^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 5$  then the function  $(a''_i)^{(5)}(T_{29}, t)$ , the FIFTH **augmentation coefficient** attributable would be absolutely continuous.

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ :

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$ :

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants

$(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30,$  satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$



$$\frac{1}{(\hat{M}_{28})^{(5)}} [ (b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)} ] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$$

(W) The functions  $(a''_i)^{(6)}, (b''_i)^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$(X) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$  :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}((G'_{35}), t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} |(G'_{35}) - (G_{35})| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(6)}(T'_{33}, t)$  and  $(a''_i)^{(6)}(T_{33}, t)$ .  $(T'_{33}, t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a''_i)^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 6$  then the function  $(a''_i)^{(6)}(T_{33}, t)$ , the SIXTH **augmentation coefficient** would be absolutely continuous.

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$  :

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} + \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$  :

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [ (a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} ] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [ (b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)} ] < 1$$

**Theorem 1:** if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

Definition of  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Definition of**  $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$$

By

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{13})^{(1)} + a''_{13}(s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[ (a_{14})^{(1)} G_{13}(s_{(13)}) - \left( (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[ (a_{15})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[ (b_{13})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[ (b_{14})^{(1)} T_{13}(s_{(13)}) - \left( (b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[ (b_{15})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + a''_{17}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + a''_{18}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + a''_{21}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

By

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

(a) The operator  $\mathcal{A}^{(1)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself. Indeed it is obvious that

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} \left( G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left( 1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left( e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ \left( (\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

(b) The operator  $\mathcal{A}^{(2)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} \left( G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} =$$

$$\left( 1 + (a_{16})^{(2)} t \right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left( e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[ \left( (\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{\left( -\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for  $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

(a) The operator  $\mathcal{A}^{(3)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} \left( G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$\left( 1 + (a_{20})^{(3)} t \right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left( e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[ \left( (\hat{P}_{20})^{(3)} + G_{21}^0 \right) e^{\left( -\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0} \right)} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

(b) The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} \left( G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$\left( 1 + (a_{24})^{(4)} t \right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left( e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[ \left( (\hat{P}_{24})^{(4)} + G_{25}^0 \right) e^{\left( -\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0} \right)} + (\hat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

(c) The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} \left( G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$\left( 1 + (a_{28})^{(5)} t \right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left( e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[ ((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

(d) The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} \left( G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left( e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[ ((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$  and to choose

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{13})^{(1)}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ ((\hat{Q}_{13})^{(1)} + T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)}$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying GLOBAL EQUATIONS into itself

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

**Definition of  $\tilde{G}, \tilde{T}$  :**

$$(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ \int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} + \\ (a''_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} + \\ G_{13}^{(2)} |(a'_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) - (a'_{13})^{(1)}(T_{14}^{(2)}, s_{(13)})| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)}t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)}(\widehat{K}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}))$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a'_{13})^{(1)}$  and  $(b'_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$  and  $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(1)}$  and  $(b''_i)^{(1)}, i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$  and  $((\widehat{M}_{13})^{(1)})_3$  :

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$$G_{13} < ((\widehat{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}, G_{15}$  and  $G_{13}, G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below.

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$  then  $T_{14} \rightarrow \infty$ .

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :



Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$  and to choose

$(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ (\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ ((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric

$$d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote

$$\underline{\text{Definition of}} \quad \widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$$

It results

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a'_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a'_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} \left( (a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)} \right) d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \right)$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(2)}$  and  $(b''_i)^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \quad \text{for } t > 0$$

**Definition of**  $((\widehat{M}_{16})^{(2)})_1$ ,  $((\widehat{M}_{16})^{(2)})_2$  and  $((\widehat{M}_{16})^{(2)})_3$  :

**Remark 3:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$ . indeed if

$$G_{16} < ((\widehat{M}_{16})^{(2)})_1 \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$ ,  $G_{18}$  and  $G_{16}$ ,  $G_{17}$  respectively.

**Remark 4:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below.

**Remark 5:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$  then  $T_{17} \rightarrow \infty$ .

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b''_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_2}$  By taking now  $\varepsilon_2$  sufficiently small one sees that  $T_{17}$  is unbounded. The same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b''_{18})^{(2)}((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$  and to choose

$(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ (\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{20})^{(3)}$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ ((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)}$$

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric

$$d\left((G_{23})^{(1)}, (T_{23})^{(1)}, (G_{23})^{(2)}, (T_{23})^{(2)}\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote

$$\underline{\text{Definition of } \widetilde{G}_{23}, \widetilde{T}_{23} : ((\widetilde{G}_{23}), (\widetilde{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))}$$

It results

$$\begin{aligned} |\tilde{G}_{20}^{(1)} - \tilde{G}_{20}^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)} \end{aligned}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ &\frac{1}{(\bar{M}_{20})^{(3)}} \left( (a_{20})^{(3)} + (a'_{20})^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)} \right) d\left( ((G_{23})^{(1)}, (T_{23})^{(1)}; (G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  and  $(\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}, i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t ((a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})) ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \quad \text{for } t > 0$$

**Definition of**  $((\widehat{M}_{20})^{(3)})_1$ ,  $((\widehat{M}_{20})^{(3)})_2$  and  $((\widehat{M}_{20})^{(3)})_3$  :

**Remark 3:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$ . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$  it follows  $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$  and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21}')^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22}')^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$ ,  $G_{22}$  and  $G_{20}$ ,  $G_{21}$  respectively.

**Remark 4:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below.

**Remark 5:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$  then  $T_{21} \rightarrow \infty$ .

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21}'')^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21}')^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to

$$T_{21} \geq \left( \frac{(a_{21}')^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \quad \text{If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left( \frac{(a_{21}')^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \quad \text{By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$  and to choose

$(\widehat{P}_{24})^{(4)}$  and  $(\widehat{Q}_{24})^{(4)}$  large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ (\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)}$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[ ((\hat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)} \right] \leq (\hat{Q}_{24})^{(4)}$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying IN to itself

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric

$$d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{27}), (\widetilde{T}_{27}) : (\widetilde{G}_{27}), (\widetilde{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} &\left( (a_{24})^{(4)} + (a'_{24})^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)} \right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right); \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$  and  $(\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}, i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \} ds_{(24)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \quad \text{for } t > 0$$

**Definition of**  $((\widehat{M}_{24})^{(4)})_1$ ,  $((\widehat{M}_{24})^{(4)})_2$  and  $((\widehat{M}_{24})^{(4)})_3$  :

**Remark 3:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$  . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)})_1 \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$  ,  $G_{26}$  and  $G_{24}$  ,  $G_{25}$  respectively.

**Remark 4:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below.

**Remark 5:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$  then  $T_{25} \rightarrow \infty$ .

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions ANALOGOUS inequalities hold also for  $G_{29}$  ,  $G_{30}$ ,  $T_{28}$ ,  $T_{29}$ ,  $T_{30}$

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$  and to choose

$(\widehat{P}_{28})^{(5)}$  and  $(\widehat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ (\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ ((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)}$$

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric

$$d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\bar{G}_{31}), (\bar{T}_{31}) : (\bar{G}_{31}), (\bar{T}_{31}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_{28}^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{-(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\bar{M}_{28})^{(5)}} &\left( (a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)} \right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right); \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (35,35,36) the result follows

**Remark 1:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$  and  $(\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}$ ,  $i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \quad \text{for } t > 0$$

**Definition of**  $((\bar{M}_{28})^{(5)})_1, ((\bar{M}_{28})^{(5)})_2$  and  $((\bar{M}_{28})^{(5)})_3 :$

**Remark 3:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$  it follows  $\frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)}G_{29}$  and by integrating

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$ ,  $G_{30}$  and  $G_{28}$ ,  $G_{29}$  respectively.

**Remark 4:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below.

**Remark 5:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$  then  $T_{29} \rightarrow \infty$ .

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29})^{(5)} - (b''_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$  which leads to

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$  and to choose

$(\widehat{P}_{32})^{(6)}$  and  $(\widehat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ (\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ ((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)}$$

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric

$$d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$



$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)}t} \}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{35}, \widehat{T}_{35}) : (\widehat{G}_{35}, \widehat{T}_{35}) = \mathcal{A}^{(6)}((G_{35}, T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} | (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)}t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}); (G_{35})^{(2)}, (T_{35})^{(2)} \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{32})^{(6)}$  and  $(b''_{32})^{(6)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  and  $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(6)}$  and  $(b''_i)^{(6)}$ ,  $i = 32, 33, 34$  depend only on  $T_{33}$  and respectively on  $(G_{35})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 69 to 32 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(6)} - (a''_i)^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$  and  $((\widehat{M}_{32})^{(6)})_3 :$

**Remark 3:** if  $G_{32}$  is bounded, the same property have also  $G_{33}$  and  $G_{34}$ . indeed if

$$G_{32} < ((\widehat{M}_{32})^{(6)}) \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If  $G_{33}$  or  $G_{34}$  is bounded, the same property follows for  $G_{32}$ ,  $G_{34}$  and  $G_{32}$ ,  $G_{33}$  respectively.

**Remark 4:** If  $G_{32}$  is bounded, from below, the same property holds for  $G_{33}$  and  $G_{34}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{33}$  is bounded from below.

**Remark 5:** If  $T_{32}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$  then  $T_{33} \rightarrow \infty$ .

**Definition of**  $(m)^{(6)}$  and  $\varepsilon_6$  :

Indeed let  $t_6$  be so that for  $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then  $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$  which leads to

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for  $T_{34}$  if  $\lim_{t \rightarrow \infty} (b_{34}'')^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

### **Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  :

(a)  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

**Definition of**  $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$  :

(b) By  $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$  and respectively  $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$  the roots of the equations  $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$  and  $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

**Definition of**  $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$  :

By  $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$  and respectively  $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$  the roots of the equations  $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$  and  $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

**Definition of**  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$  :-

(c) If we define  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$  by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined respectively

Then the solution satisfies the inequalities

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where  $(p_i)^{(1)}$  is defined

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left( \frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[ e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \leq \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[ e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[ e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[ e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

**Definition of**  $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$ :-

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

### **Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  :

(d)  $\sigma_1^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) - (b''_{17})^{(2)}(G_{19}, t) \leq -(\tau_1)^{(2)}$$

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  :

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots

(e) of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  :

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :-

(f) If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

and 
$$(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

and analogously

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and 
$$(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)}$$

Then the solution satisfies the inequalities

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$  is defined

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[ e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} \left[ e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t} \right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})} \left[ e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[ e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :-

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

### **Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

(a)  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  :

(b) By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :-

(c) If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$$

$$\left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[ e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{21}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[ e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :-

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)} (m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)} (\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

### **Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

(d)  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  :

(e) By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  :

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the

roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and  $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

(f) If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)}$  where  $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$  are defined by 59 and 64 respectively

Then the solution satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-a'_{26}^{(4)}t} \right] + G_{26}^0 e^{-a'_{26}^{(4)}t}$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[ e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

(g)  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  :

(h) By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  :

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the

roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

(i) If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$



and analogously

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } \boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)}$  where  $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$  are defined respectively

Then the solution satisfies the inequalities

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t}$$

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \right) \leq G_{30}(t) \leq \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :-

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

### **Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

(j)  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  :

- (k) By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations  
 $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$   
 and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  :

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$   
 and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

- (l) If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)}$  where  $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$  are defined respectively

Then the solution satisfies the inequalities

$$G_{32}^0 e^{((s_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(s_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((s_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(s_1)^{(6)}t}$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((s_1)^{(6)} - (p_{32})^{(6)}) - (s_2)^{(6)}} \left[ e^{((s_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(s_2)^{(6)}t} \right] + G_{34}^0 e^{-(s_2)^{(6)}t} \right) \leq G_{34}(t) \leq \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((s_1)^{(6)} - (a'_{34})^{(6)})} \left[ e^{(s_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}}$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})} \left[ e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$  :-

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

**Proof :** From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left( (a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a'_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

**Definition of**  $v^{(1)}$  :-  $v^{(1)} = \frac{G_{13}}{G_{14}}$

It follows

$$- \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$  :-

(a) For  $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t}}{1 + (C)^{(1)} e^{-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t}}, \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t}}{1 + (\bar{C})^{(1)} e^{-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t}}, \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

(b) If  $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$  we find like in the previous case,

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t}}{1 + (C)^{(1)} e^{-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t}}{1 + (\bar{C})^{(1)} e^{-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t}} \leq (\bar{v}_1)^{(1)}$$

(c) If  $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$ , we obtain

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (C)^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(1)}(t)$  :-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(1)}(t)$  :-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If  $(a''_{13})^{(1)} = (a''_{14})^{(1)}$ , then  $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$  and in this case  $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$  if in addition  $(v_0)^{(1)} = (v_1)^{(1)}$  then  $v^{(1)}(t) = (v_0)^{(1)}$  and as a consequence  $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$  this also defines  $(v_0)^{(1)}$  for the special case

Analogously if  $(b''_{13})^{(1)} = (b''_{14})^{(1)}$ , then  $(\tau_1)^{(1)} = (\tau_2)^{(1)}$  and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$  if in addition  $(u_0)^{(1)} = (u_1)^{(1)}$  then  $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$  This is an important consequence of the relation between  $(v_1)^{(1)}$  and  $(\bar{v}_1)^{(1)}$ , and definition of  $(u_0)^{(1)}$ .

we obtain

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left( (a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

**Definition of**  $v^{(2)}$  :- 
$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

It follows

$$- \left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$  :-

(d) For  $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}(v_1)^{(2)} - (v_0)^{(2)}]t}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}(v_1)^{(2)} - (v_0)^{(2)}]t}}, \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

(e) If  $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$  we find like in the previous case,

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} (v_1)^{(2)} - (v_2)^{(2)}] t}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} (v_1)^{(2)} - (v_2)^{(2)}] t}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}} \leq (\bar{v}_1)^{(2)}$$

(f) If  $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$  , we obtain

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(2)}(t)$  :-

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(2)}(t)$  :-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

**Particular case :**

If  $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$  , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if  $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}$  , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of**  $v^{(3)}$  :-

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

From which one obtains

(a) For  $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of  $(\bar{v}_1)^{(3)}$  :-**

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

(b) If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case,

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

(c) If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$  , we obtain

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of  $v^{(3)}(t)$  :-**

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)} , \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(3)}(t)$  :-**

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)} , \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{20})^{(3)} = (a''_{21})^{(3)}$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} = (v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b''_{20})^{(3)} = (b''_{21})^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$

: From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :- 
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$  :-

(d) For  $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

(e) If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case,

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

(f) If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$ , we obtain

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$  **this also defines**  $(v_0)^{(4)}$  **for the special case** .

Analogously if  $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then  $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of**  $(u_0)^{(4)}$ .

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)}$  :-  $v^{(5)} = \frac{G_{28}}{G_{29}}$

It follows

$$- \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

(g) For  $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$



(h) If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (C)^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

(i) If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$ , we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (C)^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{28}''^{(5)}) = (a_{29}''^{(5)})$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$  **this also defines**  $(v_0)^{(5)}$  **for the special case .**

Analogously if  $(b_{28}''^{(5)}) = (b_{29}''^{(5)})$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of**  $(u_0)^{(5)}$ .

we obtain

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of**  $v^{(6)}$  :-  $v^{(6)} = \frac{G_{32}}{G_{33}}$

It follows

$$- \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-

(j) For  $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

(k) If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

(l) If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$  , we obtain

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)} , \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)} , \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{32})^{(6)} = (a''_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case .**

Analogously if  $(b''_{32})^{(6)} = (b''_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

We can prove the following

**Theorem 3:** If  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  are independent on  $t$ , and the conditions

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with  $(p_{13})^{(1)}, (r_{14})^{(1)}$  as defined, then the system

If  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  are independent on  $t$ , and the conditions

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

with  $(p_{16})^{(2)}, (r_{17})^{(2)}$  as defined are satisfied, then the system

If  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  are independent on  $t$ , and the conditions

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with  $(p_{20})^{(3)}, (r_{21})^{(3)}$  as defined are satisfied, then the system

If  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  are independent on  $t$ , and the conditions

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with  $(p_{24})^{(4)}, (r_{25})^{(4)}$  as defined are satisfied, then the system

If  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  are independent on  $t$ , and the conditions

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0 ,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{28})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with  $(p_{28})^{(5)}, (r_{29})^{(5)}$  as defined satisfied , then the system

If  $(a'_i)^{(6)}$  and  $(b'_i)^{(6)}$  are independent on  $t$  , and the conditions

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{32})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with  $(p_{32})^{(6)}, (r_{33})^{(6)}$  as defined are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0$$

has a unique positive solution , which is an equilibrium solution for

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27})]T_{24} = 0$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27})]T_{25} = 0$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27})]T_{26} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

(a) Indeed the first two equations have a nontrivial solution  $G_{13}, G_{14}$  if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

(a) Indeed the first two equations have a nontrivial solution  $G_{16}, G_{17}$  if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

(a) Indeed the first two equations have a nontrivial solution  $G_{20}, G_{21}$  if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

(a) Indeed the first two equations have a nontrivial solution  $G_{24}, G_{25}$  if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

(a) Indeed the first two equations have a nontrivial solution  $G_{28}, G_{29}$  if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

(a) Indeed the first two equations have a nontrivial solution  $G_{32}, G_{33}$  if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

**Definition and uniqueness of  $T_{14}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a'_i)^{(1)}(T_{14})$  being increasing, it follows that there exists a unique  $T_{14}^*$  for which  $f(T_{14}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

**Definition and uniqueness of  $T_{17}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(2)}(T_{17})$  being increasing, it follows that there exists a unique  $T_{17}^*$  for which  $f(T_{17}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$

**Definition and uniqueness of  $T_{21}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $T_{25}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $T_{29}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $T_{33}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

(e) By the same argument, the equations 92,93 admit solutions  $G_{13}, G_{14}$  if

$$\begin{aligned} \varphi(G) &= (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - \\ &[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0 \end{aligned}$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

(f) By the same argument, the equations 92,93 admit solutions  $G_{16}, G_{17}$  if

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi((G_{19})^*) = 0$

(g) By the same argument, the concatenated equations admit solutions  $G_{20}, G_{21}$  if

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$

(h) By the same argument, the equations of modules admit solutions  $G_{24}, G_{25}$  if

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

(i) By the same argument, the equations (modules) admit solutions  $G_{28}, G_{29}$  if

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

(j) By the same argument, the equations (modules) admit solutions  $G_{32}, G_{33}$  if

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G^*) = 0$

Finally we obtain the unique solution of 89 to 94

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$



$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0$  ,  $T_{17}^*$  given by  $f(T_{17}^*) = 0$  and

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0$  ,  $T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{25}^*$  given by  $\varphi(G_{27}) = 0$  ,  $T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$  ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$  ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)} (T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)} ((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)} ((G_{35})^*)]}$$

Obviously, these values represent an equilibrium solution

### ASYMPTOTIC STABILITY ANALYSIS

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{dG_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})G_{13} + (a_{13})^{(1)}G_{14} - (q_{13})^{(1)}G_{13}^*T_{14}$$

$$\frac{dG_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})G_{14} + (a_{14})^{(1)}G_{13} - (q_{14})^{(1)}G_{14}^*T_{14}$$

$$\frac{dG_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})G_{15} + (a_{15})^{(1)}G_{14} - (q_{15})^{(1)}G_{15}^*T_{14}$$

$$\frac{dT_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})T_{13} + (b_{13})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*G_j$$

$$\frac{dT_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})T_{14} + (b_{14})^{(1)}T_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*G_j$$

$$\frac{dT_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})T_{15} + (b_{15})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*G_j$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij}$$

taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17}$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17}$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^* T_{17}$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)}) T_{16}^* G_j$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)}) T_{17}^* G_j$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)}) T_{18}^* G_j$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{21}'')^{(3)}}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b_i'')^{(3)}}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^* T_{21}$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^* T_{21}$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^* T_{21}$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)}) T_{20}^* G_j$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)}) T_{21}^* G_j$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)}) T_{22}^* G_j$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{25}'')^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b_i'')^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^* T_{25}$$

$$\begin{aligned} \frac{dG_{25}}{dt} &= -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^*T_{25} \\ \frac{dG_{26}}{dt} &= -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^*T_{25} \\ \frac{dT_{24}}{dt} &= -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*G_j \\ \frac{dT_{25}}{dt} &= -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*G_j \\ \frac{dT_{26}}{dt} &= -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*G_j \end{aligned}$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial (b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\begin{aligned} \frac{dG_{28}}{dt} &= -((a'_{28})^{(5)} + (p_{28})^{(5)})G_{28} + (a_{28})^{(5)}G_{29} - (q_{28})^{(5)}G_{28}^*T_{29} \\ \frac{dG_{29}}{dt} &= -((a'_{29})^{(5)} + (p_{29})^{(5)})G_{29} + (a_{29})^{(5)}G_{28} - (q_{29})^{(5)}G_{29}^*T_{29} \\ \frac{dG_{30}}{dt} &= -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^*T_{29} \\ \frac{dT_{28}}{dt} &= -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*G_j \\ \frac{dT_{29}}{dt} &= -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*G_j \\ \frac{dT_{30}}{dt} &= -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*G_j \end{aligned}$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations(global) and neglecting the terms of power 2, we obtain

$$\begin{aligned} \frac{dG_{32}}{dt} &= -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \\ \frac{dG_{33}}{dt} &= -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \\ \frac{dG_{34}}{dt} &= -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \\ \frac{dT_{32}}{dt} &= -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^*G_j \\ \frac{dT_{33}}{dt} &= -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^*G_j \\ \frac{dT_{34}}{dt} &= -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^*G_j \end{aligned}$$

The characteristic equation of this system is

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ & \left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ & + \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ & \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & \left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & + \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) \left( (a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \right) \\ & \left. \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right) \right\} = 0 \end{aligned}$$

+

$$\begin{aligned} &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ & \left[ ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ & \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ & + \left( ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \end{aligned}$$

$$\begin{aligned} & \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \\ & \left( (\lambda)^{(2)} \right)^2 + \left( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \\ & \left( (\lambda)^{(2)} \right)^2 + \left( (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\ & + \left( (\lambda)^{(2)} \right)^2 + \left( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\ & + \left( (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) \left( (a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \\ & \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \} = 0 \end{aligned}$$

+

$$\begin{aligned} & (\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \{ (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \} \\ & \left[ \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\ & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \\ & + \left( (\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\ & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \\ & \left( (\lambda)^{(3)} \right)^2 + \left( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\ & \left( (\lambda)^{(3)} \right)^2 + \left( (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\ & + \left( (\lambda)^{(3)} \right)^2 + \left( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\ & + \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) \left( (a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^* \right) \\ & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \} = 0 \end{aligned}$$

+

$$\begin{aligned} & (\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)} \{ (\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \} \\ & \left[ \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\ & \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \\ & + \left( (\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)} \right) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \end{aligned}$$

$$\begin{aligned} & \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(24)}T_{25}^* + (b_{25})^{(4)}s_{(24),(24)}T_{24}^* \right) \\ & \left( ((\lambda)^{(4)})^2 + ( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} ) (\lambda)^{(4)} \right) \\ & \left( ((\lambda)^{(4)})^2 + ( (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} ) (\lambda)^{(4)} \right) \\ & + \left( ((\lambda)^{(4)})^2 + ( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} ) (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\ & + \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) \left( (a_{26})^{(4)}(q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)} G_{24}^* \right) \\ & \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(26)}T_{25}^* + (b_{25})^{(4)}s_{(24),(26)}T_{24}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left( (\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)} \right) \left\{ (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \right\} \\ & \left[ \left( ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)}(q_{28})^{(5)} G_{28}^* \right) \right] \\ & \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(29)}T_{29}^* + (b_{29})^{(5)}s_{(28),(29)}T_{29}^* \right) \\ & + \left( ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)})(q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)}(q_{29})^{(5)} G_{29}^* \right) \\ & \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(28)}T_{29}^* + (b_{29})^{(5)}s_{(28),(28)}T_{28}^* \right) \\ & \left( ((\lambda)^{(5)})^2 + ( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} ) (\lambda)^{(5)} \right) \\ & \left( ((\lambda)^{(5)})^2 + ( (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} ) (\lambda)^{(5)} \right) \\ & + \left( ((\lambda)^{(5)})^2 + ( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} ) (\lambda)^{(5)} \right) (q_{30})^{(5)} G_{30} \\ & + \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) \left( (a_{30})^{(5)}(q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)} G_{28}^* \right) \\ & \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(30)}T_{29}^* + (b_{29})^{(5)}s_{(28),(30)}T_{28}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left( (\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)} \right) \left\{ (\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)} \right\} \\ & \left[ \left( ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)}(q_{32})^{(6)} G_{32}^* \right) \right] \\ & \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T_{33}^* + (b_{33})^{(6)}s_{(32),(33)}T_{33}^* \right) \\ & + \left( ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)}(q_{33})^{(6)} G_{33}^* \right) \end{aligned}$$

$$\begin{aligned} & \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T_{33}^* + (b_{33})^{(6)}s_{(32),(32)}T_{32}^* \right) \\ & \left( ((\lambda)^{(6)})^2 + ( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} ) (\lambda)^{(6)} \right) \\ & \left( ((\lambda)^{(6)})^2 + ( (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} ) (\lambda)^{(6)} \right) \\ & + \left( ((\lambda)^{(6)})^2 + ( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} ) (\lambda)^{(6)} \right) (q_{34})^{(6)} G_{34} \\ & + \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left( (a_{34})^{(6)}(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G_{32}^* \right) \\ & \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T_{33}^* + (b_{33})^{(6)}s_{(32),(34)}T_{32}^* \right) \} = 0 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

### Acknowledgments:

The introduction is a collection of information from various articles, Books, News Paper reports, Home Pages Of authors, Journal Reviews, Nature 's Letters, Article Abstracts, Research papers, Abstracts Of Research Papers, Stanford Encyclopedia, Web Pages, Ask a Physicist Column, Deliberations with Professors, the internet including Wikipedia. We acknowledge all authors who have contributed to the same. In the eventuality of the fact that there has been any act of omission on the part of the authors, we regret with great deal of compunction, contrition, regret, trepidation and remorse. As Newton said, it is only because erudite and eminent people allowed one to piggy ride on their backs; probably an attempt has been made to look slightly further. Once again, it is stated that the references are only illustrative and not comprehensive

### REFERENCES

1. A HAIMOVICI: "On the growth of a two species ecological system divided on age groups". Tensor, Vol 37 (1982), Commemoration volume dedicated to Professor Akitsugu Kawaguchi on his 80<sup>th</sup> birthday
2. FRTJOF CAPRA: "The web of life" Flamingo, Harper Collins See "Dissipative structures" pages 172-188
3. HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems (EOLSS), (Eolss Publishers, Oxford) [<http://www.eolss.net>]
4. MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204, doi:10.1029/2005JD006097
5. STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" J. Atmos. Sci., 53, 980-1006
6. FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010
7. Paul Allen Tipler, Ralph A. Llewellyn (2003-01), *Modern Physics*, W. H. Freeman and Company,



pp. 87–88, ISBN 0-7167-4345-0

8. Rainville, S. et al. World Year of Physics: A direct test of  $E=mc^2$ . *Nature* 438, 1096-1097 (22 December 2005) | doi: 10.1038/4381096a; Published online 21 December 2005.
9. F. Fernflores. The Equivalence of Mass and Energy. Stanford Encyclopedia of Philosophy
10. Relativity DeMystified, D. McMahon, Mc Graw Hill (USA), 2006, ISBN 0-07-145545-0
11. Dynamics and Relativity, J.R. Forshaw, A.G. Smith, Wiley, 2009, ISBN 978-0-470-01460-8
12. Hans, H. S.; Puri, S. P. (2003). *Mechanics* (2 ed.). Tata McGraw-Hill. p. 433. ISBN 0-07-047360-9., Chapter 12 page 433
13. E. F. Taylor and J. A. Wheeler, **Spacetime Physics**, W.H. Freeman and Co., NY. 1992. ISBN 0-7167-2327-1, see pp. 248-9 for discussion of mass remaining constant after detonation of nuclear bombs, until heat is allowed to escape.
14. Mould, Richard A. (2002). *Basic relativity* (2 ed.). Springer. p. 126. ISBN 0-387-95210-1., Chapter 5 page 126
15. Chow, Tail L. (2006). *Introduction to electromagnetic theory: a modern perspective*. Jones & Bartlett Learning. p. 392. ISBN 0-7637-3827-1., Chapter 10 page 392
16. Cockcroft-Walton experiment
17. Earth's gravitational self-energy is  $4.6 \times 10^{-10}$  that of Earth's total mass, or 2.7 trillion metric tons. Citation: *The Apache Point Observatory Lunar Laser-Ranging Operation (APOLLO)*, T. W. Murphy, Jr. et al. University of Washington, Dept. of Physics (132 kB PDF.).
18. There is usually more than one possible way to define a field energy, because any field can be made to couple to gravity in many different ways. By general scaling arguments, the correct answer at everyday distances, which are long compared to the quantum gravity scale, should be *minimal coupling*, which means that no powers of the curvature tensor appear. Any non-minimal couplings, along with other higher order terms, are presumably only determined by a theory of quantum gravity, and within string theory, they only start to contribute to experiments at the string scale.
19. G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Physical Review D* 14:3432–3450 (1976).
20. A. Belavin, A. M. Polyakov, A. Schwarz, Yu. Tyupkin, "Pseudoparticle Solutions to Yang Mills Equations", *Physics Letters* 59B:85 (1975).
21. F. Klinkhammer, N. Manton, "A Saddle Point Solution in the Weinberg Salam Theory", *Physical Review D* 30:2212.
22. Rubakov V. A. "Monopole Catalysis of Proton Decay", *Reports on Progress in Physics* 51:189–241 (1988).
23. S.W. Hawking "Black Holes Explosions?" *Nature* 248:30 (1974).
24. Lev B. Okun, *The concept of Mass*, *Physics Today* 42 (6), June 1969, p. 31–36, [http://www.physics.today.org/vol-42/iss-6/vol42no6p31\\_36.pdf](http://www.physics.today.org/vol-42/iss-6/vol42no6p31_36.pdf)
25. Max Jammer (1999), *Concepts of mass in contemporary physics and philosophy*, Princeton University Press, p. 51, ISBN 0-691-01017-X

26. Eriksen, Erik; Vøyenli, Kjell (1976), "The classical and relativistic concepts of mass", *Foundations of Physics* (Springer) **6**: 115–124, Bibcode 1976FoPh....6..115E, DOI:10.1007/BF00708670
27. Janssen, M., Mecklenburg, M. (2007), *From classical to relativistic mechanics: Electromagnetic models of the electron.*, in V. F. Hendricks, et al., *Interactions: Mathematics, Physics and Philosophy* (Dordrecht: Springer): 65–134
28. Whittaker, E.T. (1951–1953), *2. Edition: A History of the theories of aether and electricity, vol. 1: The classical theories / vol. 2: The modern theories 1900–1926*, London: Nelson
29. (Miller, Arthur I. (1981), *Albert Einstein's special theory of relativity. Emergence (1905) and early interpretation (1905–1911)*, Reading: Addison–Wesley, ISBN 0-201-04679-2
30. Darrigol, O. (2005), "The Genesis of the theory of relativity." (PDF), *Séminaire Poincaré* **1**: 1–22
31. Philip Ball (Aug 23, 2011). "Did Einstein discover  $E = mc^2$ ?". *Physics World*.
32. Ives, Herbert E. (1952), "Derivation of the mass-energy relation", *Journal of the Optical Society of America* **42** (8): 540–543, DOI:10.1364/JOSA.42.000540
33. Jammer, Max (1961/1997). *Concepts of Mass in Classical and Modern Physics*. New York: Dover. ISBN 0-486-29998-8.
34. Stachel, John; Torretti, Roberto (1982), "Einstein's first derivation of mass-energy equivalence", *American Journal of Physics* **50** (8): 760–763, Bibcode1982AmJPh..50..760S, DOI:10.1119/1.12764
35. Ohanian, Hans (2008), "Did Einstein prove  $E=mc^2$ ?", *Studies In History and Philosophy of Science Part B* **40** (2): 167–173, arXiv:0805.1400, DOI:10.1016/j.shpsb.2009.03.002
36. Hecht, Eugene (2011), "How Einstein confirmed  $E_0=mc^2$ ", *American Journal of Physics* **79** (6): 591–600, Bibcode 2011AmJPh..79..591H, DOI:10.1119/1.3549223
37. Rohrlich, Fritz (1990), "An elementary derivation of  $E=mc^2$ ", *American Journal of Physics* **58** (4): 348–349, Bibcode 1990AmJPh..58..348R, DOI:10.1119/1.16168
38. (1996). *Lise Meitner: A Life in Physics*. California Studies in the History of Science. **13**. Berkeley: University of California Press. pp. 236–237.
39. "consciousness". Merriam-Webster. Retrieved June 4, 2012.
40. Robert van Gulick (2004). "Consciousness". *Stanford Encyclopedia of Philosophy*.
41. Farthing G (1992). *The Psychology of Consciousness*. Prentice Hall. ISBN 978-0-13-728668-3.
42. John Searle (2005). "Consciousness". In Honderich T. *The Oxford companion to philosophy*. Oxford University Press. ISBN 978-0-19-926479-7.
43. Susan Schneider and Max Velmans (2008). "Introduction". In Max Velmans, Susan Schneider. *The Blackwell Companion to Consciousness*. Wiley. ISBN 978-0-470-75145-9.
44. Güven Güzeldere (1997). Ned Block, Owen Flanagan, Güven Güzeldere. ed. *The Nature of Consciousness: Philosophical debates*. Cambridge, MA: MIT Press. pp. 1–67.
45. J. J. Fins, N. D. Schiff, and K. M. Foley (2007). "Late recovery from the minimally conscious state: ethical and policy implications". *Neurology* **68**: 304–307. PMID 17242341.
46. Locke, John. "An Essay Concerning Human Understanding (Chapter XXVII)". Australia: University of Adelaide. Retrieved August 20, 2010.

47. "Science & Technology: consciousness". Encyclopedia Britannica. Retrieved August 20, 2010.
48. Samuel Johnson (1756). *A Dictionary of the English Language*. Knapton.
49. (C. S. Lewis (1990). "Ch. 8: Conscience and conscious". *Studies in words*. Cambridge University Press. ISBN 978-0-521-39831-2.
50. Thomas Hobbes (1904). *Leviathan: or, the Matter, Forme & Power of a Commonwealth, Ecclesiastical and Civill*. University Press. p. 39.
51. James Ussher, Charles Richard Elrington (1613). *The whole works, Volume 2*. Hodges and Smith. p. 417.
52. James Hastings and John A. Selbie (2003). *Encyclopedia of Religion and Ethics Part 7*. Kessinger Publishing. p. 41. ISBN 0-7661-3677-9.
53. G. Melenaar. *Mnemosyne, Fourth Series*. **22**. Brill. pp. 170–180.
54. Boris Hennig (2007). "Cartesian Conscientia". *British Journal for the History of Philosophy* **15**: 455–484.
55. Sara Hein ämaa (2007). *Consciousness: from perception to reflection in the history of philosophy*. Springer. pp. 205–206. ISBN 978-1-4020-6081-6.
56. Stuart Sutherland (1989). "Consciousness". *Macmillan Dictionary of Psychology*. Macmillan. ISBN 978-0-333-38829-7.
57. Justin Sytsma and Edouard Machery (2010). "Two conceptions of subjective experience". *Philosophical Studies* **151**: 299–327. DOI: 10.1007/s11098-009-9439-x.
58. Gilbert Ryle (1949). *The Concept of Mind*. University of Chicago Press. pp. 156–163. ISBN 978-0-226-73296-1.
59. Michael V. Antony (2001). "Is consciousness ambiguous?". *Journal of Consciousness Studies* **8**: 19–44.
60. Max Velmans (2009). "How to define consciousness—and how not to define consciousness". *Journal of Consciousness Studies* **16**: 139–156.
61. Ned Block (1998). "On a confusion about a function of consciousness". In N. Block, O. Flanagan, G. Guzeldere. *The Nature of Consciousness: Philosophical Debates*. MIT Press. pp. 375–415. ISBN 978-0-262-52210-6.
62. Daniel Dennett (2004). *Consciousness Explained*. Penguin. pp. 375.

**First Author:** <sup>1</sup>**Mr. K. N.Prasanna Kumar** has three doctorates one each in Mathematics, Economics, Political Science. Thesis was based on Mathematical Modeling. He was recently awarded D.litt. for his work on 'Mathematical Models in Political Science'--- Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Corresponding **Author:drknpkumar@gmail.com**

**Second Author:** <sup>2</sup>**Prof. B.S Kiranagi** is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150 publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent publication history.-- UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

**Third Author:** <sup>3</sup>**Prof. C.S. Bagewadi** is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. Several Books on Differential Geometry, Differential Equations are coauthored by him--- Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

### **IISTE Knowledge Sharing Partners**

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

