

Detection of *Histoplasma capsulatum* in Bats Dropping by using Polymerase Chain Reaction(PCR) at the First Time in Iraq

Neeran Obied Jasim

Clinical Laboratory Science-Pharmacy College-AL-Qadisiyia University

Email: Neranasim@yahoo.com

Abstract

In the present we isolated the fungus *Histoplasma capsulatum* as a first time from bats dropping in AL-Qadisiyah province .Results show that the macro and micro characteristics referred that these characteristics was related to the fungus *Histoplasma capsulatum* ,also this result was confirmed by PCR technique.

Keywords: *Histoplasma capsulatum* , bats dropping, PCR

Introduction:

Histoplasma capsulatum is a fungal pathogen that can result in a wide range of clinical presentations, from asymptomatic through fatal infection. It usually causes lung disease called Histoplasmosis or Darling's disease. It is called Darling's disease because it was found by Samuel Darling in histopathological specimens about a century ago.(Robert,2001) *Histoplasma capsulatum* is a biologically interesting inhabitant of soil and mammalian hosts, a clinically significant cause of respiratory and systemic infection, and an excellent fungal model of dimorphic cell development and facultative intracellular pathogenesis. (Chang Ryan, 2007) *H. capsulatum* is unique in its dimorphism. Dimorphism allows *H. capsulatum* to infect mammals by going through three significant development stages depends on the temperature shift from 25 °C to 37 °C.(Bossche,1993) In the moist soil that is rich in bird or bat guano at temperature about 25 °C, *H. capsulatum* exists in a filamentous mycelia form. However, when humans inhale *H. capsulatum* into their respiratory tracks, in order to replicate its DNA in the host at 37C, the pathogen has to be able to convert its tissue from one form to another. In this case, *H. Capsulatum* changed from fungi to yeast when it's growing in the human bodies (Kauffman, 2007). *Histoplasmosis capsulatum* is found throughout the world. It is endemic in certain areas of the United States, particularly in states bordering the Ohio River valley and the lower Mississippi River. The humidity and acidity patterns of soil are associated with endemicity. Bird and bat droppings in soil promote growth of *Histoplasma*. Contact with such soil aerosolizes the micro conidia, which can infect humans. It is also common in caves in southern and East Africa. (Ryan and Ray, 2004). Regarding the presence of the fungus in Iraq is not registered according available references we have and that the current study is the first in the country to record the presence of this fungus.

Material &Methods:

Culture media: The following media were used for isolated and identification of fungus from samples (Beneke and Rogers,1980):

*Sabouraud's Dextrose Agar &chloramphenicol

*Brain Heart Infusion Agar with cycloheximide &chloramphenicol

*Salvin's YP medium

Collection of sample : Samples of bats dropping were collected from bats nests were it's found .These samples were placed in sterile taines and transported to laboratory for culturing and testing .

Isolation &Identification of fungus: By using of method that reported by Lopez-Martines &Castaron-Oliver(1995) was isolated the fungus from samples that collected .make suspension from the samples and cultured on the media that reported above .After the incubator period (at 25C for two weeks and at 37C for seven days) the fungus was identification as a *Histoplasma capsulatum* according to macro and micro features that reported by (Ellis ,1994; Larone,1995;St-Germain & Summerbell,1996 ;Frey *et.al.*,2002)

PCR Technique:


Extract DNA: the DNA extracted from the isolates that selective on SDA medium, and Then followed the steps installed that supplied with the kit fitted from the company (Bioneer) to extract DNA, then blending part of it with dye Ethidium bromide on gel Agarose record in a concentration of 1.5% (100 ml buffer solution TEB Buffer concentration (1X) Molar according of the company (Bio Basic) for the purpose of purity way electric deportation .

Prepare the reaction solution to a polymerase chain reactions PCR:

attended the reaction mixture (size 50 Micro liters) for a polymerase **chain** reactions following the steps installed and supplied with the kit AccuPower ® PCR PerMix processed from the company (Bioneer) and by using of the *H. capsulatum*-specific primers HC-1 and HC-2 (HC-1: 5-GAGCCTCTGACCGGGAC-3;HC-2: 5-GCACGTCCCAC CCGGTCAG-3). (White *et.al.*,1990)

Results&Discussion:

Dimorphic fungi, such as *Histoplasma*, undergo a temperature induced transition between growth phases, which means that it has two distinct phenotypes. This dimorphism makes these organisms difficult to correctly identify using traditional microbiological methods. At ambient temperatures below 30°C in contaminated soils, *Histoplasma* grows in its mycelial form with the appearance of typical molds. Once the spores or micro conidia have been inhaled, the organism then undergoes a metamorphosis to its yeast or parasitic form. It is at this stage that *Histoplasma* becomes virulent. At normal body temperatures, around 37°C, the organism looks and behaves as yeast with the typical budding off of cells. This phase change, induced by the elevated temperatures of the body, is required for virulence. (Benekli,2004 ; Robert Brooks ,2011). Fig.(1) show colony with slow growing and cottony appearance , white in color .On other hand, Fig.(2) show septate hyphae with macro conidia, or tuberculate conidia, are 9 m in diameter . Identification of the tuberculate macro conidia allows to diagnosis of *Histoplasma* (Bhatti, *et.al.*, 2004) Also, this result accordant with (Kauffman, 2003) that reported , the tuberculate conidia, are 8 to 15 m in diameter and have distinctive projections on their surface. Fig (3) show moist, folded creamy colony at 37C ,that represented parasitic phase and fig(4) show , oval , unicellular budding yeast cells and buds arise at narrow neck . This results reported by (Beneke ,2004) whose reported that conversion to the yeast phase at 37C on brain heart infusion agar are necessary criteria for identification of *Histoplasma capsulatum* . Many species of fungi other than *H. capsulatum* produce similar colony and sporulation characteristics. Some examples are *Blastomyces dermatitidis* *Chrysosporium sp.*, and *Sepedonium sp.* Therefore, additional testing is needed to definitively identify the organism. One method is to convert the fungus colony from the filamentous, mycelial phase to the yeast phase by subculture to highly enriched cysteine -containing media and incubation at 35° to 37°C..(Wheat,1989).



Fig(1) White cottony colony on SDA(at 28C)





Fig.(2)Tuberculated macroconidia &small microconidia(640X)



Fig(3) Folded creamy colonyon BHI (at 37C)




Fig.(4)yeast cells with capsule (640X)

DNA based PCR technology, which uses probes specific to these organisms, is a quicker, ore accurate and cost-effective way of determining their presence or absence .Sequences that are found only in the genome of *Histoplasma* are targeted by the probes, allowing a precise, definitive result and do not require a viable organism for detection.(Hopfer,*et.al.*, 1993 ; Reid and Schafer ,1999).Fig.(5) show , Polymerase chain reaction method (PCR) have been used to detection and identification of fungus , gave rise to a single 402 bp PCR product .This result accordant with Reid and Schafer (1999) that reported A polymerase chain reaction (PCR) method was developed that allows the direct detection of *H. capsulatum* in soil and gave to a single 400 bp PCR product diagnostic of *H. cap.sulatum* .




Fig (5):gel electrophoresis that the results of polymerase chain reaction to DNA gene diagnosis of *Histoplasma* sp. Where is the M: Marker ladder 100-1500 bp and samples of No. 1-4 represent positive samples for testing an output length of 402bp.

Conclusion

The data presented in this study for our knowledge is considered the first screening data regarding the incidence of *Histoplasma* in Iraq and the high incidence was observed in bats dropping samples. PCR technique have important role of diagnosis of this fungus and this assay that correctly identified *H. capsulatum* from among a variety of fungi grown in the laboratory appears promising.

References

- Beneke, E and Rogers, 1980. Medical Mycology Manual with human mycosis monograph. 4th ed. Burgess pub. comp.
- Benekli, M. 2004. Pulmonary histoplasmosis in patients with a chronic obstructive pulmonary disease. Diagnosis J. Thrac. Cardiovasc. 127:585-586.
- Bhatti, S., L. Vilenski, R. Tight, and R. A. Smego, Jr. 2004. *Histoplasma* endocarditis: clinical and mycologic features and outcomes. J. Infect. 51: 2-9
- Bossche, Hugo, Frank Odds, and David Kerridge. 1993. Dimorphic Fungi in Biology and Medicine. 1st ed. New York and London: Plenum Press. Chang Ryan. 2007. Histoplasmosis. medicine. 10 (1): 114-122.
- Ellis, D.H. 1994. Clinical Mycology: The Human Opportunistic Mycosis. Gillingham Printers Ltd. Australia.
- Frey, D., Oldfield, R. & Bridger, R. 2002. A color atlas of pathogenic fungi. Wolfe Medical Publication P.285.
- Hopfer, R. L., Walden, P., Setterquist, S. & Highsmith, W. E. 1993. Detection and differentiation of fungi in clinical specimens using polymerase chain reaction (PCR) amplification and restriction enzyme analysis. Journal of Medical Veterinary Mycology 31, 65-75
- Kauffman, C. A. 2003. Histoplasmosis, p. 285-298. In W. E. Dismukes, P. G. Pappas, and J. D. Sobel (ed.), Clinical mycology. Oxford University Press, New York, NY
- Kauffman, C.A. 2007. "Histoplasmosis: a clinical and laboratory update. *Clinical Microbiology Reviews*. 20 (1): 115-132.
- Larone, D.H. 1995. Medically important fungi - A guide to identification. 3rd ed. ASM Press, Washington, D.C.
- Lopez-Martinez, M. & Castaron-Oliveres, L. 1995. Isolation of *Cryptococcus neoformans* from birds dropping, fruits and vegetable. Mycopathologia, 129:25-28.
- Reid, T. M and M. P. Schafer. 1999. Direct detection of *Histoplasma capsulatum* in soil suspensions by two-stage PCR. Molecular and Cellular Probes. 13, 269-273
- Rippon, J. W. 1988. Histoplasmosis. p. 381-423. In Medical mycology, the pathogenic fungi and the pathogenic actinomycetes. 3rd ed. W. B. Saunders Company.
- Robert Brooks. 2011. PCR detection of *Histoplasma capsulatum* in environmental samples. Microbac. 7:14-18.
- Robert, G. 2001. Environmental Health Services: Systemic fungal infections. Colorado State University
- Ryan, K.J., Ray, C.G. 2004. *Sherris Medical Microbiology* (4th ed.). McGraw Hill. pp. 674-6
- St-Germain, G. & Summerbell, R. 1996. Identification filamentous fungi: A clinical laboratory hand book. 1st ed. Star publishing company, Belmont, California.
- Wheat, L. J. 1989. Diagnosis and Management of Histoplasmosis. Eur. J. Clin. Microbiol. Infect. Dis. 8:480-490.
- White, T. J., Bruns, T., Lee, S. & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetic. In PCR Protocols—a Guide to Methods and Applications (Innis, M. A., Gelfand, D. H., Sninsky, J. J. & White, T. J., Journal eds) pp. 315-22. San Diego: Academic Press
- Woods, J. P., E. L. Heinecke, J. W. Luecke, E. Malsonado, J. Z. Ng, D. M. Retallack, and M. M. Timmerman. 2001. Pathogenesis of *Histoplasma capsulatum*. Semin. Respir. Infect. 16:91-10

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

