

# The Effect of Resveratrol on Swarming Differentiation and the Expression of Some Virulence Factors in *Proteus vulgaris*

Mohammed Ghaidaa<sup>1\*</sup>, Wang Yanchang<sup>1\*</sup>, and Abdallah K. Hindi<sup>2\*</sup>

<sup>1</sup>Department of Biomedical Sciences, College of Medicine,  
Florida State University, Tallahassee, Florida, 32306, USA

<sup>2</sup>Microbiology Department of Science College, Babylon University, Babel, Iraq

## Abstract

Resveratrol (3,5,4'-trihydroxy-*trans*-stilbene) is a stilbenoid, a type of natural phenol, and a phytoalexin with anti-inflammatory and antioxidant activities. It is produced naturally by several plants especially the roots of the Japanese Knotweed when under attack by pathogens such as bacteria or fungi. In this study we have verified that resveratrol has activity against *Proteus vulgaris*, an important pathogen infecting the urinary tract by investigating its effect on swarming and some virulence factor expression (haemolysin and urease). Swarming inhibition was determined on Luria Bertani agar with and without resveratrol and then bacteria was harvested to assay cell length and the production of haemolysin and urease. Resveratrol significantly inhibited swarming and virulence factor expression but its effect on growth rate was not significant.

**Keywords:** Resveratrol, *Proteus vulgaris*, phytoalexin, haemolysin, urease.

## 1. Introduction

*Proteus vulgaris* is a rod-shaped, Gram negative bacterium that inhabits the intestinal tracts infection of humans and animals. It can also be found in soil, water and fecal matter. It is grouped with the enterobacteriaceae and as an opportunistic pathogen of human, it is known to cause urinary tract infections and wound infections [1]. Individuals suffering from urinary tract infections caused by *Proteus spp.* often develop bacteriuria, cystitis, kidney and bladder stones, catheter obstruction due to stone encrustation, acute pyelonephritis, and fever [2]. Several potential virulence factors including haemolysin, swarming, adhesions, proteases, and ureases, may be responsible for the pathogenicity of *P. vulgaris*. The expression of virulence factors and the ability to invade human urothelial cells is coordinately up regulated during swarming [3,4 and 5].

Resveratrol is a phytoalexin found in grapes, wine, peanuts, cranberries, strawberries, and some other botanical sources. The discovery of resveratrol was occurred in 1940, but the first study to show the beneficial effects of resveratrol on human health was conducted in the 1990. Since then, several papers have been published annually elucidating the benefits of this molecule. Resveratrol has a wide range of biological activities and consequently it has many different targets and mechanisms of action. Resveratrol can prevent or slow the progression of several diseases; including cardiovascular, carcinogenic and neurodegenerative diseases [6,7]. Moreover, resveratrol has anti-inflammatory [8], antioxidant, and antimicrobial properties [8,9].

During the last century, antimicrobial agents have substantially reduced the threats associated with infectious diseases. The use of these drugs, combined with improvements in sanitation, housing and nutrition and the existence of comprehensive immunization programs, has allowed a radical reduction of untreatable infectious diseases, often fatal, contributing to increased life expectancy. However, the adaptation of microorganisms to antibiotics causes proliferation and persistence of drug resistance, currently a major public health problem, therefore it is urgent to discover new drugs endowed with antimicrobial activity [10,11].

In recent years, an increasing interest has been biologically active compounds including antioxidants from plants and other natural sources [13]. Thus, resveratrol, in addition to the biological activities described above, has been the subject of study for its ability to inhibit the growth of some pathogenic microorganisms such as Gram-positive and Gram-negative bacteria [14,15]. So, this study aimed to detect the effective resveratrol on the growth, swarming and virulence factor expression of *P. vulgaris*.

## 2. Methods

### 2.1. Resveratrol

Resveratrol was purchased from Indofine Chemical Company. The purity of this commercial compound is 99%.

### 2.2. Bacterial isolate and growth conditions

*P. vulgaris*, which used in this study was isolated from a Patient with a urinary tract infection and identified by biochemical tests [16]. For the bacterial growth assay, *P. vulgaris* was cultured overnight at 37°C in Luria Bertani (LB) broth, then diluted 1 in 100 in LB containing various concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60 µg/ml) and the growth rate was monitored at 1h intervals. For swarming differentiation and virulence factor assays, LB agar plates containing various concentrations of resveratrol were inoculated centrally with 5 µl of an overnight culture of *P. vulgaris* and incubated at 37°C for 7h. Bacterial cells taken from swarming plates were

suspended in 5ml of PBS; these cells were then used for morphology studies after Gram's staining, assays for urease and haemolysin production [4].

### 2.3. Swarming behavior assay

The swarming migration distance assay was performed as described previously [4,5]. Briefly, an overnight *P. vulgaris* culture (5 $\mu$ l) was inoculated centrally onto LB swarming agar plates (2 %w/v) with different concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60 $\mu$ g/ml). The plates were incubated at 37°C and the swarming migration distance was determined by measuring the swarm fronts of the bacterial cells at 7h after inoculation.

### 2.4. Measurement of growth rates

Overnight *P. vulgaris* culture was diluted 1:100 in fresh LB broth containing different concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60 $\mu$ g/ml). We also used other high concentrations of resveratrol (0.5, 1, 1.5, 2, and 2.5, 3mg/ml) to determine the effect on the growth inhibition of *P. vulgaris*. The growth rate was monitored as OD<sub>600</sub> at 1h intervals [5].

### 2.5. Measurement of cell length

Measurement of cell length was performed as described [5, 17]. Briefly, 150 $\mu$ l of stationary-phase LB cultures were spread onto LB agar plates without or with appropriate resveratrol and incubated at 37°C for various times. After incubation, cells from the entire surface of agar plates were harvested by washing into 5 ml of PBS. Bacterial cells were fixed and subjected to gram stain (Ward's Science, USA), examined by light microscopy (Carl Zeiss, Germany) at a magnification of 100X, and digitalized using a digital camera. The lengths of 100 cells in each sample were determined, and the average was calculated.

### 2.6. Haemolysin production assay

Haemolysin production was carried out by inoculating a blood agar medium containing 2% washed horse erythrocytes with bacterial cells taken after suspended in 5ml of PBS then incubated at 37°C for 24h. The appearance of clear zone around the colonies referred to a complete hemolysis ( $\beta$ -hemolysis). The appearance of greenish zone around the colonies referred to a partial hemolysis ( $\alpha$ -hemolysis), whereas no change of zone referred to non-hemolysis ( $\gamma$ -hemolysis) [4, 18, 19].

### 2.7. Urease production assay

Preparation of cells for urease assay was performed as described previously [3], in this test we inoculated the urea slant from bacterial suspension by streaking the entire slant surface, incubated the tubes with loosened caps at 37°C then color change of medium was examined after 16h incubation. Urease production was indicated by changed medium color into pink color [20].

### 2.8. Statistical analysis

All the results represent the average of three independent experiments. The data were presented as mean and analyzed by one-way analysis of variance with  $P < 0.05$  being significant, calculated using the Graph Pad Prism 5 statistical software [21].

## 3. Results

### 3.1. Inhibition of *P. vulgaris* swarming by resveratrol

After overnight incubation, the swarming behavior of *P. vulgaris* was monitored and we found that resveratrol has the ability to block the swarming migration of *P. vulgaris* in a dose-dependent manner (Fig. 1a). The swarming behavior was significantly inhibited at concentrations as low as 20 $\mu$ g/ml and was suppressed completely at 60  $\mu$ g/ml (Fig. 1a,b). The inhibitory effect of resveratrol on swarming might arise from a toxic effect on bacteria. To test this possibility, an overnight culture of *P. vulgaris* was inoculated into LB containing various concentrations of resveratrol and the growth rate of bacteria was monitored as shown in Figure 1c. The growth rate of *P. vulgaris* was inhibited slightly but not significantly because it grew in all tubes regardless of whether resveratrol was present or not, indicating that resveratrol could inhibit swarming but not the growth in *P. vulgaris*.




Fig.1.(a).Effect of resveratrol on the swarming of *P.vulgaris*.The histogram shows the migration distance of *P.vulgaris* in the presence of various concentrations of resveratrol(0, 10, 20, 30, 40, 50,60 µg/ml). The data represent the mean of three independent experiments and the differences are significant (P value <0.05).




Fig.1.(b). Halo images of swarming plates containing different concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60 µg/ml) at 7h after inoculation.




Fig.1.(c).Effect of resveratrol on the growth of *P.vulgaris*.OD<sub>600</sub> was measured overtime in the presence of various concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60 µg/ml). The data represent the mean of three independent experiments, there is no significant difference between concentrations (P value >0.05).

### 3.2.Inhibition of cell length and virulence factor expression in *P.vulgaris* by resveratrol

Cell morphology was monitored after inoculation of an overnight culture of *P.vulgaris* onto LB swarming plates containing various concentrations of resveratrol. As shown in figure 2(a), in the absence of resveratrol, the swarming cells were longer than the bacterial cells in the presence of resveratrol at the concentration 60µg/ml, suggesting that swarming differentiation was inhibited. The inhibition of differentiation started to be observed at a resveratrol concentration of 20µg/ml. Very few elongated swarming cells were observed at a resveratrol concentration of 50µg/ml. As the resveratrol concentration was increased to 60µg/ml, only short vegetative cells were observed. These results indicate that swarming differentiation of *P.vulgaris* was indeed inhibited by high concentrations of resveratrol.

To study whether the production of virulence factors (haemolysin and urease) was also influenced by resveratrol, the production of haemolysin and urease in *P. vulgaris* taken from LB agar plates containing different concentrations of resveratrol was determined. As shown in Figure 3, the production of virulence factors was not affected significantly at low resveratrol concentrations (0-40µg/ml) but was inhibited in the presence of increasing concentrations(50 and 60µg/ml).




Fig.2(a).Microscopic observation of *P.vulgaris* isolated from the LB plates without resveratrol (-) and with resveratrol 60ug/ml(+).




Fig.2.(b).Effect of resveratrol on the cell length of *P.vulgaris*.The histogram shows the cell length of *P.vulgaris* in the presence of various concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60µg/ml). The lengths of 100 cells in each sample were determined, and the average was calculated. The difference between concentrations is statistically significant (P value <0.05).




Fig.(3).The influence of resveratrol on the expression of virulence factors in *P.vulgaris*. The histogram shows the production of haemolysin and urease at different concentrations of resveratrol (0, 10, 20, 30, 40, 50, 60µg/ml). The data represent the mean of three independent experiments.The differences statistically significant (P value<0.05).

#### 4.Discussion

The emergence of bacterial strains that exhibit resistance to various antibiotics poses a major threat to public health. As a consequence, there is renewed interest in antibacterial targets which, by attenuating virulence, disrupt the capacity of pathogenic bacteria to cause infection[11].The purpose of the present study was to investigate the effect of plant extract (resveratrol) against uropathogenic *P. vulgaris*.

In this study, we found that resveratrol has the ability to inhibit *P. vulgaris* swarming significantly at a concentration as low as 20 µg/ ml and inhibited swarming completely at 60µg/ml (Fig.1a and b). Also, it had the ability to suppress the production of virulence factors(haemolysin and urease) at concentrations of 50 and 60µg/ml (Fig.3) but it did not significantly affect the growth of the bacteria at concentrations up to 60 µg/ml (Fig. 1c).Resveratrol did not affect the viability of *P. vulgaris* at a concentration of 3mg/ml(data not shown).

This means resveratrol could only inhibit swarming and virulence factor production without significant growth inhibition of *P. vulgaris*. Based on this finding we concluded that the swarming ability of *P. vulgaris* is correlated with its ability to express virulence factors and these results were similar to those reported by Allison *et al.* [3, 22]; where it has been shown that swarming differentiation of *P. mirabilis* and expression of virulence factors, such as urease, haemolysin and protease, are coordinately regulated in *P. mirabilis*.

The possible mechanism by which resveratrol could inhibit *P. vulgaris* swarming and virulence factor expression is by acting as an inhibitor compound for bacterial quorum sensing (QS). QS is the term used for the phenomenon of cell to cell communication in bacteria using secreted chemical signaling molecules called auto inducers. As environmental conditions often change rapidly, bacteria need to respond quickly in order to survive. QS is the regulation of gene expression in response to fluctuations in cell-population density and it enables bacteria to coordinate their behavior. Gram-positive and Gram-negative bacteria use quorum sensing communication circuits to regulate a diverse array of physiological activities. These processes include symbiosis, virulence, competence, conjugation, antibiotic production, motility, sporulation, and biofilm formation [23-24]. So, QS is considered a novel target for antimicrobial therapy.

The continuing emergence of multiple-drug-resistant strains of bacteria has necessitated finding novel strategies for treating bacterial infections and the discovery that a wide spectrum of organisms use quorum sensing to control virulence factor production makes it an attractive target for antimicrobial therapy. Among all the possibilities to inhibit the QS activity, the use of anti-QS compounds could be of great interest to avoid bacterial infections [25-26]. Such anti pathogenic compounds, in contrast with antibacterial compounds, do not kill bacteria or stop their growth and are assumed not to lead to the development of resistant strains [26,27]. Different mechanisms have been proposed to explain the interference of QS dependent processes by natural products. Some of these mechanisms are inhibition of signal molecule biosynthesis or acylated homoserine lactones (AHL) QS auto inducers reception (28,29), and the enzymatic inactivation and biodegradation of QS molecules (30). Therefore, we can conclude that the interference of QS control system has an anti-pathogenic effect and can be used in the treatment of bacterial infections. According to that, *Proteus spp.* swarming and virulence factor expression are generally believed to be regulated through a QS system which requires the sensing and integration of a variety of environmental, cell to cell and intracellular signals [31,32]. So the environmental changes or the presence of resveratrol in media of *P. vulgaris* has an effect on QS control system. The results from this study indicate that resveratrol has the potential to be an antimicrobial agent against *P. vulgaris* infection.

## 5. Acknowledgments

We are grateful to Biomedical Science Department of the Medicine College at Florida State University especially the laboratory of Dr. Wang. We are so grateful to Dr. Wang Yanchang, Dr. Fengshan Liang, Dr. Fengzhi Jin and Kelly McKnight for their assistance.

## References

- 1- Struble K. 2009. "Proteus Infections: Overview", *eMedicine*
- 2- Burall LS, Harro JM, Li X, Lockatell CV, Himpel SD, Hebel JR, Johnson DE, and HL Mobley. 2004. *Proteus mirabilis* genes that contribute to pathogenesis of urinary tract infection: identification of 25 signature-tagged mutants attenuated at least 100-fold. *Infect Immun.* **72**:2922–2938.
- 3- Liaw SJ, Lai HC, Ho SW, Luh KT, and WB Wang. 2000. Inhibition of virulence factor expression and swarming differentiation in *Proteus mirabilis* by P-nitrophenylglycerol. *J Med Microbiol.* **49**:725–731.
- 4- Liaw SJ, Lai HC, Ho SW, Luh KT, and WB Wang. 2001. Characterization of P-nitrophenylglycerol-resistant *Proteus mirabilis* super-swarming mutants. *J Med Microbiol.* **50**:1039–1048.
- 5- Liaw SJ, Lai HC, WB Wang. 2004. Modulation of swarming and virulence by fatty acids through the RsbA protein in *Proteus mirabilis*. *Infect. Immun.* **72**:6836–6845.
- 6- Jang JH and Surh YJ. 2000. Protective effect of resveratrol on beta-amyloid-induced oxidative PC12 cell death. *Free radical Biology and Medicine.* **34**: 1100-1110.
- 7- Zamin LL, Dillenburg-Pilla P, Argenta-Comiran R, Horn AP, Simao F, Nassif M, Gerhardt D, Frozza RL and Salbego C. 2006. Protective effect of resveratrol against oxygen-glucose deprivation in organotypic hippocampal slice cultures: Involvement of PI3-K pathway. *Neurobiology of Disease.* **24**: 170-182.
- 8- Baur JA and DA Sinclair. 2006. Therapeutic potential of resveratrol: the in vivo evidence. *Nature Reviews Molecular Cell Biology.* **5**: 493-506.
- 9- Docherty JJ, McEwen HA, Sweet TJ, Bailey E, and TD Booth. 2007. Resveratrol inhibition of *Propionibacterium acnes*. *Journal of Antimicrobial Chemotherapy.* **59**: 1182-1184.
- 10- Ferreira WFC and Sousa JCF. 2011. *Microbiologia Volume 1*. LIDEL; 1998. 1233.
- 19- Hinkelmann and Kempthorne. 2008. *Comparison of Treatments, Volume 1, Chapter 7*.
- 11- Butler MS and Buss AD. 2006. Natural products--the future scaffolds for novel antibiotics? *Biochemical*

Pharmacology. 71: 919-929.

12- Shan B, Cai YZ, Brooks JD ,andCorke H.2008.Antibacterial properties ofPolygonumcuspidatum roots and their major bioactive constituents. Food Chemistry.109: 530-537.

13- Tombola F, Campello S, De Luca L, Ruggiero P, Del Giudice G, PapiniE,andZoratti M.2003. Plant polyphenols inhibit VacA, a toxin secreted by the gastric pathogen *Helicobacter pylori*. FEBS Letters.543: 184-189.

14- MahadyGBandPendland SL.2000. Resveratrol inhibits the growth of *Helicobacter pylori* in vitro. AmericanJournalofGastroenterology. 95: 1849-1849.

15- Tegos G, Stermitz FR, Lomovskaya O, and Lewis K.2002. Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials. Antimicrobial Agents and Chemotherapy. 46: 3133-3141.

16--McFaddin JF .2000. Biochemical tests for identification of medical bacteria. 1<sup>st</sup> Ed.The Williams and Wilkins. Baltimore, USA.

17-Echeverrigaray S,Michelim L, DelamareAL,Andrade C, CostaSO ,and JZacaria.2008.The effect of Monoterpenes on swarming differentiation and Haemolysinactivity in *Proteus mirabilis*. Molecules.13:3107-3116.

18-Bulle N.2004.Bacteria from fish and other aquatic animals: A practical identification manual. Interpretation of biochemical identification test, 3, p121.

19-Ray C, George , Ryan, Kenneth J, Kenneth and Ryan .2004. Sherris Medical Microbiology: An Introduction to Infectious Diseases (4th ed.). McGraw Hill.p237.

20-Winn W, Allen S, Janda W, KonemanE, Procop G, SchreckenbergerP,andGWoods .2006.Koneman's color atlas and textbook of diagnostic microbiology, 6th ed. Lippincott Williams & Wilkins, Philadelphia, PA.

21- Montgomery.2001.The analysis of variance.Experiments with a single factor: Analysis of the fixed effects model.Section 3:3.

22-Allison C, Lai HC, and C Hughes.1992.Co-ordinate expression of virulence genes during swarm-cell differentiation and population migration of*Proteusmirabilis*.MolMicrobiol. 6:1583–1591.

23-DekievitTR andIglewski BH .2000.Bacterial quorum sensing inpathogenic relationships. Infect Immun. 68:4839–4849.

24-ZhangLH .2003.Quorum quenching and proactive host defense.Trends Plant Sci. 8:238–244.

25-AdonizioAL,Downum K, Bennett BC,andMathee K.2006. Anti-quorum sensing activity of medicinal plants in southern Florida. J. Ethnopharmacol.105:427–435.

26-Maria V, Alvarez, Maria R, Morier A, and P Alejandra .2012.Antiquorum Sensing and antimicrobial activity of natural agents with potential use in food. Journal ofFood Safety ISSN 1745-4565.

27-Fulghesu L, GiallorenzoC, and Savoia D.2007.Evaluation of different compounds as quorum sensing inhibitors in *Pseudomonas aeruginosa*.JChemother. 19(4):388-91.

28-Vattem D, Mihalik K, Crixell S.,and Mclean R.2007. Dietary phytochemicals as quorum sensing inhibitors.Fitoterapia.78(4):302-310

29-SturgillGand Rather PN .2004. Evidence that putrescine acts as an extracellular signal required for swarming in *Proteus mirabilis*. Mol. Microbiol. 51:437–446.

30-Defoirdt T, Boon N, Bossier Pand WVerstraete.2004. Disruption of bacterial quorum sensing: An unexplored strategy to fight infections in aquaculture. Aquaculture,240(1–4):69–88.

31-SturgillGand Rather PN .2004. Evidence that putrescine acts as an extracellular signal required for swarming in *Proteus mirabilis*.MolMicrobiol. 51:437–446.

32-RatherPN.2005.Swarmer cell differentiation in *Proteus mirabilis*.EnvironMicrobiol. 7:1065–1073.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

### CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

**Prospective authors of journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

### MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

### IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

