

Life Cycle and Abundance of Rice Leaf Folder, *Cnaphalocrocis medinalis* (Guenee) - A Review

Raveesh Kumar Gangwar

Lecturer and Head Department of Plant Protection, Faculty of Agriculture, Doon College of Education,
Saharanpur (U.P.) India
raveeshgangwar@yahoo.com

ABSTRACT

The Rice leaf folder, (RLF) *Cnaphalocrocis medinalis* (Guenee) is widely distributed in rice growing areas in all over the world. The leaf folder has been considered a miner pest of rice growing areas but after the existence, high yielding and Basmati rice varieties increasing the importance of these pests. Feeding rates of the first three larval instars rice leaf folder on rice plants were significantly lower than those of the fourth and fifth instars. The first three instars consumed less than 10% of the larva's total consumption. Feeding also decreased with increasing plant age. To understand about life cycle and management's tactics is help to reduce the yield losses. In present review articles discusses about life cycle and abundance of rice leaf folder.

Key word *Cnaphalocrocis medinalis*, life cycle, abundance

Introduction:

Rice (*Oryza sativa*) acts as a staple food for much of the global population (FAO 2004; Kumar S. A. V., et al., 2009) particularly in Asia where about 90% of people live on rice (Khush, G. S, Brar. D. S, 2002) Uttar Pradesh is responsible for the production of 14.5% of the country's rice (McCarthy, S., et al., 2008). However, in commercial production, rice productivity and quality are adversely affected by many biotic stresses, particularly insect pests. Insect pests damage rice crop at different stages of its growth. Yield losses in global rice output to pests (diseases, animal pests, and weeds), range from up to 20% to at least 30% of the attainable yield (Savary, et al., 2000b). For instance, some 500 species of insects and spiders may appear in a rice field in a particular season out of these, only few are potential threat. The stem borer, rice gandhi bug, brown plant hopper, white backed plant hopper, rice leaf folder etc cause damage to rice fields. Among them, leaf feeding insect have major importance because of their ability to defoliate or to remove the chlorophyll content of the leaves leading to considerable reduction in yield. Paddy leaf folder is one of the most important insect pests (Gunathilagaraj, K. and M. Gopalan, 1986).

Lepidopteron insect-pests are the main class of pests causing significant damage to crop plant yields. The rice leaf folder (RLF), *Cnaphalocrocis medinalis* (Guenee) (Lepidoptera:Pyralidae), is a predominant foliage feeder and one of the most destructive pests, affecting in all the rice ecosystems in Asia (Luo, S. J., 2010). Rice leaf folder was considered as pests of minor importance have increased in abundance in late 1980's and have become major pests in many parts of India. The yield loss is from 30 to 80 per cent due to leaf folder epidemic situation (Nanda, V. K. and R. C. Bisoi, 1990; Shah, et al., 2008). The rice leaf folder was considered as minor or sporadic pest in the past in many Asian countries. However, now it has been assumed as one of important insect pests and become a major threat to rice production in tropical and subtropical Asia. (Kushwaha and Singh, 1984; Shrivastava, 1989).

An increase in *C. medinalis* population could be attributed to the large scale cultivation of high yielding varieties, application of fertilizers, and continuous use of insecticides leading to outbreak of this pest in several countries, including India (Khan, Z. R., et al., 1988; Shanmugam T. R., et al 2006; Kaushik, C., 2010). The larvae fold the leaves longitudinally by stitching the leaf margins and feed by scraping the green mesophyll tissue from within the folded leaves. This feeding causes linear, pale white stripes that result in membranous patches. Out of the eight species of leaf folder, the most widespread and important one is *Cnaphalocrocis medinalis* cause significant losses to rice ecosystem (Bhatti, M. N. 1995). In their studies, 17.5% damaged leaves resulted in 16.5% yield loss, and 21.3% yield loss corresponded to 26.6% damaged leaves (Bautista, R. C., et al., 1984). Second instars leaf folder larvae glues the growing paddy leaves longitudinally for accommodation and feeds on green foliage voraciously which results in papery dry leaves (Khan, et al., 1989). A single larva can damage a number of rice leaves, disturbing photosynthesis and reducing the rice yield (Alvi S. M, et al., 2003). Loss incurred to the growing paddy crop is insurmountable (Ahmed et al., 2010). Feeding often results in stunting, curling or yellowing of plant green foliage (Alvi et al., 2003). The extent of loss may extend up to 63 to 80 percent depending on agro-ecological situations as reported by (Rajendran et al., 1986; Muragesan & Chellish, 1987). The heavy use of insecticides and high fertilizer rates seem to favors leaf-folder population outbreaks (Gottfried and Fallil 1986).

Bio- ecology of Rice leaf folder (*Cnaphalocrocis medinalis*):

The heavy use of (nitrogen) fertilizer persuades rapid multiplication of the leaf folder population in rice field. High humidity, high temperature and shady areas of the field favor their growth and developments. The presence of alternate host in rice fields and around the field provided shelter for insect and create favorable environment for complete several generation in one growing season. Expanded rice areas with irrigation systems, multiple rice cropping and insecticide induced resurgences are important factors in the leaf folder's abundance. The adults of leaf folder are nocturnal they hide in rice leaf during the day time to escape predation. They are active year-round and moths fly short distances when disturbed.

Life cycle of leaf folder:

For control of rice leaf folder there is very important to know about their life cycle because it's played a significant role to manage the population of *C. medinalis* in field and increased the yield.

Adult: Adult female moth is golden yellow in color with brown margins on both the wings. While resting, the body shape is like that of an equal-sided triangle. It has three bands traversing entire forewing. Middle band was short, comma like and curved outside. The adult moths are 10-12 mm long. Wing expanse is 13-15 mm. Male moths are smaller than female moths and have a prominent patch of dark brown scales along the mid Costa of forewing. They generally mate between dusk and midnight. Oviposition starts two-three nights after mating. Adults usually emerge in the evening and mating occurs at night. Pre-oviposition period of 3 days and oviposition period of 4 days was observed.

Egg: Eggs of leaf folder are transparent, yellowish white, 0.90 mm long and 0.39 mm wide, and almost flat with a slightly convex surface. The eggs hatch 3-4 days after oviposition. Eggs are laid singly or in groups of 3 to 8 along the midrib of young leaves. A female lays about 135 to 175 eggs.

Larva: Larvae are yellow, turn yellowish-green with brown head as they mature, and are about 12-25 mm long. Immediately after hatching, neonate larvae move to the unopened leaf at the center of the plant and feed gregariously by scraping the green matter. From second instars onwards, the larva folds the leaves by stitching with silken threads keeping open both the ends of the fold. Larval feeding by scraping the green matter staying within the fold results in the development of longitudinal white streaks. Sometimes 2-3 leaves were stitched together and larvae feed from within this fold. Larva turns pinkish white just before pupation. *C. medinalis* completes its larval development in 14 to 18 days after passing through five instars. The caterpillar secretes a series of threads and uses these to connect the two margins of a leaf blade. The threads contract as they dry and bring the two leaf margins together, turning the leaf blade into a tubular structure. The full-grown larva is yellowish green with a dark brown head and is 20-25 mm long and 1.5-2 mm wide.

Pupa: Pupa is light brown to bright brown and turns reddish-brown just before adult emergence. The larva cuts the leaf margin transversely up to the midrib, folds the cut portion of the leaf lamina over the other half and pupates inside the white silken webbing in leaf fold after stitching it completely on all sides. Pupation occurs mostly at the base of the plant and a single leaf was folded for pupation. The newly formed pupa is light brown, but turns reddish brown toward moth emergence in 6-10 days.

Nature of Damage:

The leaf folder damages the crop in its larval stage. Mostly Damage is caused by the Second instars larvae due to the folding of leaves and feeding on the green mesophyll tissue which caused the loss of output by 20% to 30% generally (Shen *et. al.*, 1988; Shepard *et. al.*, 1995). From within the folds which results in papery dry leaf (Chatterjee, P. B., 1979). However, the young larvae feed on open leaves but later feed inside the rolled leaf formed by folding the leaf longitudinally with a sticky substance. The larvae chew inside the fold by scraping the green matter. The scraped leaves become membranous, turn whitish and finally wither (Mishra, B. K., *et. al.*, 1998). This results in longitudinal white streaks causing reduced photosynthesis and plant vigor, ultimately affecting the plant growth and yield. At vegetative stage, crop generally recovers from the leaf folder damage but at reproductive stage, feeding damage on the flag leaf significantly reduces the grain filling resulting in yield loss. Generally one larva was found in each leaf fold and after feeding on that leaf it moves to another leaf. Thus single larvae feeds on a number of leaves can damage during its growth period. Heavily infested fields show many folded leaves and a scorched appearance of leaf blades. The damaged plants also pre-dispose the plants to fungal and bacterial infection. Severe infestations may annihilate the plant totally (Ramasubbaiah, K., *et. al.*, 1980). Losses that incurred to the growing paddy crop are insurmountable (Singh, H. M., *et al.*, 2003).

Seasonal occurrence and abundance:

Although moths are recorded in the warm tropics region around the years but during the rainy season they usually are most abundant. In cool regions, the insect is active from May to October, during these periods they completes four to five generations; the later generations usually overlap. High humidity and optimum temperature are the important factors for the insect's abundance. Every year the initial population migrates to these temperate countries from tropical regions. Leaf folders have importance both in upland as well as lowland rice growing fields in the last few decades, particularly in areas where modern and scented varieties are extensively grown. Expanded rice areas with new irrigation systems, multiple rice cropping, reduced genetic

variability of the modern semi dwarf varieties, application of high levels of nitrogenous fertilizers, and insecticide resistance and insect resurgence is increased the leaf folder problem.

References:

1. Ahmed, H., R. B. Khan, D. Sharma, V. V. S. Jamwal, and S. Gupta, 2010. Seasonal incidence, infestation and trap catch of *Cnaphalocrocis medinalis* (Guenee) in rice. *Annals Pl. Prot. Sci.* 18(2):38-383.
2. Alvi S. M, Ali M. A, Chaudhary S, Iqbal S (2003) Population trends and chemical control of rice leaf folder, *Cnaphalocrocis medinalis* on rice crop. *Int J Agri Biol* 5: 615–617.
3. 'Bautista, R. C., Heinrichs, E. A. and Rejesus, R. S. (1984) Economic injury levels for the rice leaf folder *Cnaphalocrocis medinalis* (Lepidoptera: Pyralidae). Insect infestation and artificial leaf removal. *Environ. Entomol.* 13:439-443.
4. Bhatti, M. N. 1995. Rice leaf folder (*Cnaphalocrocis medinalis*): A review. *Pak. Entomol.* 17: 126– 131.
5. Chatterjee, P. B., 1979. Rice leaf folder attack in India. *Intl. Rice Res News.* 4(3): 21.
6. Food and Agriculture Organization of the United Nations (FAO) (2004). The state of food security in the world. pp. 30–31.
7. Gunathilagaraj, K. and M. Gopalan, 1986. Rice Leaf folder Complex in Madurai, TN, India. *Intl. Rice Res. Notes* 11(6): 24.
8. <http://www.rkmp.co.in>
9. Kaushik C. 2010. Extent of damage by leaf folder, *Cnaphalocrocis medinalis* (Guenee) in paddy cultivars at Raiganj, Uttar Dinajpur, West Bengal. *Current Biotica* 4: 365–367.
10. Khan Z. R, Barrion A. T, Litsinger J. A, Castilla N. P, Joshi R. C. 1988. A bibliography of rice leaf folders (Lepidoptera: Pyralidae). *Insect Science and its Application* 9: 129–174.
11. Khan, M. R., M. Ahmad and S. Ahmad, 1989. Some studies on biology, chemical control and varietal preference of rice leaf folder, *Cnaphalocrocis medinalis*. *Pak. J. Agric. Sci.* 26: 253–63.
12. Khush G. S, Brar D.S, (2002) Biotechnology for rice breeding: progress and potential impact. In: Proc. 20th Session of the Int. Rice Commission: 23rd–26th July 2002; Thailand: Bangkok.
13. Kumar S. A. V, Jocqué, R and Sebastian, P. A (2009), "A new species of the ant spider genus *Suffasia* (Araneae: Zodariidae) from the Western Ghats, India with a key to the species of the genus", *Zootaxa*, Vol. 2203, pp. 59-64.
14. Kushwaha K. S, Singh R. 1984. Leaf folder (LF) outbreak in Haryana. *Int Rice Res Newsl*, 9(6): 20.
15. Luo, S.J. (2010) Occurrence of rice leaf roller in China and its identification and prevention. *Plant Diseases and Pests* 1: 13–18.
16. Maragesan, S. and S. Chellish, 1987. Yield losses and economic injury by rice leaf folder. *Indian J. Agri. Sci.*, 56: 282–5
17. McCarthy S, Singh D. D and Schiff H (2008), "Value Chain Analysis of Wheat and Rice in Uttar Pradesh, India", ACIDI/VOCA for World Vision, Lucknow, India.
18. Mishra, B. K., B. Senapati and P. R. Mishra, 1998. Chemical control of rice leaf folder, *Cnaphalocrocis medinalis* in Orissa. *J. Insect Sci.*, 11: 137-40.
19. Nanda, V.K. and R. C. Bisoi, 1990. Bionomics of rice L7 *C.M.* Orissa *J. Agrl. Res.*, 3(2): 130-135.
20. Rajendran, R., S. Rajendran and P. C. Sandra, 1986. Varietals resistance of rice of leaf folder. *Intl. Rice Res. News* 11: 17.
21. Savary, S., Willocquet, L., Elazegui, F. A., Castilla, N., Teng, P. S., 2000b. Rice pest constraints in tropical Asia: quantification of yield losses due to rice pests in a range of production situations. *Plant Dis.* 84, 357e369.
22. Shah, S., Ali, M., Rehman, U. H., Rehman, A., Abbasi, F. M., Khalil, I. H., and Ali, A. 2008. Characterization of wild rice species in response to leaf folder (*C. medinalis*). *Sarh. J. Agri.* 24(1):69-74.
23. Shanmugam T. R, Sendhil R, Thirumalvalavan V. 2006. Quantification and prioritization of constraints causing yield loss in rice (*Oryza sativa*) in India. *Agricultura Tropica et Subtropica* 39: 194–201.
24. Shen C. X, Zhang G. F, Zhao B. G, Xue J. J, 1988. The present situation of study of *Cnaphalocrocis medinalis* in the world. *China Plant Prot*, 3(10): 34–36. (in Chinese)
25. Shepard B. M, Barrion A. T, Litsinger J. A. 1995. Rice Feeding Insects of Tropical Asia. Manila, Philippines: IRRI: 70–90.
26. Shrivastava S. K. 1989. Leaf folder (LF) damage and yield loss on some of the selected rice varieties. *Int Rice Res Newsl*, 14(5): 10.
27. Singh S, Kushwaha K. S, Sharma P. D, 1995. Resurgence of rice leaf folder *Cnaphalocrocis medinalis* Guenee due to application of carbofuran granules. *Indian J. Ent.* 57, 366–372.
28. Singh, H. M., R. K. Srivastava, S. M. A. Rizvi, F. A. Elazegui, N. P. Castilla and S. Savary, 2003. Yield reduction due to brown spot and leaf folder injuries and various levels of fertilizers and water supply to rice crop. *Annals of Plant Prot. Sci.*, 11: 16-19.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

