

Impact of Flaxseed Consumption on Dyslipidemia

Kusumaneela Bolla, Santhi Sri .K.V

Department of Food and Nutritional Sciences, Acharya Nagarjuna University

P.Krishnaveni, Kusuma

Department of Appliednutrition,S.D.M.S.Mahila kalasala

Aim: The Present study is design to evaluate the effect of flaxseed consumption on lipid profile.

Methodology: Dyslipidemia subjects were selected and encouraged them to take 25gms of flaxseeds in their diet, anthropometric and biochemical parameters were collected before and after completion of study period.

Conclusion: By This study we are conduting that the intake of flaxseeds shows a significant reduction of the serum lipid($p < 0.001$) i.e: triglycerides, total cholesterol, low density lipoproteins.

Keywords: Dyslipidemia, Flaxseeds, Coronary Heart Diseases, Polyunsaturated Fatty Acids, Blood Pressure.

Introduction

Coronary heart disease(CHD) is a leading cause of morbidity and mortality in developed countries. These in the higher socioeconomic groups are the first ones to adopt to an adverse life style such as sedentary life style and intake high saturated fatty acids diet, cigarette smoking. They are also the first to revert to a healthier life style. Saturated fatty acid raise the levels of LDL and total blood cholesterol level. Both of these effect increases the risk of CHD. The activity of LDL receptors its ability to mediat the entry of LDL appears to be suppressed by saturated fatty acids. When LDL receptor activity decreases, LDL catabolism decreases and blood levels of LDL increases. Parallel changes in life style and diet have led to an increase in life expectancy and a greatly increases burden of CVD and other chronic disease.

Dyslipidemia is one of the modifiable risk factors of CHD, characterised by high triglycerides(TG), decreased high density lipoprotein cholesterol(HDL-C), increased lipoprotein cholesterol(LDL-C)and increased lipoproteins(LPa). LPa provides a carrier system for low density lipoprotein cholesterol facilitating cholesterol accumulation in cells. Blood concentration of antioxidants affects the susceptibility of LDL and lipoprotein(a) at oxidation, because it is the oxidation LDL that attracts circulating monocytes which then adhere to the arterial wall forming atherosclerotic plaque. Decreased level if these substance that protect against oxidation increase atherosclerotic risk. Although flaxseed act as a antioxidant, as flaxseed contains all sorts of healthy components and owes its primary health reputation to there of them: omega 3 fatty acids, lignans, fiber. Researchers at the Iowa state university nutrition and wellness research center found that cholesterol level lowered among men who included flaxseed in their diet. While flaxseed shown to reduce evidence that flaxseed and flaxseed oil may lower cholesterol levels. hence the present study aims to investigate the effect of flaxseed consumption on dyslipidemia

Methodology

In the present study 75 dyslipidemic subjects in the age of 40-60years were recruited. Dyslipidemic subjects with elevated levels(triglycerides, total cholesterol, LDL, HDL) who were visited hospital for first time as a regular health check ups. The methodology consisted a pre tested questionnaire for the subjects to collect general information, nutritional status i.e: height, weight, bmi, blood pressure, and other health issues, dietary pattern and life style. At the beginning of the study baseline values for blood pressure, body weight and BMI were recorded. Blood samples were collected from the selected subjects for triglycerides, total cholesterol, high density lipoproteins, low density lipoproteins was estimated by different procedures. After completion of supplementation period estimation of blood lipid profile, measurement of BMI was done.

Result

Mean reduction of 2.89% of body weight was noticed in the selected subjects. A significant difference($p < 0.01$) in the baseline and final value of the body weight and BMI of the subjects were observed. A significant decrease in the both systolic and diastolic blood pressure.

Table I

Effect of flaxseeds consumption on anthropometric and blood pressure measurements

Parameters	Before	After	Mean changes %	P value
Body weight	76.12	73.92	2.89	0.000
BMI	28.48	27.61	3.05	0.000
Systolic blood pressure(mm of hg)	135.76	134.56	0.88	0.044
Diastolic blood pressure (mm of Hg)	88	86	1.910	0.034

Table II
Effect of flaxseed consumption on lipid profile

Parameters	Before	After	Mean changes %
Total cholesterol	266.8	223	16.19
Triglycerides	192.68	165.7	13.9
HDL	37.2	39.1	4.94
LDL	134.6	107.8	19.93
VLDL	47.48	38.96	17.94

There is a significant reduction in the lipid profile by the supplementation of the flaxseeds for the period of 3 months in their diet.

Discussion

Dyslipidemia is a state that accelerates the beginning and development of atherosclerosis. Dyslipidemia can often be prevented or treated with life style management and proper nutrition. After supplementation of flaxseeds to dyslipidemic subjects indicated that interventions were well tolerated by subjects and the dropout rate was minimum. In the present study supplementation of flaxseeds to dyslipidemic subjects resulted in a significant decrease in body weight, BMI, systolic and diastolic blood pressure.

Regular intake of flaxseed in their diet of dyslipidemic subjects brought significant fall in the blood pressure, which may be attributed to the presence of PUFA, mainly ALA, soluble fiber, Lignans precursors and estrogen in flaxseeds. Intake of flaxseeds exhibited anti-hyperlipidemic effect. Conclusion of the highly significant reduction in total cholesterol, TG, LDL, VLDL levels and elevated HDL levels.

By our findings we are concluded that intake of regular flaxseeds shows the beneficial effect on the lipid profile and cardiac health.

Reference

1. Eriksson H, Pinsky JL, Larsson B, et al. the epidemiology of heart failure: the Framingham study. *J Am Coll Cardiol* 1993; 22:6A-13A
2. FOSSATI P, Prencipe L. Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clin Chem* 1982;28:2077-80.
3. McVeigh BL, Dillingham BL, Lampe JW, Duncan AM. Effect soy protein varying in isoflavone content on serum lipid in healthy young men. *Am J Clin Nutr* 2006;83:592-600.
4. Srilakshmi B. dietetics, New Age International (p) limited publisher; 2005, page: 377-385.
5. Charu katara, sonali saxena, amelioration of selected cardiac risk factors through supplementation of diet with flaxseed and soyabean, *IJN, pharmacology* oct-dec2013, vol3, issue4.
6. Lucas EA, Wild RD, Hammond LJ, Khalil DA, Juma S, Daggy BP, et al. flaxseed improves lipid profile without altering biomarkers of bone metabolism in postmenopausal women. *Indian J Clin Biochem* 2010;25:201-7.
7. Begg DP, Sinclair AJ, Stahl LA, Premaratna SD, Hafandi A, Jois M, et al. hypertension induced by omega 3 polyunsaturated fatty acid deficiency is alleviated by alpha - linolenic acid regardless of dietary source. *Hypertens res.* 2010;33:808-13.