

Analysis of Heavy Metals Found in Vegetables From Some Irrigated Gardens in Katsina, Daura and Funtua Metropolis, Katsina State, Nigeria.

Zahraddeen Hassan Yusuf^{1*} Muhammad Danjuma² Ibrahim Adam Tahir³

1. Department of Biology, School of Science Federal College of Education, Katsina State Nigeria.
2. Department of Biology, School of Science Federal College of Education, Katsina State Nigeria.
3. Department of Biology, School of Science Federal College of Education, Katsina State Nigeria.

* E-mail of the corresponding author: deeneecrc@gmail.com

Abstract

This study was conducted to determine the concentration of heavy metals cadmium (Cd) chromium (Cr) copper (Cu), iron (Fe) manganese (Mn) zinc (Zn), lead (Pb) and in five vegetables viz; (Bitter-leaf), Cabbage, Cucumber, Lettuce and Spinach and soil collected from selected farm and market sites in industrial, residential and commercial areas of Lagos state. The heavy metals present in the vegetables were analyzed using Atomic Absorption Spectrophotometer (AAS). The results reveal that all the heavy metals were detected in all the vegetables from the various sites. Most of the concentrations are below the World Health Organisation (WHO) and Food and Agriculture Organisation (FAO) safe limit of 0.1 mg/kg, 0.1 mg/kg, 40 mg/kg, 1.3 mg/kg, 1.5 mg/kg, 6.61 mg/kg, 3 mg/kg, 60 mg/kg for cadmium (Cd), cobalt (Co) (Cu), chromium (Cr), iron (Fe), lead (Pb) and zinc (Zn), and respectively in the vegetables. However the concentration of cobalt in spinach and lead in cabbage from Katsina are above the WHO and FAO safe limits. There is a significant differences in the concentration of Cd in the vegetables from the sampling sites ($P < 0.05$), while there is no significant difference in the concentration of Cr, Co, Cu, Fe, Mn, Pb, Zn in the vegetables from the sampling sites ($P > 0.05$). If the practice of treating the soils in the irrigation gardens with contaminated waters is not controlled, it may lead to health hazard on the part of consumers of the vegetables on the long term. Therefore, there is the need to continually monitor, control and take necessary policy decisions so as to limit and ultimately prevent these avoidable problems.

Keywords: Irrigation, Vegetables, Heavy metals, Atomic Absorption Spectrophotometry (AAS), Gardens

1. Introduction

The term "heavy metals" refers to any metallic element that has a relative density greater than 4g/cm^3 (Grant and Grant, 1987). Heavy metals include; Lead (Pb), Cadmium (Cd), Zinc (Zn), Mercury (Hg), Arsenic (As), Silver (Ag), Chromium (Cr), Copper (Cu), Iron (Fe) and the Platinum group elements. They are non-biodegradable and persistent environmental contaminants which may be deposited on the surface and then absorbed into the tissues of vegetables. They might accumulate in the food chain with risks to the health of animals and humans which are less sensitive to metal toxicity compared to plants but are capable of concentrating heavy metals. The uptake of metals from the soil depends on different factors such as their solubility, soil pH, plant growth stages, fertilizer and soil (Sharma et al., 2006; Ismail et al., 2005). Plant species have various ways of removing and accumulating heavy metals, hence there are reports indicating that some species may accumulate specific heavy metals, causing a serious risk to human health when plant-based food stuff are consumed (Wenzel and Jackwer, 1999). Vegetables are edible plants which store up food reserves in their roots, stems, leaves and fruits. They play an important role in maintaining general good health due to the presence of mineral elements such as calcium, iron, sulphur, potassium and vitamins such as vitamins A, B, and C. These substances help to build bone, teeth and protect the body from diseases. They also regulate body processes on which vitality and good health depend. Leafy vegetables are widely used for culinary purposes. They are used to increase the quality of soup and for dietary purposes (Sobukola et al., 2007). They contain cellulose and form roughage which helps the bowel to function regularly in the elimination of unwanted matter from the body. They also contain 70-75% water which is essential to the body system. They are very important protective foods, useful for the maintenance of health, prevention and treatment of various diseases (D'Mello, 2003). Waste water irrigation is known to contribute significantly to the heavy metal contents of soils

(Mapanda et al., 2005). Irrigation is the artificial addition of water to soils in order to meet plants' needs to overcome drought limitations and improve the crops' yields. However, other factors such as soil and water quality and management practices are also important.

1.1 Statement of the Problem/Justification

Vegetables constitute important part of the human diet since they contain carbohydrates, proteins, vitamins, minerals as well as trace elements. Accumulation of heavy metals by vegetables may depend on plant species as well as concentrations of heavy metal. These heavy metals are not abundant in soil, but they may be accumulated through disposal of sewage water. Disposal of sewage water is a great problem. These sewage effluents are considered not only a rich source of organic matter and other nutrients but also they elevate the level of heavy metals like Cu, Pb, Cr and Cd in receiving vegetable of soils (Singh K.P et al., 2004). Elevated concentration of heavy metal can negatively affect human being. Heavy metals are not easily biodegradable and consequently can be accumulated in human vital organs leading to unwanted side effect (Sathawara N.G et al., 2004). This situation causes varying degrees of illnesses based on acute and chronic exposures. Wastewater irrigation is known to contribute significantly to the heavy metal contents of soils (Mapanda et al., 2005; Devkota and Schmidt, 2000). A vast majority of the vegetables consumed in Katsina state are produced through irrigation, and the water used by the farmers is mostly waste water from gutters and other waste water outlets. There is therefore the need for periodic monitoring of these vegetables for the presence of heavy metals. Results obtained from this study will reveal if the vegetables consumed by the inhabitants of Katsina state are contaminated or not when compared with world health organization safe limits and standards.

2. Methodology

2.1 Description of Study Area

1. Katsina is a city in Nigeria. Its area is 52 sq mi 142 km². The city's population is 318,459. Katsina is located at 12.5139° N, 7.6114° E.
2. Daura is a town and local Government area in Katsina state, Northern Nigeria, with a population of 178,277. Daura is located at Latitude: 13° 01' 58.76" N and Longitude: 8° 19' 24.64" E.
3. Funtua is a town and local government area in Katsina state, Northern Nigeria. It has an area of 448 km² and a population of 225,571 (at the 2006 census) and 420,110 according to 2012 estimate. Funtua has a conducive weather condition as it lies on the latitude and longitude 11° 32' N and 7° 19' E respectively. The city has an average temperature of 32°C and humidity of 44%.

2.2 Sample Collection and Analysis

The vegetables analyzed include spinach, cucumber, cabbage, lettuce and bitter leaf. The vegetables were washed with distilled water to remove dust particles and then cut to separate the root, stems and leaves using a knife. The leaves and soil were air-dried and then oven-dried at 65°C. Dried vegetable samples were ground into a fine powder using a commercial blender (TSK WestPoint, France) and stored in polyethylene bags until they were used for digestion. 2g of each vegetable was weighed into a digestion flask and treated with 9ml of an acid mixture made up of concentrated nitric acid (HNO₃), hydrochloric acid (HCl) and sulphuric acid (H₂SO₄). A blank sample was prepared by applying 9ml of concentrated HNO₃, HCl and H₂SO₄ into an empty digestion flask. The samples were mixed and heated for 30 minutes on an electric hot plate at 80-90°C at which they were brought to boil and a clean solution was obtained. After cooling, the solution was filtered with whatman No. 4 filter paper and then transferred quantitatively to a 100ml volumetric flask by adding 50ml of de-ionized water. The solution was then preserved in a universal bottle for further analysis. All reagents used were of analytical grade and the Atomic Absorption Spectrophotometer (AAS, Perkin Elmer model 2130) was used to determine the heavy metals {Copper (Cu), Cobalt (Co) Zinc (Zn), Lead (Pb) Cadmium (Cd), Chromium (Cr), and Manganese (Mn)} in the digested solution.

2.3 Data Management and Analysis

Data were reported as mean ± standard deviation (SD). One way analysis of variance (ANOVA) was used to determine significant difference (p<0.05) between groups using SPSS version 19.0.

3. Results and Discussion

Vegetables are known to accumulate heavy metals either from waste water or dump site (Arora et al., 2008). The concentrations of Cd, Co, Cu, Cr, Fe, Mn, Pb and Zn from the three sampling sites are shown in Table 1 and their means in Figure 1.

3.1 Heavy Metals Concentration Across all the Vegetables

Cadmium is a non-essential element in foods and natural waters and it accumulates principally in the kidneys and liver (Divrikli et al. 2006.). Various sources of environmental contamination have been implicated for its presence in foods. Differing values have been previously reported in leafy vegetables which include 0.090mg/kg for fluted pumpkin by Sobukola et al. (2010) and 0.049mg/kg by Muhammad et al. (2008) for lettuce. In this study, Cadmium was detected at low concentrations of 0.0102 – 0.0147mg/kg in vegetables purchased from Funtua in spinach and Katsina in Lettuce. This is below the FAO/WHO safe limit of 0.1mg/kg for cadmium consumption in vegetables. The order of increasing of concentration of Cd is as follows: Katsina: lettuce<cabbage<cucumber<spinach<bitterleaf.Funtua:cucumber<lettuce<cabbage<bitterleaf<spinach. Daura: lettuce<cucumber<cabbage<spinach<bitterleaf. Cobalt shows highest concentration of 0.224 mg/kg in spinach from Katsina and lowest concentration of 0.0612 mg/kg in cabbage from Katsina. With the exception of spinach from Katsina, all the samples are above the FAO/WHO safe limits of 0.1 mg/kg.

The order of increasing of concentration of Co is as follows: Katsina: cabbage<cucumber<bitterleaf<lettuce<spinach.Funtua:lettuce<cucumber<bitterleaf<cabbage<spinach. Daura: lettuce<cabbage<cucumber<bitterleaf<spinach. The concentration of copper ranged from 0.014 mg/kg in spinach from Funtua to 0.9951 mg/kg in lettuce from Katsina. All the results obtained are below the FAO/WHO safe limits of 40 mg/kg. This is higher than the findings of Divrikli et al. (2006) and Ozcan (2004) who reported Copper concentrations of 0.02mg/kg and 0.0081mg/kg, respectively for Indian Basil. The order of increasing of concentration of Cu is as follows: Katsina: spinach<bitterleaf<cabbage<cucumber<lettuce.Funtua:spinach<lettuce<cucumber<bitterleaf<cabbage.Funtua:spinach<bitterleaf<cabbage<cucumber<lettuce.

The concentration range for Cr was found to be 0.013 to 0.3406 mg/Kg with the highest concentration recorded in Funtua spinach and the lowest in the spinach from Katsina. All the results obtained are below WHO/FAO safe limits. The order of increasing of concentration of Cr is as follows: Katsina: spinach<cucumber<lettuce<cabbage<bitterleaf.Funtua:spinach<cucumber<lettuce<cabbage<bitterleaf. Daura: spinach<cucumber<lettuce<cabbage<bitterleaf. Interestingly all the vegetables show similar trend in the accumulation Cr across all the sampling locations. Fe concentration ranged from 0.0116 mg/kg from Katsina in spinach to 0.1247 mg/kg in the cabbage from Katsina. All the vegetables are considered safe, as the Fe concentration of below the WHO/FAO safe limits of 150 mg/kg. The order of increasing of concentration of Cr is as follows: Katsina: bitterleaf<spinach<cabbage<lettuce<cucumber.Funtua:bitterleaf<spinach<cabbage<lettuce<cucumber.Daura:Spinach<lettuce<bitter leaf<cabbage<cucumber. Maximum concentration of Mn of 0.037 mg/kg was found in cabbage from Katsina and the lowest concentration of 0.0013 mg/kg was found in lettuce from Funtua. All the results obtained are below WHO/FAO safe limits. The order of increasing of concentration of Mn is as follows: Katsina: cabbage<spinach<bitterleaf<lettuce<cucumber. Funtua: lettuce<cucumber<spinach<bitterleaf. Daura: lettuce<spinach<bitter leaf<cucumber<cabbage. Lead, a nonessential toxic metal showed highest concentration of 0.3481 mg/kg in cabbage sampled from Katsina and lowest concentration of 0.0031 mg/kg bitter leaf from Katsina. The concentration of lead is above the WHO/FAO safe limits of 0.3 mg/kg in cabbage and cucumber sampled from Katsina. There are other concentrations that are also close to the WHO/FAO safe limits in most of the vegetables sampled. The elevated levels of Pb in certain leafy vegetables may also occur due to contaminants in irrigation water, soil or industrial and vehicular emission as lead occurs in the fuel as anti-knocking (Mebale A.A et al., 2014).The traffic is so voluminous around the study sites that the air pollution could convert to soil pollution in short term. The introduction of lead into the food chain may affect human health and may cause disruption of the biosynthesis of hemoglobin and anemia, rise in blood pressure, kidney damage, miscarriages and subtle abortions, disruption of nervous systems and brain damage. Thus studies concerning lead accumulation in vegetables have increased important. The order of increasing of concentration of Pb is as follows: Katsina: bitter leaf<lettuce<spinach<cucumber<cabbage. Funtua: bitter leaf<cucumber<spinach<cabbage<lettuce. Daura: bitter leaf<spinach<lettuce<cucumber<cabbage. Finally the results indicated the mean concentration range of Zn to be 0.0065 to 0.0003 mg/Kg with the highest concentration found in Funtua spinach onions and the lowest in bitter leaf from Funtua. However, the highest value obtained is still below the WHO/FAO safe limit of Zn (50 mg/Kg) in fresh vegetables. The results obtained in this study are comparable with some literature values of similar studies reported previously (Onianwa et al., 2001; Erwin and Ivo, 1992; Pennington et al., 1995). The order of increasing of concentration of Pb is as follows: cabbage<lettuce<bitter leaf<cucumber<spinach. Funtua: bitter leaf<cabbage<cucumber<lettuce<spinach. Daura: bitter leaf<cabbage<cucumber<lettuce<spinach.




Table 1: Result of Heavy Metals Concentrations of Vegetables from Katsina, Funtua and Daura

Sampling Sites	Metals (mg/kg)							
	Cd	Co	Cu	Cr	Fe	Mn	Pb	Zn
Bitter leaf								
Katsina	0.0119	0.1656	0.0037	0.1347	0.1128	0.0144	0.0031	0.0028
Funtua	0.0147	0.2053	0.0018	0.3406	0.0205	0.0223	0.1178	0.0003
Daura	0.0133	0.18545	0.00275	0.23765	0.06665	0.01835	0.06045	0.00155
Mean	0.18545	0.00275	0.23765	0.06665	0.01835	0.06045	0.00155	0.00155
STD	0.016207	0.000776	0.084058	0.037681	0.003225	0.046826	0.001021	0.001021
Cabbage								
Katsina	0.0108	0.0612	0.0044	0.1327	0.1247	0.0048	0.3481	0.0004
Funtua	0.0129	0.2097	0.0019	0.0726	0.0256	0.037	0.2439	0.0021
Daura	0.01185	0.13545	0.00315	0.10265	0.07515	0.0209	0.296	0.00125
Mean	0.13545	0.00315	0.10265	0.07515	0.0209	0.296	0.00125	0.00125
STD	0.060625	0.001021	0.024536	0.040457	0.013146	0.042539	0.000694	0.000694
Cucumber								
Katsina	0.0113	0.1524	0.0086	0.0301	0.094	0.0314	0.2522	0.0034
Funtua	0.0122	0.1964	0.0015	0.0431	0.0496	0.0061	0.1449	0.0034
Daura	0.01175	0.1744	0.00505	0.0366	0.0718	0.01875	0.19855	0.0034
Mean	0.1744	0.00505	0.0366	0.0718	0.01875	0.19855	0.0034	0.0034
STD	0.017963	0.002899	0.005307	0.018126	0.010329	0.043805	0	0
Lettuce								
Katsina	0.0102	0.208	0.9951	0.0732	0.053	0.0218	0.1011	0.0006
Funtua	0.0127	0.1386	0.0014	0.0484	0.0496	0.0013	0.2501	0.0054

Daura	0.01145	0.1733	0.49825	0.0608	0.0513	0.01155	0.1756	0.003
Mean	0.1733	0.49825	0.0608	0.0513	0.01155	0.1756	0.003	0.003
STD	0.028332	0.405676	0.010125	0.001388	0.008369	0.060829	0.00196	0.00196
Spinach								
Katsina	0.0116	0.224	0.0015	0.0013	0.0116	0.0116	0.1115	0.0062
Funtua	0.0147	0.2179	0.0014	0.0052	0.0256	0.0124	0.1563	0.0065
Daura	0.01315	0.22095	0.00145	0.00325	0.0186	0.012	0.1339	0.00635
Mean	0.22095	0.00145	0.00325	0.0186	0.012	0.1339	0.00635	0.00635
STD	0.00249	4.08E-05	0.001592	0.005715	0.000327	0.01829	0.000122	0.000122
WHO/FAO STDS	0.1	0.1	40	1.3	150	6.61	0.3	60

Statistical Analysis

The P values for the heavy metals were shown in Table 2. There is a significant differences in the concentration of Cd in the vegetables from the sampling sites ($P < 0.05$), while there is no significant difference in the concentration of Cr, Co, Cu, Fe, Mn, Pb, Zn in the vegetables from the sampling sites. This could be attributed to the similarities of the activities and topography of the sampling sites. This made the null hypothesis made from this study acceptable.

Table 2: P-values for the Heavy Metals Sampled in Vegetables from Katsina, Funtua and Daura Towns

Metals	Cd	Co	Cr	Cu	Fe	Mn	Pb	Zn
P value	0.0075	0.5573	0.5553	0.9107	0.1141	0.9889	0.9538	0.8424

4. Conclusion

This study further confirms the increased danger of growing vegetables on soils irrigated with contaminated industrial and domestic wastewaters. However, the levels of the metals in most of the vegetables are currently within the WHO/ safe limits guidelines. But lead is found in all the vegetables with some concentrations such as in cabbage and cucumber. If the practice of treating the soils in the irrigation gardens with contaminated waters is not controlled, it may lead to health hazard on the part of consumers of the vegetables on the long term. Therefore, there is the need to continually monitor, control and take necessary policy decisions so as to limit and ultimately prevent these avoidable problems. However, in the meantime, farmers from the study areas are hereby encouraged to use well or borehole water for irrigation in their gardens instead of contaminated streams.

References

- Devkota B, Schmidt GH (2000). Accumulation of heavy metals in food plants and grasshoppers from the Taigetos Mountains, Greece. *Agric. Ecosyst. Environ.*, 78: 85-91
- D'Mello JPF (2003). Food safety: Contamination and Toxins. CABI Publishing, Wallingford, Oxon, UK, Cambridge, M.A. p. 480.
- Divrikli U, Horzum N, Soylak M, Elci L (2006). Trace heavy metal contents of some spices and herbal plants from western Anatolia, Turkey. *Int. J. Food Sci. Technol.* 41: 712-716.
- Erwin JM, Ivo N (1992). Determination of Lead in tissues: A pitfall due to wet digestion procedures in the presence of sulphuric acid. *Analyst.* 17: 23-26.
- Grant R, Grant C (1987). Grant and Hackh's Chemical Dictionary, McGraw-Hill, New York.
- Ismail BS, Fariyah K, Khairiah J (2005). Bioaccumulation of heavy metals in vegetables from selected agricultural areas. *B. Environ. Contam. Tox.* 74:320-327..
- Mebale A.A., Ndong R.O., Affane A.L.N., Omanda H.M., Nziengui P.P., Biyogo R.M., Ondo J.A. (2014) "Assessment of Metal Content in Leafy Vegetables Sold in Markets of Libreville, Gabon". *Int J Cur Res Rev*, 06 (01). 28-33.
- Mapanda F, Mangwayana EN, Nyamangara J, Giller KE (2005). The effect of long term irrigation using wastewater on heavy metal contents of soils under vegetables in Harare, Zimbabwe. *Agric. Ecosys. Environ.*

107:151-165.

Naveh Z, Steinberg EH, Chaim S (1979). The use of bio-indicators for monitoring of air pollution by fluorine, ozone and sulphur dioxide: In Environmental Bio-monitoring, Assessment, Prediction and Management of certain case studies and related Quantitative issues. Ed. Cains, G.I.P. and Waters, W.E. International Cooperative Publishing House, Fatiland, USA, pp. 21-47

Onianwa PC, Adaeyemo AO, Odowu EO, Ogabiela EE (2001). Copper and Zinc contents of Nigerian foods and estimates of the adult dietary intakes. *J. Chem.*, 72: 89-95.

Ozcan M (2004). Mineral contents of some plants used as condiments In Turkey. *Food Chem.* 84:437-440.

Pennington JAT, Schoen SA, Salmon GD, Young B, John RD, Matrt RW (1995). Composition of core foods of the USA foods of the USA food supply 1982-1991. *J. Food Compos. Anal.*, 8(2): 129-169

Sharma RK, Agrawal MM (2006). Heavy metals contamination in vegetables grown in waste water irrigated areas of Varanasi, India. *Ecotoxicology and Environmental Safety* 66: 258-266.

Sathawara N.G., Parikh D.J., Agarwal Y.K., Bull (2004) *Environ. Contam. Toxicol.*, 73, 264-269.

Sobukola OP, Adeniran OM, Odedairo AA, Kajihansa OE (2010). Heavy metal levels of some fruits and leafy vegetables from selected markets in Lagos, Nigeria. *Afr. J. Food Sci.* 4(2): 389 - 393

Wenzel W, Jackwer F (1999). Accumulation of heavy metals in plants grown on mineralized solids of the Austrian Alps. *Environ. Poll.* 104: 145-155.