

Evaluation of Arsenic Concentration in Rocks of Kaltungo Area, Upper Benue Trough, Nigeria

M.N Mboringong¹, A.Olasehinde^{1*}, R.P.Tabale¹ A. Yusuf¹ and E.C Ashano²

1. Gombe State University, P.M.B 127, Gombe, Nigeria.

2. University of Jos, P.M.B 2084, Jos, Nigeria.

* yinka516@gmail.com

1. Abstract

This research aimed at investigating the concentration of arsenic (As) due to their potential environmental impact and contamination, in the rocks (Coarse Porphyritic Granite, Biotite Granite, Bima Sandstone and basalt) of the Kaltungo area which is part of the Upper Benue Trough. Eight (8) fresh samples (two from each lithologic unit) were collected and analysed using Inductively Coupled Plasma Optical Emission Spectrometer (ICPOES) for arsenic (As). The results were presented using graphical presentations and this revealed that rocks in the study area host high concentration of arsenic (As) (87.54-237.65 ppm) compared with the average crustal abundance of 2ppm. These values can be attributed to high concentration of the element in the initial magma and/or hydrothermal and weathering processes which acted on the rocks. The high concentration of arsenic (As) in the rocks of Kaltungo area calls for further studies on its distribution and concentration in the soils, stream sediments and ground water.

Keywords: Arsenic, Kaltungo, Upper Benue trough, environment.

2. Introduction

There is a considerable interest in the study of arsenic (As) distribution within the kaltungo area due to their potential environmental impact and contamination. In the study area a lot is known about the tectonic evolution and origin but little has been done on the natural variability of the toxic elements in the rocks. Reasons for this include the interest in the exploration for oil which include sedimentary, stratigraphy and paleontological studies (Allix and Poppof, 1983; Benkhelil, 1982; Benkhelil and Robineau, 1983; Benkhelil, 1986; Benkhelil et al., 1988; Enu, 1980; Guiraud, 1990; Guiraud, 1993; Lawal and Moullade, 1986; Maurin *et al.*, 1986; Zaborski, 1990; 1993; Zaborski *et al.*, 1997) and limited value for chemical classification or petrogenitic studies.

The present work tend to study the distribution of arsenic (As) in the rocks of Kaltungo to evaluate its concentration due to their potential for environmental contamination, which will in turn serve as baseline for further studies in its distributions in the soil and ground water respectively.

3. The geology

The Benue Trough of Nigeria is an intracontinental Cretaceous basin about 1000kilometres in length elongated in the NE-SW directions which overlies the Precambrian shield of the West African mobile belt (Maurin *et al.*, 1986; Guiraud, 1990) . Reviews and the origin and the evolution of this linear NE-SW trending megastructure were provided by Benkhelil ,1989; Freeth, 1990; Guiraud and Maurin, 1991; 1992, Genik, 1992, 1993. The sedimentary sequence infilling the Upper Benue Trough includes continental and marine deposits ranging from Upper Aptian to Paleocene in age. The Bima Formation were laid unconformably on the basement in the fitted Wrench – fault basin. The study area consists partly of the basement complex and cretaceous sediments in the Upper Benue Trough (Figure 1). This transgressive episode caused the deposition of Yolde, Dukul, Jessu, Sekule and Numanha sedimentary Formations, all of which outcrop as inliers to Bima Formation in Dadiya syncline (Figure 2). These Formations are shallow marine depositions of limestone, shale and mudstone and they are foundmainly as narrow strips of rocks in the study area, such as Banju, Guyuk, and Shellen. The Longuda Basalt of tertiary age terminates the lithological succession in the area. The Coarse Porphyritic Granite, Biotite Granite, Bima Sandstone and Basalts represent the rocks of the Kaltungo area (Figure 3).

4. Chemistry of Arsenic

Arsenic is situated in the 33rd spot on the periodic table, right next to Germanium and Selenium. It has atomic weight of 74.92, with Electron Configuration [Ar] 4s23d104p3, Density 5.727 g/cm³, Oxidation States 5, 3, 2, 1,-3 and Rhombohedral Crystal Structure. It is a chalcophile element being usually found in sulphide minerals. They exhibit similar or identical atomic radii (As, 1.5 Å; Sb, 1.6 Å and Bi 1.7Å), electronegativity (2, 1.9, 1.9) but dissimilar ionic radii (0.47,0.62. 0.77 Å)(Heydemann, 1969).

5. Methodology

Eight (8) Fresh surface rock samples were collected from outcrops of all lithologic units (two per each lithologic unit) in the study area (Figure 3) which includes the Coarse Porphyritic Granite, Biotite Granite, Bima Sandstone and Basalts. Arsenic (As) was determined using the Inductively Coupled Plasma optical Emission Spectrophotometer, (ICPOES), after HF / HClO₄ dissolution at the PTDF Geochemistry Laboratory of Department of Geology and Mining, University of Jos. Analytical precision for the instrument was 0.5 – 1% and an accuracy of 2% - 5%. Quality control procedure was by the simultaneous analysis of international standards and blanks and calculated after the method of Ashano (1996).

6. Result and Discussion

The result of the analysis is presented table 1. The concentration Arsenic in all the rock types shows much higher concentration than the Average Crustal Abundant (ACA) of 2ppm. The concentration varies from 87.54ppm in K3 to 235.2 ppm in K1 with mean concentration 194.05ppm in K1, 237.65ppm in K2, 158.120ppm in K3 and 163.10 ppm respectively.

K1 = Coarse Porphyritic Granite; K2= Biotite Granite; K3= Bima Sandstone; K4= Basalts

CA= Crustal Abundant

Arsenic may be lost from soils by two sources, either by streams, which happens rapidly but only low levels of arsenic accumulate, or by groundwater, which is very slow, however comparatively high levels of arsenic may build up. Arsenic is released into ground and surface waters by the erosion, dissolution and weathering of rocks which is highly concentrated in the rocks of the study area. Arsenic can stick to mineral surfaces from circulating groundwater.

7. Human health effects

According to a 1999 study by the National Academy of Sciences, arsenic in drinking water causes bladder, lung and skin cancer, and may cause kidney and liver cancer. The study also found that arsenic harms the central and peripheral nervous systems, as well as heart and blood vessels, and causes serious skin problems. It also may cause birth defects and reproductive problems. According to this same study, an arsenic concentration of 50 ppb, gives you a 1/100 lifetime chance of developing cancer. Even an arsenic concentration as low as 1 ppb gives you a 1/1000 lifetime chance of developing cancer, which is 10 times higher than the EPA standard. Therefore it is necessary that people have access to water which complies with the WHO standard for arsenic of <10 ppb and the arsenic concentration should be kept as low as possible.

8. Conclusion

we have reported the anomalous concentration of arsenic in the rocks of kaltungo area which varies from 87.54-237.65ppm. This work would serve as a baseline for further studies of the concentration of this element in soil, stream sediments and ground water of the area.

References

- Allix, P. and Poppof, M (1983): Le Cretace inferieur de la partie nord-orientale du fosse de la Benoue (Nigeria). Un exemple de la relation etroite entre tectonique et sedimentation. *Bulletin des Centres de Recherches Exploration –production Elf- Aquitaine* 7, 349-359
- Ashano, E. C., (1996): Sampling Variability in Exploration Geochemistry: An Estimation Model. *Jour. Min and Geol.* 32, 27 – 33.
- Benkhelil, J., (1982): Benue Trough and Benue Chain. *Geological Magazine* 119, 155-168 Benkhelil J, Robineau B (1983) Le fosse de la Benoue est il um rift? *Centre Rech Eplor Prod Elf Aquitaine Bull* 7, 315–321
- Benkhelil, J.,(1986): Structure et Evolution geodynamique du basin intracontinental de la Benoue (Nigeria). *Theses de Doctrat d' Etat, Universite de Nice*, 226p
- Benkhelil, J., Dainelli, P., Ponsardd, J.F., Popoff, M., Saugy, L.,(1988): The Benue Trough: Wrengh fault related basin on the border of the Equatorial Atlantic. In: Manspezier, W., (Ed), *Triassic- Jurassic rifting-continental break-up and the origin of the Atlantic Ocean and passive margins, Development in geotectonics*, 22, 789-819.
- BenkhelilJ(1989)The origin and evolution of the Cretaceous Benue Trough, Nigeria. *J. Afr Earth Sci* 8, 251–

- Coulon, C., Vida, P., Dupuy, C., Baudin, P., Popoff, M., Maluski, H., and Hermite, D.,(1996). The Mesozoic to early Cenozoic agmatism of the Benue trough (Nigeria); geochemical evidence for the involvement of the St Helena plume. *J. Petrol.*, 37, 1341-1358.
- Enu, E.L. (1980): Sukuliye (Sekule) Formation, facies equivalent of the Numanha Formation in the upper Benue Trough, Nigeria. *J. min. Geol* 17, 91-95
- Freeth, S.J.(1990): The origin of Benue Trough-In: Ofueogbu, C.O(ed): The Benue Trough: Structure and evolution. Vieweg, Braunschweig, 217-228
- Genik, G.J.(1992): Regional framework, structural and petroleum aspect of rift basins in Niger, Chad and the Central African republic (C.A.R). *Tectonophysics* 213, 169-185
- Genik, G.J.(1993): Petroleum geology of Cretaceous-Tertiary rift basins in Niger, Chad, and Central Africa Republic. *Bull. Amer.Assoc.Petroleum Geol.* 77,1405-1434
- Guiraud, M.,(1990): Tectono-sedimentary framework of Early Cretaceous Continental Bima Formation(Upper Benue trough NE Nigeria) *Journ. Afri. Earth Sciences(Special publication)* 10, 341-353
- Guiraud, R. and Maurin, J.C,(1991): Le rifting en Afrique au Cretace inferieur: Synthese structurale, mise en evidence de deux etapes dans la genese des basins, relations avec les ouvertures oceanique peri-africaines. *Bull.Soc.geol.France*, 16, 811-823
- Guiraud, R. and Maurin, J.C.,(1992): Early Cretaceous rifts of Western and Central Africa: An overview. *Tectonophysics* 213, 811-823
- Guiraud, M (1993) Late Jurassic – Early Cretaceous rifting and Late Cretaceous transpressional inversion in the Upper Benue Basin, NE Nigeria. *Bull Rech Elf Aquitaine* 17, 371–383.
- Heydemann, A. 1969. In: Wedepohl, K.H. (ed.) *Handbook of geochemistry*. 1. Springer-Verlag, Berlin, 376-412.
- Hughes, M. F. (2002). Arsenic toxicity and potential mechanisms of action. *Toxicol. Lett.* Vol 135, 1–16.
- Maurin, J.C., Beenkhelil, J., and Robineau, B.(1986): Fault Rocks of the Kaltungo Area Lineaments N.E Nigeria, and their relationship with the Benue Trough Tectonics. *Journ. Geol. Soc. London.* 143,587-599
- Samaila, N.K, Dike, E.F.C., Obaje, N.G.,(2007): Microstructure in the Cretaceous Bima Sandstone, Upper Benue Trough, N.E. Nigeria: Implication for hydrocarbon migration. *Journ. Of Afri. Earth Sci.* 50, p6-15
- Zaborsky, P.M., Ugodunluwa, F., Idornigie, A., Nnabo, P., Ibe, K.,(1997): Stratigraphy and structure of the Cretaceous Gongola Basin(N.E Nigeria). *Bulletin des centres de Recherches Exploration Production Elf-Aquitaine* 21 (1), 153-177.

Table 1. Concentration of arsenic (As) within the Rocks of Kaltungo

| Samples (ppm) | Coarse Porphyritic Granite (K1) | Biotite Granite (K2) | Bima Sandstone (K3) | Basalts (K4) |
|---------------|---------------------------------|----------------------|---------------------|--------------|
| 1 | 152.900 | 232.200 | 228.700 | 174.600 |
| 2 | 235.200 | 243.100 | 087.540 | 151.600 |
| Mean | 194.0500 | 237.650 | 158.120 | 163.100 |
| Median | 194.0500 | 237.650 | 158.120 | 163.100 |

K1 = Coarse Porphyritic Granite; K2 = Biotite Granite; K3= Bima Sandstone; K4= Basalts

Table 2. Summary of health effect of arsenic (Adopted from Hughes 2002).

| System | Effect |
|----------------|---|
| Skin | Skin lesions—keratosis, hyperkeratosis |
| Cardiovascular | Blackfoot disease |
| Nervous | Peripheral neuropathy, encephalopathy |
| Hepatic | Cirrhosis, hepatomegaly |
| Hematological | Bone marrow depression |
| Endocrine | Diabetes |
| Renal | Proximal tubule degeneration, papillary and cortical necrosis |




Figure 1. The Benue trough (after Coulon et al 1996)

| AGE | PALAEO-ENVIRONMENT | GONGOLA BASIN | YOLA BASIN | LAMURDE-LAU BASIN |
|----------------|--------------------------|-------------------|----------------------------------|-------------------|
| Quaternary | Continental | Biu Basalts | Longuda Basalts | |
| Pliocene | | | | |
| Miocene | | | | |
| Oligocene | | | | |
| Eocene | | | | |
| Paleocene | | Kerri-Kerri Fm | | |
| Maastrichtian | Continental/Transitional | Gombe Sandstone | | |
| Campanian | Marine | Pindiga Formation | Lamja Sandstone | Lamja Sandstone |
| Santonian | | | | |
| Coniacian | | | Numanha Fm | Numanha Fm |
| Turonian | | | Sekuleye Fm | Sekuleye Fm |
| Cenomanian | | | Jessu Fm | Jessu Fm |
| | | | Dukul Fm | Dukul Fm |
| | | | Yolde Formation | |
| Upper Albian | Continental | | Bima Sandstone (B ₃) | |
| Late Aptian | | | Bima Sandstone (B ₂) | |
| Early Aptian | | | Bima Sandstone (B ₁) | |
| Late Jurassic? | | | | |
| Pre-Cambrian | | | Basement Complex | |

Figure 2. Stratigraphic succession of the Upper Benue Trough (after Samaila et al, 2007)




Figure 3. The Geological map of the study area showing sampling location




Figure 4. Histogram plots of Arsenic distribution in the rocks of katungo K1, K2, K3, and K4 in ppm




Figure 5. Line plots of the Arsenic concentration in the rock samples of Kaltungo compared with the Crustal Abundant of 2.0 ppm

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

