

# Review on Effect of Soil Acidity on Barley (*Hordeum vulgare* L.) Producing Central Highlands of Ethiopia

Tigist Shiferaw and Mesfin Tadele  
Holetta Agricultural Research Center, P.O. Box 31, Holetta, Ethiopia

## Abstract

Barley is an important food crop grown in diverse agro-ecology, different seasons and production systems of Ethiopia. Soil acidity is now a serious threat to barley production in most high lands of Ethiopia and about 43% of the Ethiopian high land with altitude of >1500 arable land are affected by soil acidity. Acidic soils affected areas of Ethiopia found at western and southwestern, northwestern part of the country. Barley is considered to be more sensitive to acidic soils. The initial toxic effects of acid soil are stunted and shortened root growth then delay barley germination and initiation. This continued by delay of days to heading and days to maturity, decrease of plant height and increase severity of foliar diseases. Finally the overall effect of soil acidity is significantly expressed on biomass and grain yield and yield components barley.

**Keywords:** Barley, Soil acidity stress

**DOI:** 10.7176/JNSR/9-1-03

**Publication date:** January 31<sup>st</sup> 2022

## 1. INTRODUCTION

Barley (*Hordeum vulgare* L.) is the fourth most important cereal crop grown in the world after maize, wheat and rice with productivity of 3.3 ton ha<sup>-1</sup>. On the African continent, Ethiopia, Algeria, Morocco, South Africa and Tunisia are the top five largest barley producers (FAO, 2020). In Ethiopia barley is the most important cereal crop following teff, maize, wheat and sorghum in bulk production and area coverage and fifth in yield (ton ha<sup>-1</sup>) after maize, rice, wheat and sorghum with around 0.92 million ha and 2.3 million tons. In addition, the productivity of barley is 2.53 ton ha<sup>-1</sup> (CSA, 2021).

Barley is an important grain food crop grown in diverse agro-ecology from 1,500 to 3,500m altitude for many purpose in different seasons and production systems and a common food grain, especially for highlands of Ethiopia (Berhane *et al.*, 1996; Muluken, 2013).

The abiotic stresses were indicated among the most important constraints in barley production (Bayeh and Berhane, 2011). From abiotic stress, soil acidity is one of the major constraints for barley production. The acidic soils mainly Nitosols or Oxisols are common in Ethiopian highlands, where the rainfall intensity is high and crop cultivation has been carried out for centuries (Desta, 1987). Acid soils cause nutritional disorders, deficiencies or unavailability of essential nutrients such as calcium, magnesium, molybdenum and phosphorus and toxicity of aluminium, manganese and hydrogen ions activity in the soil (Foy *et al.*, 1978). Toxic elements like aluminium and manganese are the major causes for crop failure in acid soils. These elements are a problem in acid soils because they are more soluble at pH below 5 (Birhanu, 2008). Barley grows well on neutral to mildly acid soils with a pH range of 6–7. Increasing soil acidity may lead to poor plant vigor and crop growth, stunted root growth, persistence of acid-tolerant weeds, increased incidence of diseases, abnormal leaf color and reduced yield (Somani, 1996). Yields of the major cereal crops, particularly barley is reduced by 0.5 ton ha<sup>-1</sup> as a result of soil acidity (Paoulos, 2001). Among small grain crops barley is one of the most sensitive species to acid soil among small-grain crops, but differences in acid soil tolerance exist among the genotypes (Zhao *et al.*, 2003).

Therefore, this research was conducted to assess the effect soil acidity on barley producing areas of Ethiopia.

## Soil Acidity Extent and Distribution in Ethiopia

Soil acidity is one form of chemical degradation of soils. The problems of acid soils are high in acidity and low amount of exchangeable cations especially calcium and it is considered to be one of the most important factors that affect the soil chemical fertility. Acid soils are phototoxic because of nutritional disorders, deficiencies or unavailability of essential nutrients such as calcium, magnesium, molybdenum and phosphorus and toxicity of aluminium, manganese and hydrogen activity (Foy *et al.*, 1978).

The main factors giving rise to increased soil acidity in Ethiopia include climatic factors such as rainfall and leaching, temperature, severe soil erosion, anthropological factors, acidic parent material, organic matter decay, removal of products from the farm or paddock and inappropriate use of nitrogenous fertilizers. Soil acidity is more easily understood when there is an abundance of acidic cations, like hydrogen (H<sup>+</sup>) and aluminium (Al<sup>+++</sup>) compared to the alkaline cations like calcium (Ca<sup>++</sup>), magnesium (Mg<sup>++</sup>), potassium (K<sup>+</sup>) and sodium (Na<sup>+</sup>) (Johnson, 1914; Temesgen *et al.*, 2011).

In Ethiopia, about 43% of the Ethiopian high land with altitude of >1500 arable land are affected by soil acidity (Ethiosis, 2014). The most strongly acidic soils are found in western and southwestern parts of Ethiopia,

the central highlands and high rainfall areas of northwestern part of the country. Nevertheless, moderately acidic soils (pH 5.5-6.5) are distributed through much of the rest of the country (Taye, 2008). In moving from central (West Shewa) to Western Ethiopia (West Wellega), the degree of soil acidification is increased ( $ASP > 60$ ). In Western and eastern Wellega zones, the largest proportion of exchangeable acidity was due to exchangeable  $Al^{+++}$  while in West Shewa zone it was due to Exchangeable  $H^+$ . The acidity problem in East and West Wellega zone of Oromia region is critical (Abdennaet *et al.*, 2007).




Figure 1. Distribution of acid soils in Ethiopia (Schlede, 1989, Behailu Kasahun, 2015)

The optimum soil pH for plant production is (pH 6-6.5), at this pH, soil microorganisms are most active and plant nutrients are readily available. Beyond to this delicate balance is disturbed and plant nutrients becomes deficient to plant growth. Some essential nutrients such as calcium, phosphorous, magnesium and molybdenum become unavailable if the soil pH becomes too acid (Eshetu, 2011).

Crop production on acid soils can be sustained by application of lime. The most important benefit of liming acid soils is reduced in solubility of the potentially toxic elements hydrogen, aluminium ( $Al^{3+}$ ) and manganese ( $Mn^{2+}$ ). Optimum nutrient uptake by most crops occurs at a soil pH near 7.0, and the availability of fertilizer nutrients such as nitrogen, phosphorous and potassium generally is reduced as soil pH decreases (Mark *et al.*, 2001). Thus, developing cultivars with improved tolerance to acid soil stress is a solution to address this problem (Scott and Fisher, 1989).

#### 2.4. Effect of Soil Acidity on Growth, Development and Yield of Barley

The most important abiotic stress constraining the production of barley includes low soil fertility, low soil pH, poor soil drainage, frost and drought. From these abiotic stresses, soil acidity is one of the most important constraints in barley production, mainly on Nitisols or Oxisols, of the Ethiopian highlands (Taye and Höfner, 1993).

Barley is considered to be more sensitive to acidic soils than rye, oat, rice and wheat (Bona *et al.*, 1993). The acid tolerance order reported as maize > rye > triticale > wheat > barley (Polle and Konzak, 1985). The initial toxic effects of acid soil are stunted and shortened root growth then limited growth and productivity of barley by restricting water uptake and nutrient absorption (Wang *et al.*, 2006). Because of inhibitory effect of soil acidity on root development, it decreases tolerance of plants to acidity and decreases use of subsoil nutrients and suppression of photosynthetic capacity of shoots. This is associated with cellular and ultra structural modifications in leaves, reduced stomatal opening and  $CO_2$  assimilation, reduced chlorophyll concentration, chlorosis and leaf necrosis (Miyasaka *et al.*, 2007).

Soil acidity cause delay in barley germination and initiation. The principal factor responsible for delay emergence in acidic soil did not delay radical initiation, but delay initiation of hypocotyls elongation and these elongation of hypocotyls was highly associated with rate of tap root growth (Ritchey and Carter, 1993). Soil acidity had a reduction effect on agro-morphological traits of the studied barley genotypes compared to limed soil such as, delay of days to emergence, days to heading and days to maturity, decrease of plant height and increase severity of foliar diseases (Tigist and Thomas, 2021)

Tiller numbers may also be reduced by poor nutrient availability (Baque *et al.*, 2006). Mean value of number of fertile tillers per plant at both limed and not limed soil is statically different and the limed have a higher mean value of fertile tiller (Temesgen, 2014). The number of tillers plant<sup>-1</sup> significantly ( $P < 0.05$ ) differed between cultivars as well as lime treatments. The number of tillers plant<sup>-1</sup> were higher for plants grown in lime treated soil than in lime untreated soil (Tenaye and Tesfaye, 2014). Moreover, barley grown under acidic soil condition is remarkably shorter than lime treated soil. Plant height increased under lime treated soil; this is related to the increase in soil fertility and reduction of the toxic concentration of acidic cations (Chimdi *et al.*, 2012).

Gallardo *et al.*, (1999) reported that, the overall effect of soil acidity is significantly expressed on biomass and grain yield of crops. On barley 50% and 30% reduction of grain yield, respectively, for sensitive and tolerant cultivars when they were grown in naturally acidic soil (pH 4.9) with a large amount of extractable Al compared to that grown in non acidic soil (pH 5.8). The pH of the soil less than 5.5 results in very low yield of barley (0.5 ton ha<sup>-1</sup>) compared with barley growing soils of the country (Taye and Höfner, 1993).

Furthermore, non-amended acidic soil produced the lowest harvest index (HI). The increased percent in HI obtained on acidic soils treated with different lime rate as compared to respective controls (non amended). This is highly likely associated with reduction of concentration of exchangeable acidity and enhancement of exchangeable bases, CEC and available P of the soils (Chimdi *et al.*, 2012). All limed treatments had higher mean values of thousand seed weight (TSW), number of seeds per spike (NSPS) and hectoliter weight (HLW) relative to no lime (Temesgen, 2014). Soil acidity had also a significant effect on yield and yield components of barley genotypes (Table 5). Genotypes grown in unlimed soil had low number of fertile tillers per plant, spike length, numbers of kernels per spike, grain and biomass yields, harvest index, thousand kernel weight and hectoliter weight comparing to genotypes grown in limed soil (Tigist and Thomas, 2021).

#### Genetics Mechanism of Barley (*Hordeum vulgare* L.) Resistance to Soil Acidity

Barley (*Hordeum vulgare* L.) is considered one of the most Al<sup>3+</sup> sensitive cereal crops. Two main mechanisms of resistance have been proposed: exclusion mechanism and resistance mechanism. The exclusion mechanism prevents Al<sup>3+</sup> from entering cells and minimizes Al toxicity, while the resistance mechanism allows plants to take up Al<sup>3+</sup> and accumulate Al<sup>3+</sup> within their cells (Kochian *et al.*, 2004).

The Al<sup>3+</sup> induced secretion of organic acid anions from roots is the best example of exclusion mechanisms in higher plants, such as wheat (*Triticum aestivum*) and sorghum (*Sorghum bicolor*) (Ryan *et al.*, 2009; Tovkach *et al.*, 2013). In wheat, both malate and citrate secretion from roots has been associated with Al<sup>3+</sup> resistance (Ryan *et al.*, 2009), whereas in barley, only citrate exudation from roots has been identified (Furukawa *et al.*, 2007). In addition, aluminium resistance between resistant wheat and barley was significantly different. (Zhou *et al.*, 2013). Thus, a variety of transporters and proteins have been demonstrated to be involved in Al<sup>3+</sup> resistance in plants. Recently, Dai *et al.* (2013) identified 35 Al<sup>3+</sup>-associated proteins from wild barley which were involved in metabolism, cell growth, energy, protein storage, protein biosynthesis, signal transduction, and transporters, etc. There were four proteins specifically expressed in wild barley that were expressed in the Al<sup>3+</sup>.

## CONCLUSION

Soil acidity is now a serious threat to barley production in most high lands of Ethiopia. The extent of acidity is increased in 2.1% within the past three decades, mainly due to continuous cropping system and use of acidifying fertilizers. However, the assessment of genetic variability in response to soil acidity based on the response mechanism among barley genotypes is indispensable to identify tolerant breeding plant materials to develop soil acid tolerant genotypes.

## REFERENCES

- Abdenna Deressa, Negassa Chewaka & Tilahun Geleto. 2007. Inventory of Soil Acidity Status. In Crop Lands of Central and Western Ethiopia. "Utilisation of diversity in land use systems: Sustainable and organic approaches to meet human needs".
- Achalu Chimdi, Heluf Gebrekidan, Kibebew Kibret & Abi Tadesse. 2012. Response of barley to liming of acid soils collected from different land use systems of Western Oromia, Ethiopia. *J. Biodiversity and Environmental Sciences (JBES)*. Vol. 2, No. 5, p. 37-49.
- Baque, M.A., M.A. Karim, A. Hamid & H. Tetsushi. 2006. Effects of fertilizer potassium on growth, yield and nutrient uptake of wheat (*triticum aestivum*) under water stress conditions. *South Pacific Studies*, 27: 25-35.
- Bayeh Mulatu & Berhane Lakew. 2011. Barley research and development in Ethiopia an overview In: Mulatu, B. and Grando, S. (Eds.). *Barley Research and Development in Ethiopia. Proc. the 2nd National Barley Research and Development Review Workshop*. November 28-30, 2006. HARC, Holetta, Ethiopia. pp. 3.
- Behailu Kassahun. 2015. *Soil Fertility Mapping and Fertilizer blending*. ppt.8
- Berhane Lakew, Hailu Gebrea & Fekadu Alemayehu. 1996. Barley production and research. In G. Hailu and J. van Leur (eds.). *Barley Research in Ethiopia. Past work and future prospects*. Proceedings of the First Barley Review work shop, 16- 19 October 1993, Addis Ababa: IAR/ICARDA.
- Birhanu Agumas. 2008. Soil acidity problems and comparison of lime requirement determination methods in different land uses systems: The Case of Fagetalekoma Woreda, Awi Zone. MSc. Thesis, Faculty of Environmental Science, Addis Ababa University, Addis Ababa. pp.19.
- Bona L, Wright RJ, Baligar VC & Matuz J. 1993. Screening wheat and other small grains for acid soil tolerance. *J. Landscape and Urban Planning*; 27:175-178.
- CSA (Central Statistical Agency). 2021. Agricultural sample survey: area and production of major crops, meher

- season. Vol. I. Addis Ababa, Ethiopia.
- Desta Beyene.1987. Effects of liming and nitrogen and phosphorus fertilizers on grain yield of barley. *Ethiopian Journal of Agricultural Science*, 9: 1–13.
- Dai,H.X.,Cao,F.B.,Chen,X.H.,Zhang,M.,Ahmed,I.M.,Chen,Z.H.,etal.2013.Comparative proteomic analysis of aluminium tolerance inTibetan wild and cultivated barleys. *PLoS ONE* 8:e63428. doi:10.1371/journal.pone.0063428
- Eshetu Lemma. 2011.Assessment of soil acidity and determination of Lime requirement for different land use types:The case of degem wereda, north shoa zone. MSc.Thesis Faculty of Environmental Science.Addis Ababa University, Addis Ababa.
- Ethiosis 2014.Soil fertility mapping and fertilizer blending.Agricultural Transformation Agency (ATA) Report, Ethiopia soil information system (Ethiosis).Ministry of Agriculture, Addis Ababa.
- FAO (food and agriculture organization). 2020. Food balance sheets. Faostat. Rome(<http://www.fao.org/faostat/en/#data/QC/visualize>).(Accessed on 12 February 2018).
- Foy, C.D., R.L. Chaney& M.C. White.1978.The physiology of metal toxicity in plants. *Ann. Rev. Plant Physiol.* 29:511-566.
- Furukawa, J.,Yamaji,N.,Wang,H.,Mitani,N.,Murata,Y.,Sato,K.,etal.2007. An aluminum-activated citrate transporter in barley. *Plant CellPhysiol.* 48, 1081–1091. doi:10.1093/pcp/pcm091
- Gallardo F, Borie F, Alvear L&Baer EV.1999. Evaluation of aluminium tolerance of three barley cultivars by two short-term screening methods and field experiments.*J. Soil Sci Plant Nutr*,45:713–719
- Jonhson C.V, 1914. Cause and effects of soil acidity.Oklahoma Cooperative Extension Facts sheets PSS-2239.
- Mark L. Mc Farland, Vincent A. Haby, Larry A. Redmo& David H. Bade. 2001.Managing Soil Acidity. *Department of Soil and Crop Sciences*. The Agriculture Program, Texas A&M University System.
- Miyasaka SC, Hue NV & Dunn MA.2007.Aluminum. In: Barker AV, Pilbeam DJ (eds) Handbook of Plant Nutrition. Tayler and Francis Group, Boca Raton, pp. 439-497.
- Muluken Bantayehu. 2013. Study on malting barley genotypes under diverse agro ecologies of north western Ethiopia: Adet agricultural research centre, p. O. Box 08, Bahirdar, Ethiopia.
- Paoulos D. 2001.Northeast African Studies, (ISSN 0740-9133) 8:1 (New Series), P.27-52.
- Polle E, Konzak CF. 1985. A single scale for determining Al tolerance levels in cereals. *Agron Abstr.*67.
- Ryan,P.R.,Raman,H.,Gupta,S.,Horst,W.J.,andDelhaize,E.2009. Asecond mechanism for aluminium resistance in wheat relies on the constitutive efflux of citrate from roots. *Plant Physiol.* 149, 340–351.doi:10.1104/pp.108.129155.
- Schlede H.1989. Distribution of acid soils and liming materials inEthiopia. Ethiopian Institute of Geological Surveys, Ministry of Mines and Energy. AddisAbaba, Ethiopia.
- Scott B.J. & Fisher JA.1989.Selection of Genotypes Tolerant of Aluminium and Manganese. In: Robson AD, editor. Soil Acidity and Plant Growth. Australia: *Academic Press*. pp. 167–203.
- Somani, L.L. 1996. Crop Production in Acid Soils. 1st ed. Agrotech Publishing Academy, Udaipur (Rajasthan), India.
- Taye Bekele & Höfner, W. 1993.Effects of different phosphate fertilizers on yield of barley and rapeseed on reddish-brown soils of the Ethiopian highlands.*Fertilizer Research*, 34: 243–250.
- Taye Bekele. 2008. Estimation of lime requirement. Training manual for regional soil testinglaboratory heads and technicals.National soil Testing Center, Ministry of Agriculture and Rural Development.
- Temesgen Desalegn, Getachew Alemu, Tesfu, K., Tolessa, D.&Mesfin, T., 2011.Past and present soil acidity management research in Ethiopia.A review.Paper presnted at the ‘2nd National Soil and Water Management Workshop’, 28 -31. Addis Ababa, Ethiopia.
- Temesgen Desalegn.2014. Soil acidity-induced land use/cover change and management systems on soil quality parameters in the central highlands of Ethiopia.PhD Thesis.Sustainable Forest Management Research Institute University of Valladolid-INIA.Spain.pp170-173.
- Tenaye Sisay & Tesfaye Balemi. 2014. Screening of Barley Cultivars (*Hordeum vulgare ssp.vulgare* L.) for Acid Soil Tolerance Under Greenhouse Condition.*Ethiop. J. Appl. Sci. Technol.* 5(1): 58-84.
- Tigist Shiferaw and Thomas Tsige.2021. Assessment of mean reduction in morphological trait of Ethiopian barley (*Hordeum vulgare* L.) genotypes grown under limed and unlimed soil at Holetta, Central Ethiopia. *International Journal of Novel Research in Life Sciences* Vol. 8, Issue 1, pp: (15-17)
- Tovkach,A.,Ryan,P.R., Richardson,A.E.,Lewis,D.C.,Rathjen, T. M.,Ramesh,S.,etal.2013.
- Transposon-mediatedalte ration of TaMATE1B expression in wheat confers constitutive citrate efflux from root apices. *Plant Physiol.* 161, 880–892.doi:10.1104/pp.112. 207142
- Wang Jun-ping, Raman Harsh, Zhang Guo-ping, Mendham Neville&Zhou Mei-xue. 2006. Aluminium tolerance in barley (*Hordeum vulgare* L.):physiological mechanisms,genetics and screening methods. *Journal of Zhejiang University science b issn 1673-1581*.
- Zhou, G.F.,Pereira,J.F.,Delhaize,E.,Zhou,M.X.,Magalhaes,J.V.,andRyan, P. R.(2014). Enhancing the aluminium

tolerance of barley by expressing the citrate transporter genes SbMATE and FRD3. *J. Exp.Bot.* 65, 2381–2390.doi: 10.1093/jxb/eru121  
Zhao ZQ, Ma JF, Sato K& Tekeda K.2003. Differntal Al resistance and citrate secretion in barley (*Hordueum vulgare* L.) *Planta*,217:794-800.