

## A Preliminary Yield Model for Natural *Yushania Alpina* Bamboo in Kenya

M. N Muchiri\* and M. O. Muga

Kenya Forestry Research Institute, P.O Box 20412-00200, Nairobi.

\* E-mail of the corresponding author: mbaemuchiri2002@yahoo.com

*The research is financed by the Kenya Forestry Research Institute*

### Abstract

Natural bamboo resources in Kenya are about 150,000 ha and they are mainly found on the mountain slopes in the high potential areas in Mt Kenya, Aberdares ranges, Mau escarpment, Cherangani hills and Mt Elgon at high elevations between 2300 and 3200 meters above sea level. There is only one indigenous bamboo species, *Yushania alpina* (formerly *Arundinaria alpina*). It grows in single culms grouped into clumps. The main use of *Y. alpina* bamboo in Kenya is construction and fencing. There is high demand for bamboo for use in horticulture farming, handcraft, residential fencing, and cottage industry for making furniture, baskets, tooth picks and match sticks. Some companies have expressed interest in using large amounts of bamboo for bio-energy, panels, particle boards and pulp. However, bamboo raw material is not available for use because of a Presidential ban imposed on harvesting bamboo resources in all Government and Trust lands. But even if the ban was lifted, harvesting of bamboo in the natural forests cannot be prudent because there is very little information on the quantity, quality and geographical distribution of the available amount. Many entrepreneurs have continued to request the Government through Kenya Forest Service (KFS) to be allowed to extract natural bamboo for various uses. On the other front Kenya Forestry Research Institute (KEFRI) has vigorously continued to promote growing and intensive use of bamboo. These two noble ideas cannot be realized in the absence of the bamboo resource data. The objective of this study was to develop a yield model that can be used to estimate total bamboo culms biomass in a given area and the proportion that can be harvested on a sustainable basis. This study used bamboo data from 25 rectangular temporary sample plots of 10 m x 20 m in natural stands of *Y. alpina* and trees at Kieni, Kamae and Kinale in Kiambu County to estimate density and model the dependence of culms merchantable green weight and oven dry weight on diameter at breast height. The culms mean stoking density was 21000 culms ha<sup>-1</sup>. The coefficient of determination (R<sup>2</sup>) for green weight model is 0.82 and the root mean square error (RMSE) is 1.45 kg. The model calculates mean weight of old *Y. alpina* culm as 9.6 kg.

**Keywords:** Bamboo, stocking, yield, weight, culms, green weight model

### 1. Introduction

Bamboo is the fastest growing plant on earth with a short growth cycle. Some species grow as much as 1.2 m per day. Its yield is up to 25 times higher than that of fastest growing timber trees. It can be harvested in 3-5 years cycles versus 10-50 years for most softwoods and hardwoods. One bamboo clump can produce 200 poles in the five years it takes one tree to reach maturity.

Bamboo is a viable replacement for wood and is one of the strongest building materials, with a tensile strength that rivals steel. It is a critical element in the balance of oxygen and carbon dioxide in the atmosphere. It helps reduce the carbon dioxide gases blamed for global warming with some bamboo sequestering up to 12 tons of carbon dioxide from the air per hectare, which makes it an extremely efficient replenisher of fresh air (Isagi *et al* 1997). Bamboo also mitigate water pollution due to its high nitrogen consumption, making it the perfect solution for excess nutrient uptake of waste water from manufacturing, intensive livestock farming, and sewage treatment facilities.

The natural bamboo resources in Kenya are about 150,000 ha and are divided into six categories: bamboo with trees and shrubs, bamboos with trees, bamboos with shrubs, pure bamboo, shrubs with bamboos and alpine vegetations, moorland and bamboos (Kant *et al* 1992). They are mainly found on the mountain slopes in the high potential areas in Mt Kenya, Aberdares ranges, Mau escarpment, Cherangani hills and Mt Elgon at high elevations between 2300 and 3200 meters above sea level.

There is only one indigenous bamboo species, *Yushania alpina* (formerly *Arundinaria alpina*) in Kenya (Figure 1). It grows in single culms grouped into clumps. Results of studies by (Wimbush 1947, Kant *et al* 1992 and Kigomo 1994, 1995, 1988 and 2007) showed that *Y. alpina* attains a mean total height of 14 to 15 m, mean diameter of 6.0 cm. The study by Wimbush reports the average air-dry weight of *Y. alpina* culms is 100 tons per ha (Wimbush 1947) comprising of 10000 to 17000 culms ha<sup>-1</sup>. The same study reports the average cellulose content of *Y. alpina* culm is 47.5 per cent. A study by Ongugo *et al* (1999) reports the average length of a *Y. alpina* culm as 10.2 m and a dbh of 7.5 cm. Results of the study by Kant *et al* (1992) gives the mean weight of *Y. alpina* culm as 10 kg.

The main use of *Y. alpina* bamboo in Kenya is construction and fencing. Currently there is high demand for bamboo for use in horticulture farming, handcraft, residential fencing, and cottage industry for making furniture, baskets, tooth picks and match sticks. Some companies have also expressed interest in using large amounts of bamboo for furniture, bio-energy, panels, particle boards and pulp. However, bamboo raw material is not available for use because of a Presidential ban imposed on harvesting of wood and bamboo resources in all Government and Trust lands in 1986. But even if the ban was lifted, harvesting of bamboo in the natural forests cannot be prudent because there is very little information on the quantity, quality and geographical distribution of the available amount. The only information available on bamboo yields indicates that a well-managed natural stand of *Y. alpina* can produce 6 to 10 tons of mature and 1200 new culms  $\text{ha}^{-1}\text{yr}^{-1}$  (Kant *et al* (1992).



Figure 1. *Yushania alpina* at Kieni Forest

As mentioned above, many entrepreneurs have continued to request the Government through Kenya Forest Service (KFS) to be allowed to extract natural bamboo for various uses. On the other front Kenya Forestry Research Institute (KEFRI) has vigorously continued to promote growing and intensive use of bamboo. These two noble ideas cannot be realized in the absence of the bamboo resource data. The objective of this study was to develop a yield model that can be used to estimate total bamboo culms biomass in a given area and the proportion that can be harvested on a sustainable basis.

## 2. Materials and methods

The field data were collected from 25 temporary sample plots in natural pure *Y. alpina* and *Y. alpina* and trees vegetation types at Kieni (Thika district), Kamae and Kinale (Kiambu district) forest stations in June 2008. The plots were rectangular (10 m x 20 m) with the longer side expressing the slope. They were geo-referenced and located on the flat ground on the ranges where bamboo is likely to be harvested.

All bamboo clumps in a plot were identified, classified and tallied in four categories: old, mature, young and dead. The old were those covered by mold and greenish brown in colour, mature those greenish brown in colour and without mold, young were green in colour and without mold and dead those dry without foliage. However, the classification was not discrete in regards to transition from old to mature and young. From each of the first three categories, three culms were randomly selected and diameter at breast height (dbh) measured. The culms were then felled and the following parameters determined: total height, merchantable height to a top diameter of 2.5 cm and green weight (Figure 2).



Figure 2. Measurement of green weight of bamboo culms

Samples for determination of moisture content and basic density were obtained by strategically extracting billets to include the portion 10 cm below and 10 cm above the first node at the upper side of the dbh measurement point. The samples were packed in water proof polythene bags (Figure 3) and transported to KEFRI laboratory at Karura Research Centre. Two discs, each 25 mm long were extracted from each billet, weighed separately and immersed in water to obtain green volume. The discs were then oven dried at 103<sup>0</sup> C until a constant weight was obtained.

The oven dry moisture content and basic density were calculated using the following formulae:

$$\text{Moisture content (\%)} = [(\text{Initial weight} - \text{Oven dry weight}) \times 100] / \text{Oven dry weight} \text{Eq. 1}$$

$$\text{Basic density (Kgm}^{-3}\text{)} = [\text{Oven dry weight} / \text{green volume} ] \dots\dots\dots \text{Eq. 2}$$



Figure 3. Samples of Bamboo culms packed in water proof polythene bags

Graphical presentations of the culm merchantable weight as a function of dbh and merchantable height were used to identify the general form of the functions to be fitted to the data. The linear regression functions were found most suitable. The culm merchantable green weight was estimated as a function of dbh, merchantable height, basic density, and moisture content using SPSS version 15 to estimate parameters in the equations. The contribution by each of the dependent variables was tested stepwise and those found significant (at  $P=0.05$ ) retained. The best model was selected based on the following criteria:

- the adjusted  $R^2$ , – models with the highest adjusted  $R^2$  were preferred;
- residual mean square (RMS) – models with lowest RMS; and

The following five models were fitted:

- $w = a+bd$
- $w = a+bh$
- $w = a + bd*h$
- $mc = a+bd$
- usable  $wg = a+bd*h-mc$

where,  $w$  = culm merchantable green weight,  $d$  = culm dbh,  $h$  = culm merchantable height,  $mc$  = culm moisture content,  $d$  = culm basic density at dbh, and  $a$  and  $b$ , are estimated regression coefficients.

### 3. Results

More than half of the sample plots (Table 1) were located at Kieni at an elevation between 2401 to 2643 m.a.s.l, which is within the range for natural forests bamboo in East Africa.

Table 1. Geographical location of the sample plots

Forest Station	Plot No.	GPS readings		
		Northing	Easting	Altitude
Kieni	1	00 51750	3641920	2467
Kieni	2	00 51782	3641048	2479
Kieni	3	00 51817	3641227	2459
Kieni	4	00 51796	3643135	2401
Kieni	5	0051807	3641239	2460
Kieni	6	0051776	3641136	2484
Kieni	7	0051464	3640270	2509
Kieni	8	0051447	3640299	2511
Kieni	9	0051441	3640295	2521
Kieni	10	0051258	3640596	2537
Kieni	11	0052306	3640525	2516
Kinale	12	0052403	3640556	2543
Kieni	13	0052449	3640476	2553
Kinale	14	0052526	3640449	2552
Kieni	15	0052541	3640238	2566
Kieni	16	0052550	3640136	2517
Kinale	17	0052610	3640128	2540
Kieni	18	0052619	3640390	2562
Kieni	19	0052599	3640013	2566
Kieni	20	0052643	3639893	2558
Kamae	21	0050872	3638122	2612
Kamae	22	0050903	3638171	2611
Kamae	23	0050921	3638223	2643
Kamae	24	0050908	3638265	2622
Kamae	25	0050866	3638291	2628

#### 3.1 Stocking

The culms mean stoking density calculated from the data in 25 sample plots was  $20916 \text{ ha}^{-1}$  and varied from 12650 to 36900 (Table 2). The distribution by age class was: 8.4 % dead; 60.1 % old; 21.6 % mature; and 9.9 % young. The mean dbh ranged from 5.2 cm for young to 5.4 cm for the old ones. However, the mean dbh for the dead culms was not included.

#### 3.2 Merchantable green weight

The culms mean merchantable green weight ranged from 8.3 kg for the young ones to 9.6 kg for the old (Table 3). Merchantable height ranged from 8.1 m for the old to 8.4 m for the mature ones. Analysis of variance (Tables 4 and 5) showed that there were no significant differences ( $P<0.005$ ) among the means of the culms by age class in dbh, merchantable green weight, merchantable height and total height.

### 3.3 Moisture content and basic density

The mean moisture content of sampled culms ranged from 29.3 % to 457.2 % with an overall plot mean of 154.4 % (Tables 5). The young culms had the highest mean moisture content of 283.2 %, while the old ones had the least at 85.8 % (Table 6). The culms mean basic density was  $0.5\text{gcm}^{-3}$  and ranged from 0.2 to  $1.0\text{gcm}^{-3}$ . Basic density increased significantly ( $P<0.005$ ) with age with the young culms having a mean of  $0.3\text{gcm}^{-3}$  and the old  $0.7\text{gcm}^{-3}$  (Table 6).

Table 2. *Yushania alpina* stocking at Kieni, Kamae and Kinale forest stations

Plot No.	No. of culms	No. of culms $\text{ha}^{-1}$	Percentage			
			Old	Mature	Young	Dead
1	423	21150	68.3	22.7	1.9	7.1
2	352	17600	64.8	24.7	6.5	4.0
3	401	20050	45.4	29.4	17.2	8.0
4	412	20600	52.4	15.5	13.1	18.9
5	559	27950	49.0	21.1	15.9	14.0
6	354	17700	55.6	22.0	10.2	12.1
7	463	23150	78.6	10.2	8.6	2.6
8	500	25000	56.4	31.2	7.0	5.4
9	491	24550	56.6	28.7	10.4	4.3
10	327	16350	48.6	41.9	1.5	8.0
11	378	18900	51.6	24.9	0.0	23.5
12	501	25050	48.1	23.0	15.6	13.4
13	738	36900	43.4	24.8	16.0	15.9
14	509	25450	59.9	13.9	11.6	14.5
15	487	24350	57.3	28.7	5.3	8.6
16	253	12650	62.5	28.1	0.0	9.5
17	398	19900	45.7	36.2	7.8	10.3
18	422	21100	60.0	27.7	5.9	6.4
19	281	14050	74.7	6.8	3.6	14.9
20	431	21550	78.7	10.4	4.2	6.7
21	306	15300	68.0	20.3	9.8	2.0
22	262	13100	67.6	13.4	19.1	0.0
23	342	17100	76.6	3.5	19.9	0.0
24	347	17350	75.5	8.6	15.9	0.0
25	521	26050	58.0	21.3	20.7	0.0
<b>Mean</b>	418	20916	60.1	21.6	9.9	8.4

Table 3. Descriptive statistics of *Yushania alpina* culms at Kieni, Kamae and Kinale forest stations

Culm class	Statistic	Dbh, cm	Merchantable height, m	Total Height, m	Weight, kg
Old	Mean	5.4	8.1	10.5	9.6
	N	75.0	75.0	75.0	75.0
	Std. Deviation	0.7	1.7	1.7	3.4
	Minimum	4.0	4.6	6.0	3.0
	Maximum	7.0	13.7	13.6	19.5
	Std. Error of Mean	0.1	0.2	0.2	0.4
Mature	Mean	5.3	8.4	11.0	9.1
	N	75.0	75.0	75.0	75.0
	Std. Deviation	0.8	1.9	1.9	3.5
	Minimum	3.3	4.2	4.8	3.0
	Maximum	7.4	13.5	14.0	17.5
	Std. Error of Mean	0.1	0.2	0.2	0.4
Young	Mean	5.2	8.2	10.9	8.3
	N	69.0	69.0	69.0	69.0
	Std. Deviation	1.0	2.2	2.0	3.5
	Minimum	2.8	2.4	4.8	0.6
	Maximum	7.0	12.4	14.2	16.5
	Std. Error of Mean	0.1	0.3	0.2	0.4

Table 4. Analysis of variance of dbh, merchantable height, total height and green weight of *Yushania alpina* culms at Kieni, Kamae and Kinale forest stations

Variable	Source	Sum of Squares	df	Mean Square	F	Sig.
Dbh, cm	Between Groups	1.2	2.0	0.6	0.9	0.4
	Within Groups	151.1	216.0	0.7		
	Total	152.3	218.0			
Total Height, m	Between Groups	9.7	2.0	4.9	1.4	0.3
	Within Groups	751.6	216.0	3.5		
	Total	761.3	218.0			
Merchantable height, m	Between Groups	2.7	2.0	1.3	0.4	0.7
	Within Groups	819.1	216.0	3.8		
	Total	821.8	218.0			
Green weight, kg	Between Groups	54.2	2.0	27.1	2.3	0.1
	Within Groups	2576.0	216.0	11.9		
	Total	2630.2	218.0			

Table 5: Descriptive statistics of moisture content and basic density of *Yushania alpina* culms at Kieni, Kamae and Kinale forest stations

Plot No.	N	Moisture content, %					Basic density, gm <sup>-3</sup>				
		Mean	Std. Dev	Std. Error	Min	Maxi	Mean	Std. Dev	Std. Error	Minimum	Maximum
1	9	136.2	96.4	32.1	35.0	271.2	0.54	0.20	0.07	0.29	0.84
2	9	143.0	111.1	37.0	58.0	368.6	0.56	0.21	0.07	0.23	0.78
3	9	157.6	104.2	34.7	29.3	319.1	0.49	0.18	0.06	0.25	0.69
4	9	151.0	99.8	33.3	59.2	348.3	0.53	0.22	0.07	0.23	0.95
5	9	158.1	97.0	32.3	60.8	312.9	0.51	0.20	0.07	0.26	0.78
6	8	147.8	102.8	36.3	57.3	318.1	0.58	0.23	0.08	0.25	0.80
7	9	160.3	85.9	28.6	80.2	324.7	0.48	0.16	0.05	0.24	0.70
8	9	147.6	71.8	23.9	71.0	254.3	0.49	0.15	0.05	0.29	0.71
9	9	142.3	85.1	28.4	66.0	296.8	0.52	0.18	0.06	0.26	0.74
10	9	151.9	47.7	15.9	86.3	251.5	0.46	0.10	0.03	0.28	0.65
11	6	82.6	16.0	6.6	60.5	101.9	0.65	0.08	0.03	0.56	0.77
12	9	166.8	136.1	45.4	72.1	457.2	0.51	0.20	0.07	0.18	0.69
13	9	148.7	122.1	40.7	62.9	357.2	0.55	0.21	0.07	0.22	0.75
14	9	120.8	85.2	28.4	53.4	257.5	0.62	0.23	0.08	0.29	0.85
15	9	179.4	138.6	46.2	61.1	366.0	0.51	0.23	0.08	0.22	0.78
16	6	113.8	50.7	20.7	75.0	209.1	0.57	0.13	0.05	0.35	0.71
17	9	145.8	103.4	34.5	48.4	333.2	0.55	0.22	0.07	0.24	0.85
18	9	195.7	139.8	46.6	76.8	395.0	0.47	0.21	0.07	0.21	0.69
19	9	187.0	124.9	41.6	68.2	373.8	0.48	0.21	0.07	0.22	0.72
20	9	165.9	127.8	42.6	55.0	354.9	0.54	0.25	0.08	0.22	0.81
21	9	158.6	82.8	27.6	83.5	280.3	0.48	0.16	0.05	0.27	0.65
22	9	165.6	88.0	29.3	84.5	307.0	0.47	0.15	0.05	0.26	0.65
23	9	156.7	106.0	35.3	77.4	346.2	0.51	0.18	0.06	0.23	0.67
24	9	161.6	80.6	26.9	92.6	310.2	0.47	0.14	0.05	0.25	0.61
25	9	176.1	99.2	33.1	90.0	315.3	0.45	0.15	0.05	0.25	0.62
Mean	218	154.4	98.2	6.6	29.3	457.2	0.52	0.19	0.01	0.18	0.95

Table 6. Descriptive statistics of oven dry weight, basic density and moisture content of *Yushania alpina* culms by age class at Kieni, Kamae and Kinale forest stations

Parameter	Culm class	N	Mean	Std. Dev	Std. Error	Min	Max
Oven dry weight, kg	Old	75.0	5.2	2.0	0.2	1.9	12.1
	Mature	75.0	4.6	1.9	0.2	1.3	12.2
	Young	69.0	2.3	1.1	0.1	0.2	6.1
	Total	219.0	4.1	2.1	0.1	0.2	12.2
Basic density, gm <sup>-3</sup>	Old	75.0	0.7	0.1	0.0	0.4	0.9
	Mature	74.0	0.6	0.1	0.0	0.4	1.0
	Young	69.0	0.3	0.1	0.0	0.2	0.8
	Total	218.0	0.5	0.2	0.0	0.2	1.0
Moisture content, %	Old	75.0	85.8	26.6	3.1	29.3	215.2
	Mature	74.0	103.8	32.1	3.7	35.0	209.1
	Young	69.0	283.2	63.8	7.7	67.0	457.2
	Total	218.0	154.4	98.2	6.6	29.3	457.2

**Linear Regression models for predicting merchantable green weight and oven dry weight of *Y. alpina***

The best linear regression model for predicting green weight of *Yushania alpina* culms based on dbh (Table 8) is:  
 $W = -1.11 + 0.36 D^2$  ..... Eq. 3

Where, W = culm merchantable green weight, D = culm dbh, cm.

Table 8: Linear regression models for predicting merchantable green weight of *Yushania alpina* culms based on dbh, merchantable height, and total height

Model	N	Constant	Coefficient	Adjusted R Square	Std. Error of the Estimate of R	F-value	Significance
Dbh <sup>2</sup>	218	-1.11	0.36	0.82	1.45	976.02	0.00
Dbh	218	-10.64	3.74	0.81	1.52	914.25	0.00
Dbh*merchantable height	218	0.26	0.20	0.79	1.58	834.47	0.00
Merchantable height	218	-2.29	1.37	0.59	2.23	312.46	0.00
Merchantable height <sup>2</sup>	218	3.26	0.08	0.57	2.31	276.17	0.00
Total height <sup>2</sup>	218	2.03	0.06	0.41	2.66	154.16	0.00
<b>Total height</b>	218	-3.28	1.14	0.37	2.75	130.90	0.00

The best linear regression models for predicting oven dry weight of *Yushania alpina* culms based on green weight and dbh is equation 4 and the statistics are presented in Table 9.

$ODW = 1.04 + 0.06(dbh*gw)$  -----Eq. 4

Where, ODW = culm merchantable oven dry weight, kg, dbh = diameter at breast height and gw = green weight.

Table 9. Linear regression models for predicting oven dry weight of *Yushania alpina* culms based on green weight, dbh and total height

Model	N	Constant	Coefficient	Adjusted R Square	Std. Error of the Estimate of R	F-value	Significance
Dbh <sup>2</sup>	218	-0.213	0.151	0.39	1.66	140.6	0.000
Dbh	218	-4.33	1.16	0.39	1.65	142.43	0.000
Merchantable height	218	-0.86	0.60	0.30	1.78	93.25	0.000
green weight	218	-0.21*	0.47	0.60	1.34	329.07	0.000
green weight <sup>2</sup>	218	1.58	0.04	0.28	1.80	54.89	0.000
<b>Green weight*dbh</b>	218	1.04	0.06	0.55	1.42	267.67	0.000

\*Not significant

**4. Discussion**

The culms mean stoking density (21000 culms ha<sup>-1</sup>) in this study is one and half times that reported in earlier studies in the same geographical area (Wimbush, 1945; Kigomo 1995, 1988, 2007, and Kant *et al* 1992). The possible reason is that there has been no harvesting of bamboo culms in the study area for more than a decade, which is more than the life cycle of bamboo. This may also tell why the culm mean dbh is lower than that reported by earlier studies. The mean weight of old *Y. alpina* culm (9.6 kg) calculated in this study is in line with that reported by Kant *et al* (1992).

The total number of the old culms and their green merchantable weight is 12600 and 121 tons ha<sup>-1</sup>, respectively. The mean oven dry weight of the same culms is 70 tons ha<sup>-1</sup> and is in line with the 100 kg ha<sup>-1</sup> air dry biomass reported by Wimbush (1945). The merchantable green weight of culms is comparable to 116.5 tons ha<sup>-1</sup> of the above ground biomass of the *Phyllostachys pubescens* (the largest bamboo species in the World) reported by Isagi (1994).

Just like the case with all living things, *Y. alpina* culm stops growing as they age. Consequently, the old and dead culms which accounts for 60.1 % and 8.4 % of the total green merchantable culms, respectively do not put on new growth. It would therefore be prudent that these culms are harvested to supply raw material to many individuals and companies that have expressed interest in harvesting large amounts of bamboo for various uses. Nonetheless, the proponents of the environment may raise concern that cutting 68.5 % of bamboo may expose the environment to rainfall run-off, and loss of soil and ground moisture. This would interfere with the water availability in the region because natural bamboo forest is only found in the country's five water towers. With the foresaid notwithstanding, the alternative of not cutting is a worse scenario because bamboo has very poor regeneration if left unattended. Indeed, cutting bamboo triggers prolific regeneration as evidenced along the

main road at Kieni forest station (Figure 1). The compromise would be to cut slightly less than 50 % of the old and dead culms which is in agreement with results of many silvicultural studies that reducing density of a stand by half does not result in growth shock or expose the environment to harsh weather conditions.

## 5. Conclusion

The results of this study lead to the following conclusions about bamboo in undisturbed stands at Kamae, Kieni, Kinale and Ragia Forest Stations:

- The mean stocking density of the bamboo culms is 21000 culms ha<sup>-1</sup>;
- 68 % of bamboo culms are either old (60 %) or dead (8 %) and are therefore suitable for immediate use;
- The mean diameter at breast height, mean merchantable height to 2.5 cm top diameter and mean green weight of utilizable bamboo culms are 5.4 cm, 8.1 m and 9.6 kg, respectively;
- 50 % of the culms of a bamboo stand comprising the dead and old stock can be harvested without negatively affecting the environment;
- One ha of undisturbed bamboo stand can produce 10 500 bamboo culms with green and oven dry weight of 100 tons and 55 tons, respectively.
- Selective cutting of bamboo triggers regeneration;

## 6. Recommendations

- Validate the model using data collected in other bamboo forests in Kenya;
- The authors recommend that, the Kenya Forest Service considers using the results of this study to embrace scientific management of bamboo resources and lifting of the ban on harvesting of bamboo culms; and
- Undertake further studies to incorporate the potential of *Y. alpina* in carbon sequestration.
- The linear regression models developed for estimation of bamboo culm weight can be tabulated for easy application by those interested in calculating bamboo culm weight for various purposes.

## References

- Isagi, Y. 1994. Carbon stock and cycling in a bamboo *Phyllostachys bambusoides* stand. Ecological-Research. 1994, 9: 1, 47-55.
- Isagi, Y; Kawahara, T; Kamo, K; and Ito, H. 1997. Net production and carbon cycling in a bamboo *Phyllostachys pubescens* stand. : Plant-Ecology. 1997, 130: 1, 41-52.
- Kant, H., Kigomo, B. N. and Ndambiri. J.K. (1992). Development of Bamboo in Kenya. Kenya Forestry Master Plan.
- Kigomo, B.N. 1988. Distribution, cultivation and research status of bamboo in Eastern Africa Kenya Forestry Research Institute Ecological Series; Monograph No. 1: October, 1995. Nairobi, Kenya. 19 pages.
- Kigomo, B.N. 1994. J.M. Were, and G.O. Sigu, 1994. The establishment and growth of *Dendrocalamus hamiltonii* Nees and Arn under field conditions in Kenya". East African Agricultural Forestry Journal.
- Kigomo, B.N. 1995. Guidelines for establishment and management of bamboo in Kenya. Kenya Forestry Research Institute Occasional (Management) Paper No. 1: October, 1995. Nairobi, Kenya. 31 pages.
- Kigomo, B.N. and Sigu G.O., 1996. Establishment and growth yield of field trials of exotic and indigenous bamboo species in Kenya. East African Agricultural Forestry Journal 61 (3) 213-223.
- Kigomo, B.N. 2007. Guidelines for growing bamboo. KEFRI Guideline Series.: No. 4. Kenya Forestry Research Institute; Nairobi, Kenya.
- Ongugo, O, Sigu, G.O, Kariuki, J.G, Luvanda, A.M., and Kigomo, B.N. 1999. Production-to-consumption System: A case study of the Bamboo Sector in Kenya.
- Wimbush, S.H. (1947). The African Alpine Bamboo. The East African Agricultural Journal, Vol. 24, No 1, 1945.



This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

## CALL FOR JOURNAL PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

## MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

