

Assessment the Status of Agro-Chemicals Utilization and Honeybee Poisons Plants in Sidama Zone, SNNPR, and Ethiopia

Dinku Negash¹ Abebe Jenberie² Asaminew Tassew²
1.Southern Agricultural Research Institute
2.Bahir Dar University

Abstract

The study was conducted with the objective of identifying honeybee poisoning plant and assessing the status of agro-chemicals utilization in Sidama zone, Southern Ethiopia. Semi-Structured questionnaires were developed and 160 individuals (80 beekeepers and 80 non beekeepers) were interviewed on voluntary basis. The result of this assessment indicated that 75 (93.75%) of beekeepers complained presence of plant poisoning on honeybees in the study area. Ten plants were complained having poisoning effect on honeybees by beekeepers in the study area. Among these plants *Lanthana camara*, *Euphorbia continifolia*, *Climatis flammula*, *Ranclus multifides*, *Discopodium penninervium*, *Climatisinte grifolia*, *Datura metal linn*, *Sesbania sesban*, *Phytolaca americana* and *Justitia schemperina*, were the most frequently complained toxic plants in the study area. Seven different agro-chemicals (2,4-D, Agrothoate 40% Malathion 50%, pyriban 48%, Diazinon 60%, Macezeb 80% and Pallas 45 OD) were commonly applied on various crop in the study area. The main sources of these agro-chemicals are cooperative, legal and illegal traders. 95% of farmers used empty containers for household purpose. 3.25% of the respondents left in farm while 1.25% of the respondents indicated that they have burnt. Most of agro-chemical users (80%) of the respondents had no training on how to apply Agro-chemicals safely to honeybees, themselves and environment and 97.5 % of respondents applied agro-chemicals without following the recommended instructions in addition 95.62% of the respondents did not use protective clothes when spraying. Therefore, further study on complained poisonous plants and toxic chemicals in the study area, and proper utilization agro-chemicals are important to minimize poisoning of honeybee.

Keywords: Agro-chemicals, Honeybee poison plants, Sidama zone

DOI: 10.7176/JRDM/85-04

Publication date: May 31st 2022

Chapter 1: INTRODUCTION

1.1. Background and Justification

Naturally, there are certain species of honey plants whose pollens, nectars and honeys are toxic to bees. Accordingly, poisonous plants have caused extensive losses to the livestock industry in many parts of the world mainly east Africa including Ethiopia since the days of early settlements (Abera Dereje, 2015). Further, its effect is still a significant threat in many parts of the globe. The toxic effects of poisons plants in livestock after ingestion or absorption includes, physical disturbance, decrease productivity and death (Abera Dereje, 2015).

The poisonous nature of whole plant or any plant part may be due to production of toxic substances such as alkaloids, glucosides, amines, toxalbumins, picrotoxins, resins and saponins many of which are harmful to man and animal life, at least under certain conditions (Tom, 2011). Nectar and pollen collected from flowering plants are the main food resources of social bees. The nutritive value of pollen grain and nectar is known to be a potential protein and carbohydrate sources for the colony respectively (Queiroz and Contrera, 2014). On the other hand, various plant species contain secondary compounds in nectar and pollen that could be toxic to pollinators, including bees. For instance, the Almond tree (*Amygdalus communis L- Rosaceae*) contains the cyanogenic glycoside amygdalin that releases cyanide. *Amygdalin* is found in the nectar and pollen of almond trees and consumption of this pollen can be toxic to honeybees (Kevan, 2005).

The introduction of pesticide in Ethiopia to control agricultural pests' dates back to the 1960's (EPA, 2004). Although chemical pesticide use in Ethiopia was historically low and the volume fluctuates across the pesticide types, recent developments in increased food production and expansion in floriculture industry have resulted in the importation of about 3346.32 metric tons of agro-chemicals annually and higher consumption (Gizachew Assefa, 2011). The most devastating phenomena that curtails the productivity of honeybee colonies, is poisoning of honeybees by agro-chemicals such as fungicides, pesticides, and herbicides. This daunting challenge not only affect the wellbeing of honeybees but also the wellbeing of human beings who utilize its products and also the ecology in which honeybees are main actors in pollination of plants to keep the ecology balances (Assemu Tesfa *et al.*, 2013). Poisoning of honeybees by agro-chemicals has been increased from time to time. Some beekeepers lost totally their colonies due to agro-chemicals (Kerealem Ejigu *et al.*, 2009). The promotion of some agricultural inputs such as pesticides and herbicides for cereal crops production and the use of deadly chemicals for malaria eradication program have substantially reduced honey production (Gezahegn Tadesse, 2001). According to Desalegn Begna (2015), there is a growing pesticides grievance on honeybee population and their

products decline with considerable economic impacts on beekeepers. Indiscriminate uses of pesticides caused fatalities 22987 honeybee colonies and incurred economic loss amounting of \$819291.37 USD in Bure districts of Amhara Region.

Similarly beekeepers and beekeeping experts of the SNNPR have always blame the indiscriminate use of agro-chemicals and honeybee poisonous plants for the loss of honey bee colonies in the area. They repeatedly reported that honeybee colony population and swarms, honey production had declined in the area. According to their reports these are critical problems particularly during September–November when most agrochemicals are applied in cultivated field and dearth period honeybees are exposure to poisonous plants in the area. However, there were no substantial quantitative data on these cases. Therefore this study was initiated to assess major honeybee poisonous plants and document utilization of the agro-chemicals in Sidama zone

2. Objectives

2.1.1. General objective

The general objective of the study was to identify the honeybee poisoning plants and commonly used agro-chemicals, which are negatively affecting honeybee health and evaluating the economic importance of the unwise chemical application in the study area.

2.1.2. Specific Objectives

- To identify honeybee poisonous plants in the study area
- To assess the status of agrochemical utilization among farmers in the study area

Chapter 3: MATERIALS AND METHODS

3.1. Description of the Study Area

The Southern nations, nationalities and peoples region (SNNPR) is one of the regional states of Ethiopia located in the southern parts of the country, covering a total area of 105,887.18square kilometers. Geographically, it lies between 4°43'N - 8°58'N, and 34°88'-E 39°14'E. SNNPR is structured into 13 administrative zones, 133 woredas and 3512 kebeles and its capital is Hawassa. The total population surpasses 15,745,000, of which about 89% of the population lives in rural areas of the region (CSA, 2008). According to CSA (2015/16) report, the region owns 7,938,490cattle population, 10,228,460 small ruminants 6,586,140poultry, 640620equines and 726,960honeybee colonies.

Sidama zone, located in the SNNPR state at a distance of 273 km South of Addis Ababa with GPS coordinates of latitude: 5° 45" and 6° 45"north and longitude, 38° and 39° East. Sidama is bordered on the South by the Oromia Region (except for a short stretch in the middle where it shares a border with Gedeo zone), on the West by the Bilate River, which separates it from Wolayita zone, and on the North and East by the Oromia region. The zone has a total area of 10,000 km² and 2,954,136 total population (of which 1,491,248 are males and 1,462,888 are females). When we see its land use features, the zone's total land falls in to the following categories: 48.70 % cultivated , 2.29 % forest, 5.04 % shrub and bush land , 17.47 % grazing land and 18 % uncultivated land (MoARD, 2016).

Sidama has a variety of climate conditions; warm climate covers 54% of the area. Elevation of the area ranges from 1500 to 3500 m.a.s.l. and mean annual rainfall varies between 1200 mm to 1999 mm with 15°C to 19.9 °C mean annual temperature. The main crops cultivated in the zone are coffee, Enset, Teff, Barley, Wheat, Maize, Chickpea, Bean, Pea, Lentil and Haricot bean. The major livestock raised in the zone are cattle, horses, donkey, goats, sheep, mules, and chicken and honeybee colonies (MoARD, 2016).

Hula district is geographically located between 6°28'60N and 38°31'0E in the southern parts of Sidama zone with an altitude ranging from 2759 to 2829 m.a.s.l. The district, representing the highland agro-ecology of the study area, has a total human population of 129,263of which64,551 are men and 64,712 are women; whereby 6,410 or 4.96% of its population is urban dweller. The district has high potential for beekeeping and crop production when compared to the other highland district

Shebedino district is geographically located between 6°50'N and 38°30'E in the northern part of Sidama zone at a distance of 15 km from Hawassa. The district is representing the midland agro-ecology of the study area. The altitude of the district ranges from 1980 to 2500 m.a.s.l where 15°C to 30°Cis the average daily air temperature with 558 mm mean annual rainfall. This district has a total population of 233,922, of which118,026 are men and 115,896 women; whereby 11,831 or 5.06% of its population are urban dwellers (Sidama zone Agricultural and Rural Development office).

Loka Abaya district, representing the lowland agro-ecology of the study area, is located at the western parts of Sidama zone. Geographically, it is located between 6° 50' 0" N and38° 30"E with an altitude of 1200 to 2300 m.a.s.l. The districts estimated total population of 99,233, of which50, 603 is men and 48,630 are women; whereby 1,059 or 1.07% of its population is urban dwellers.




Figure 1. Maps of study areas

3.2. Sampling Methods and Sample Size Determination

This particular study was conducted in three representative districts of Sidama zone selected from each of the three agro-ecologies (highland, midland and lowland). Agro-ecology representation, beekeeping and crop potential, status and intensity of agrochemical application and accessibility were used as districts and kebele selection criteria. Accordingly, eight Kebeles (3 from lowland, 3 from midland and 2 from highland) from three representative districts namely Hula (representing highland), shebedino (midland) and Loka Abaya (lowland) were selected using purposive random sampling technique.

In order to collect primary data from respondents, a formula described by Yamane 1967 was employed to select sample respondents. Accordingly, 160 respondents (80 beekeepers and 80 non beekeepers) out of 270 model farmers identified in the study area were selected using a purposive and stratified random sampling technique.

$$n = \frac{N}{1 + N(e)^2}$$

Where:

n = sample size

N = total population

e = sampling error (*e*=0.05)

In this case, a total of 20 sample respondents per kebele were selected purposively for primary data collection based on their relative and/or better beekeeping experiences, intensity of agro-chemicals application and representation of each agro-ecology. Checklists were prepared and framers' perceptions were assessed through focused group discussions from each kebele. In addition, beekeeping experts and technicians, crop protection experts, agro-chemical traders at various levels and other relevant stakeholders were considered as a part of the data collection procedure during the survey work.

Table 1. Number of respondents in respective kebeles and agro-ecologies

Kebeles	woredas	Agro-ecology	Sample size	Total sample size
Gasy		Highland	20	
Buchesa Gobe	Hula	Highland	20	40
Remeda		Midland	20	
Teremesa	Shebedino	Midland	20	60
Sedeka		Midland	20	
Sela kebedo		Lowland	20	
Arede Galye	Loka Abaya	Lowland	20	60
Danshy Gambela		Lowland	20	
Total sample size			160	160

3.3. Data Sources and Methods of Data Collection

3.3.1. Data Sources

3.3.1.1. Primary data collection

Primary data was collected through semi-structured questionnaire, check lists, laboratory results and observation. Beekeepers, non-beekeepers crop growers, bee experts, crop protection experts, agrochemical whole sellers and retailers were a part of the interview.

3.3.1.2. Secondary data collection

The study also included secondary data from various sources. Secondary information was obtained from documents of Regional Beuro of agriculture and respective districts office of agriculture and rural development and from various published and unpublished literatures.

3.3.2. Methods of data collection

Both quantitative and qualitative methods were used to obtain validated and substantiated findings in offsetting weaknesses inherent within one method with the strengths of the other method. Quantitative data were collected through survey and laboratory analysis.

3.3.2.1. Survey

Information was collected using semi-structured questionnaire by the researcher and trained enumerators. Before the actual survey the questionnaire was pre-tested. The main data was collected through the survey included:

- Socio-economic characteristics of the households: - gender, age, family size, education level, land holding and livestock holding.
- Beekeeping production system: Source of colony, trend of honey production and honeybee colony
- Uses of agro-chemicals: The type of agro-chemicals, extents of uses, means of application and the hazards it might cause and level of experiences in using agro-chemicals
- Constraints of apiculture in the areas: Trends of agro-chemicals application, identified poisonous plants, Pests and diseases data were collected.

3.4. Data Management and Statistical Analysis

The data collected from the survey were coded and stored into computer loaded SPSS software programs version 20, and cleaned for consistency and accurateness. The statistical analysis used in the study varied depending on the type of variable and information obtained. Summarized data were presented in the form of tables and figures. Chi-square was used to test the significance difference between or among values whenever necessary and the constraints of the apiculture in the study areas were prioritized using rank index.

Chapter 4: RESULTS AND DISCUSSION

4.1. Socio-Economic Characteristics of Household

The general characteristics associated with beekeeping and non-beekeeping households distributed by gender, age, marital status, farmland holding and educational level are presented in Table 2 and figure 1. Of the total respondents, 96.25% and 100% of the beekeepers and non-beekeepers respectively were male while the remaining 3.75% of beekeepers were females. This is in line with the finding of Addis Getu and Malede Birhanu (2014) and Dereje Shibru *et al.*(2016) who reported that majority of farming activities are duties of male society. Moreover, the limited numbers of female participation in beekeeping activities agree with Tessega Belie (2009) who indicated that only 1.7% were female. So that few women's are engaged in beekeeping activity in the area and not economically empowered through beekeeping.

The age of about 81.25% of beekeepers and 83.75% of non-beekeepers interviewed in the study area ranges

between 25-60 years old. This result showed that people are engaged in beekeeping activities both at the younger and older ages. This result correlates with Tewodros Alemu *et al.*(2015), Dereje Shibru *et al.*(2016), Sintayehu Fetene (2016), Wolaye Kiros and Teklberhan Tsegay (2017) results, who reported that beekeeping learnt through generation and practiced by all economically active age groups ranges between 25-60 years old. Of the total interviewed households, 98.1% were married and 1.9% unmarried. Marriage helps farmers to sustainably engage in crop production and beekeeping practices to ensure the livelihood of their families and option for asset building. This result agreed with Haftu Kebed and Gezu Tadess (2014), who stated that high percentage of the respondents (96.8%) were married and engaged to agricultural production.

Concerning to level of education, the highest percentage 27 (33.75%) and 14 (17.5%) of non-beekeepers and beekeepers were illiterate respectively. This result indicates that most of beekeepers are educated compared to non-beekeepers. In other districts, numbers of illiterate in non-beekeepers are more than the beekeepers. Time and thereby improve the productivity and production of their agricultural activities. Education affects technology adoption, household income and socio-economic status of the family. Figure 2, explains why most of the farmers engaged in beekeeping activity are more literate compared to non-beekeepers who are slower to adopt other agricultural activities. This result agree with Malede Birhan (2015), who stated that education is an important and one entry point for fast transfer of knowledge on improved beekeeping.




Figure 2. Educational status of respondents

Table 2. Household Socio-demographic characteristics

Category	Variables	Beekeepers(n=80)		Non beekeepers (n=80)	
		N	%	N	%
Gender	Male	74	96.25	80	100
	Female	6	3.75	0	0
	Total	80	100	80	100
Age	Below 25	4	5	-	-
	25–60	65	81.25	67	83.75
	>60	11	13.75	13	16.25
	Total	80	100	80	100
Marital status	Married	77	96.25	80	100
	Divorced	-	-	-	-
	Widowed	-	-	-	-
	unmarried	3	3.75	-	-
Total		80	100	80	100

The average farmland holding of the respondents were 1.43±0.07ha. The data has described that the overall mean of land holding in the study area is similar with the mean national average (1-1.5 ha). Similarly, Chala Kinati (2010) and Bekele Tesfy (2015) reported that average land holding of farmers in Oromiya region was 1.48±0.09 hectares.

Based on this study the overall mean family size of the beekeepers were 5.6±0.13 per household with a maximum of 9 and a minimum of 2 peoples. According to Bekele Tesfy (2015) result, family sizes of Bale zone

of Oromiya region was 8.01 ± 0.27 . So that, this finding agreed with the above report.

Table 3. Family size and land holding of the respondents

Variables	Total sample size (N=160)							
	Shebedino(N=60)		Hula (N=40)		Loka Abaya(N=60)		Over all	
	Range	Mean±SE	Range	Mean±SE	Range	Mean±SE	Range	Mean±SE
Family size	2-8	5.2±0.23	3-9	5.85±0.26	2-9	5.9±0.22	2-9	5.6±0.13
Total land holding	0.5-3	1.15±0.09	0.5-2	1.52±0.15	0.5-3	1.65±0.12	0.5-3	1.43±0.07

N = Number of cases, SE = Standard error of mean

The main income sources for beekeepers were crop 0.35(1st), beekeeping 0.33 (2nd) and livestock 0.32(3rd) ranked in decreasing order. Honeybee products marketed locally to provides incomes for various beekeepers in the study areas. As indicated in the table below beekeepers had not full of confidence in beekeeping as main income source rather than crop production in the study area. This is due to traditional production system, lack of knowledge and migratory nature of the colony was the major challenges for hindering beekeeping thought as main income source in the study area. whereas, crop and livestock were the 1st and 2nd ranks income source for non-beekeepers.

Table 4. Major income source of beekeepers and non-beekeepers

Farming activity	Bee keepers				Non beekeepers			
	1st	2nd	3rd	Rank index	1st	2nd	3rd	Rank index
Crop production	26	22	34	0.35(1 st)	80	0	0	0.52(1 st)
Livestock production	7	44	29	0.32(3 rd)	0	75	0	0.48(2 nd)
Bee keeping	47	14	18	0.33(2 nd)	0	0	0	0(3 rd)

4.2. Beekeeping activities and Honey production trends in the study area

4.2.1. Source of colonies

Honeybee colonies can be obtained from different sources. Though it varies between beekeepers and different localities. The study indicated that there is a huge indigenous knowledge in beekeeping and different of source of honeybee colonies due to various reasons. According to the survey result, 92.5% and 7.5% of respondents replied that active season swarm catching and gift from parents are sources of their colonies respectively (Table 5). Interestingly, colony purchase has never been used as a source of honeybee colonies to start beekeeping in the study area. This is most probably because of poor extension services system, poor adoption of improved beekeeping technologies, high costs of beekeeping equipment. This result is partially in line with findings of Addis Getu and Malede Birhan (2014), Haftu Kebede and Gezu Tadesse (2014), Teklu Gebretsadik and Dinku Negash (2016), Welay Kiros and Tekleberhan Tsegay (2017), Abebe Mitikie (2017). They have stated that swarms catching are the main sources of honeybee colonies to start beekeeping in various parts of the country. Moreover, bee colony marketing is not a trend or is not common in the study area. This is due to the fact that farmers could catch colonies easily when reproductive swarming is active.

Table 5. Sources of honeybee colonies

Sources of honeybee colonies	Percentage of respondents
Catching swarms	92.5%
Gift from Parents	7.5%
Purchasing	-
Total (N=80)	100%

4.2.2. Trends of bee colonies in the study area

From the total respondents, about 78.8% of beekeepers replied that honeybee colonies in the study area is decreasing over the past year due to indiscriminate use of agro-chemicals, shortages of bee forages and pests and predators are the major limiting constraints. The remaining 16.3 % and 5% of the respondents replied that beekeeping practice is increasing and remain in the same in the past years respectively (Fig.3). Even if the government is giving a special attention to beekeeping and high involvement of different NGO to this sector is higher, number of bee colonies is steadily decreasing from time to time due the trends of agrochemicals application increase.




Figure 3. Trends in the numbers of honeybee colonies

4.2.3. Trends of honey yield in different hive type

According to the survey result, the trend in honey yield from different hive types has decreased during 2013-2016 but increased in 2017. The collected data revealed that the average honey yield obtained from traditional hives has steadily reduced from 5.72 kg/hive/year to 5 kg/hive/year during 2013-2016 and increased to 6.06 kg/hive/year in 2017. Similarly, the average honey yield obtained from transitional hive reduced 12.16 kg/hive/years to 10.64 kg/hive/year and 19.54 kg/hive/year to 16.91kg/hive/year in movable frame hive through year 2013 to 2016. whereas, honey yield increased to 11.5 kg/hive/year from a transitional hive and to 18.91 kg/hive/year from movable frame hives in 2017(Fig.4). Honey yield achieved better performance in 2017 rather than the years before. This is due to availability of ample bee forage and suitable climatic conditions for honeybees to provide high honey yield in the study area. This results is comparable to Tewodros Alemu *et al.* (2017) result, who stated that hive productivity trend has increased from 2011 to 2014 in Sekota district. Honey yield in modern hives (zander) and transitional hives (Kenya top bar) was increasing from 2011 to 2014




Figure 4. Trends of honey yield in various hive types

4.2.4. Mean amount of honey yield from different hive types per annum (kg)

According to the survey result, traditional, transitional and frame hives showed a very significant difference in honey yield at ($p < 0.05$). Over all mean amount of honey yield from traditional, transitional and movable frame hives were 5.74 ± 0.17 kg, 11.37 ± 0.25 kg and 18.32 ± 0.64 kg per annum respectively (Table 6). This result comparable with results of Welay Kiros and Teklebirhan Tsegaye (2017) who showed that honey yield from traditional hive was significantly lower than transitional and frame hives. Similarly Atsbaha Hailemariam *et al* (2015) also explained that there was significant difference between traditional (12.79 kg) and frame (28.29 kg) hives in his study area. Moreover, productivity of frame and transitional hives in this study were much higher than that of the traditional hives. This is due to the fact that transitional and frame hives did get better management such as provision of wax foundation sheets, recycling of harvested combs and higher frequency of honey harvesting. In addition, the variation in productivity of traditional, transitional and frame hive might be attributed to the suitability of the improved hives to improved management.

Table 6. Mean amount of honey produced from different hive types per annum (kg).

Type of hive	Total sample size (N=80)							
	Shebedino(N=30)		Hula (N=20)		Loka Abaya(N=30)		Over all	
	Range	Mean±SE	Range	Mean±SE	Range	Mean±SE	Range	Mean±SE
Traditional	3-8	5.37±0.16	2-7	5.36±0.20	3-8	5.67±0.15	2-8	5.47±0.17 ^a
Transitional	8-15	11.24±0.26	8-14	11.83±0.27	9-15	11.04±0.23	8-15	11.37±0.25 ^b
MF	10-30	18.82±0.82	10-25	17.29±0.48	14-30	18.87±0.62	10-30	18.32±0.64 ^c

Superscript indicate significantly different at $p < 0.05$

N = Number of cases, SE = Standard error of mean and MF= Movable Frame

4.3. Major agricultural constraints in the study area

Agriculture, not only in the study area but also in nationwide is facing various constraints in its major components (crop production, livestock production and beekeeping). More specifically, shortage of cropland, drought, disease and parasite, lack of inputs, lack of draft animals, availability of different weeds and low soil fertility in crop production (Table 8) and shortage of animal feeds, shortage of grazing land, prevalence of various diseases and low productivity of local animals in livestock production (Table 8), were the majorly identified agricultural culprits in the study area.

Similar to the other components of agriculture, beekeeping has been explained to receive various constraints hindering honey production. Accordingly, respondents in this study have clearly identified that, random use of agro-chemicals (15.63%), presence of pests and predators (15.27%), high rate of absconding (14.54%), increased cost of production (14.36%), lack of bee forage plants (14%) and drought (13.45%) were the major constraints in beekeeping (Table 7). Even if their level of importance varied from place to place, miss use of agro-chemicals ranked first as an important constraint in honeybee production system, which was negatively affecting colony productivity in the study area. This finding agrees with results suggested by Gidey Yirga and Kibrom Ftwi (2010), Yetemwork Gebremeskel (2015), Dereje Shibru *et al.*(2016), Teklu Gebetadik and Dinku Negash (2016), Desalgn Begna *et al.*(2017) who identified agro-chemicals, poisonous as the first important constraint mentioned as reasons for hive productivity and colony population decline in different parts of the country.

Table 7. Major crop production and livestock production problem

Major crop constraint	Farming category	
	Beekeepers (%)	Non beekeepers (%)
Shortage of farm land	38 (47.5)	23 (28.75)
Drought	15 (18.75)	17 (21.25)
Shortage of oxen	10 (12.5)	9 (11.25)
Soil fertility	3 (3.75)	8 (10)
Input	2 (2.5)	9 (11.25)
Weed	5 (6.25)	5 (6.25)
Pest	4 (5)	1 (1.2)
Disease	1 (1.25)	3 (3.75)
Rodent	2 (2.5)	5 (6.25)
Livestock production constraints		
Shortage of animal feeds	35 (43.75)	43 (53.75)
Shortage of grazing land	29 (25)	25 (31.25)
Diseases	11(13.75)	9 (11.25)
Shortage of improved breeds	5 (6.25)	3 (3.75)

Table 8. Major beekeeping problems in the study areas

Constraints	Respondents rank (n=80)							Percentage	Rank
	1 st	2 nd	3 rd	4 th	5 th	6 th	7 th		
Agro-chemicals	9	5	21	5	13	15	18	15.63%	1 st
Poisonous plants	11	23	16	23	7	2	2	15.27%	2 nd
Absconding	5	9	13	14	18	11	10	14.54%	3 rd
Increase production cost	6	10	4	12	10	14	23	14.36%	4 th
Shortage of bee forage	9	20	11	21	4	7	5	14%	5 th
Drought	38	13	15	4	3	1	0	13.45%	6 th
Pest and predators	2	0	0	1	15	30	22	12.7%	7 th

4.4. Honeybee plant species and their availability

4.4.1. Honeybee flora trends in the study area

Majority of the respondents (90%) confirmed that honeybee forage is decreasing from time to time (Fig 5). Whereas, 6.25% and 3.75% of the respondents explained that honeybee forage availability did not change and increased in the last years respectively. According to the respondents, agrochemicals application and deforestation were undesirable effects on honeybee forages and the main reason for the decline of honeybee forage. This result is in line with findings of (Amssalu Bezabeh *et al.*, 2012; Sintayehu Fetene 2016) who showed that decrease in honeybee forage resources is recognized to be due to deforestation and agrochemical application for various purposes. In general, honeybees are facing shortage of forages associated with seasonal variation and resulted in low production and productivity.




Figure 5. Honeybee forage trends in the study areas

4.4.2. Major crops/horticultural plants used as honeybee forage

During the survey work, respondents have identified that coffee (*Coffea arabica*), avocado (*Persea americana*), papaya (*Carica papaya*), mango (*Mangifera indica*), soya bean (*Glycine max*) and maize (*Zea mays*) are the major Honeybee floral resource plants in the study area (Table 9). However, it has been explained also that honeybees are exposed to agro-chemicals poisoning, as these plants are dependent on chemical application. This finding is in line with the general facts that considerable numbers of honeybees are killed when bee forage plants are sprayed with agro-chemicals during flowering (Mark and Libby, 2003; US EPA, 2017). Though availability of supplementing seasonal bee forages could result in higher honey production, there was limited number of bee forage plants during times of dearth. Moreover, bee forage development and improvement, in the study area, except in watersheds/closure areas, is too minimal. The followings are the major bee plants available in the study area with flowering periods as indicated by respondents.

Table 9. The main bee floras crops of the study area and their flowering calendar

Local name (In Sidamigna)	Common Name	Scientific Names	Flowering periods	Number of days on flowering	Potential source (pollen/nectar)
<i>Buna</i>	Buna	<i>Coffee Arabica</i>	January-February	Ten days	Nectar
<i>Abukato</i>	Avocado	<i>Persea America</i>	Sept-Dec	Twenty five days	Pollen and nectar
<i>Mango</i>	Mango	<i>Mangifra indica</i>	Sept-Dec	Fifteen days	Pollen
<i>Wahe</i>	Soy bean	<i>Glycine max</i>	September	Ten days	Pollen
<i>Denicha</i>	Dinich	<i>Solanum tuberosum</i>	August- Sept	Ten days	Pollen and nectar
<i>Badala</i>	Bokolo	<i>Zea mays</i>	Jun –August	Fifteen days	Pollen
<i>Baqela</i>	Bakela	<i>Vacia foba</i>	Sept-October	ten days	Pollen
<i>Timatie</i>	Timatim	<i>Lycopersicon esculentum</i>	year round	Five days	Pollen and nectar
<i>Atara</i>	Ater	<i>Pisum sativum</i>	Sept-October	Ten days	Pollen
<i>Lome</i>	Lomi	<i>Citrus limon</i>	Sept –October	Twenty days	Pollen and nectar
<i>Papaya</i>	Papay	<i>Carica papaya</i>	Sept-Nov	Twenty day	Pollen and nectar
<i>Chate</i>	chat	<i>Catha edulis</i>	Year round	Fifteen days	Pollen

4.4.3.1. Beekeepers knowledge on honeybee poisoning plants

In this study, 75 (93.75%) of the beekeeping respondents have explained that they do have awareness on honeybee poisoning plants while 5 (6.25%) of the respondents do not have information on honeybee poisonous plants (Table 10). This result indicated that majority of the beekeeping farmers had the knowledge on honeybee poisoning plants in the study area. However, all of the beekeeping farmers confirmed that they do not know how to differentiate the types of honey made from poisonous plants. Moreover, 87.5% and 97.5% of the respondents did not have information on the effects of poisonous plants on other animals in addition to honeybees and the importance of these plants in the local conditions respectively (Table 10). This implied that the majority of the beekeepers in the study area did not have any information on the general backgrounds of these poisoning plants other than their toxicity to honeybees.

Table 10. Farmer's awareness on honeybee poisoning plants

Respondent perception	Yes	No	unknown
Is there any poisonous honeybee floral species in your area?	75(93.75%)	5(6.25%)	-
Are there any consumers' preferences on the types of honey sourced from poisonous plants?	-	-	80(100%)
Do they have impacts on other animals; despite Honeybees and /or human beings?	10(12.5%)	70(87.5%)	-
Despite poisoning effects, do they have any other importance locally?	2(2.5%)	78(97.5%)	-

4.4.3.2. Major toxic plants that affect honeybees

In the study area, respondents have identified ten plants to be poisonous to honeybees. despite the fact that a laboratory based confirmation is required, among these plants *Lanthana camara*, *Euphorbia continifolia*, *Climatis flammula*, *Ranclus multifides*, *Discopodium penninervium*, *Climatisinte grifolia*, *Ddataura metal linn*, *Sesbania sesban*, *Phytolaca americana* and *Justitia schemperina* were the most frequently complained toxic plants in the study area (Table 11). This list of poisonous plants comparable with results suggested by different researcher. According to Yetimwork Gebremeskel *et al.*(2015) reported that plants like *Acacia saligna*, *Euphorbia species*, *Melia azedarach* and *Azadirachta indica* were identified as poisons in Kilte Awulalo district of eastern Tigray. Additionally, Hailegebriel Tesfaye (2014) and Abebe Mitike (2017) have reported that *Sesbania sesban* and *Justitia schemperina* were known to cause poisoning of honeybees in Ethiopia. Various

plants grow in different areas that have different climatic conditions. These factors may contribute to the chemical compositions of the plants, which account for existence of different toxic plants in different geographical areas.

The respondents revealed that honeybees show various symptoms when consuming the poisonous plants. Because of poisonous flowering plants, honeybees have increased their aggressiveness, have died tremendously in and around the hive entrance and lack of foraging force and paralysis. Moreover, this survey result revealed that flower and stem parts of poisonous plants are sources of toxic effects supported by 97.5% and 2.5% of respondents respectively (Table 13). However, leaves, seeds, and fruits of poisonous plants were not identified as a source of toxic substances to honeybees. On the other hand, Ketawa (2008) reported that whole parts of *Datura metal Linn* were toxic to livestock. Wise management or clearing of these identified poisonous plants during flowering and most active periods around apiaries and development of other potential non poisonous bee forage plants have been suggested as an option to prevent or minimize the side effects in the study area.

Table 11. Summary of complained honeybee poisoning plants in the study areas

Local name (In Sidamigna)	Scientific name	Flowering period	Symptom
Che'atyguma	<i>Lanthana camara</i>	Year round	Paralysis
Du'emo	<i>Euphorbia continifolia</i>	Early summer	increased defensiveness
Tolchoomy	<i>Climatis flammula</i>	January	Dead honeybees in front of the hives,
Beetibetoo	<i>Ranclus multifides</i>	August - February	Lack of foraging
Laalunity	<i>Discopdium penninervium</i>	January	Lack of foraging and Repellent
Fiidee	<i>Climatisinte grifolia</i>	January	Lack of foraging and Repellent
Booriborich	<i>Datura metal linn</i>	July - October	Increased defensiveness
Taasfanii	<i>Sesbania sesban</i>	June	Lack of foraging
Raafoo	<i>Phytolaca americana</i>	December–February	Dead honeybees in front of the hives
Seensele	<i>Justitia schemperina</i>	November	Increased defensiveness

Table 12. Summary of plants that are poisonous with their botanical parts of exposure

Flower	Steam
<i>Lanthana camara</i>	<i>Climatis flammula</i>
<i>Euphorbia continifolia</i>	
<i>Climatis flammula</i>	
<i>Ranclus multifides</i>	
<i>Discopdium penninervium</i>	
<i>Climatisinte grifolia</i>	
<i>Datura metal linn</i>	
<i>Sesbania sesban</i>	
<i>Phytolaca Americana</i>	
<i>Justitia schemperina</i>	

Level of abundance for honeybee poisoning plants in the study area were vary; about 50 (62.5%), 40 (40%), 30 (37.5%) and 29 (36.25%) of the respondents revealed that *Euphorbia continifolia*, *Ranclus multifides*, *Datura metal linn*, and *Justitia schemperina* are more abundant toxic plants in the study area respectively. While the other toxic plants were abundant and rarely abundant in the area




Figure 6. Level of abundance for different poison plants in the study area

4.5. Respondents' practice in Agro-chemicals application

4.5.1. Purpose of Agro-chemicals application

According to the respondents, 93.75%, 78.12% and 59.37% of them were using agro-chemicals for the control of pests, for the control of weeds in their farmland and for the control of fungi/rust respectively. According to the result, majority of the respondents were using agro-chemicals for control of crop pests (Fig 7). This results agrees with Kalayou Hiluf and Amare Ayalew (2016) who have reported that about 93.26%, 89.93% and 37.5% chemical application in Amhara region was for the control of weeds, pests and fungi respectively.




Figure 7. Purpose of Agro-chemicals application

4.5.2. Respondents' utilization of Agro-chemicals

The result indicates that 75% of beekeepers and 87.5% of non-beekeepers had used agro-chemicals in the study area respectively (Table 13). It was expected that beekeepers had more information on toxicity of agro-chemicals to honeybees than non-beekeepers, our statistical analysis confirmed that there is no significant difference between beekeepers and non-beekeepers in agro-chemicals utilization at $p < 0.05$ ($X^2 = 0.068$). Surprisingly, it was found that beekeepers aware of the toxic effects of agro-chemicals on honeybees due to various reasons being household labor shortage and higher prevalence of pests and parasites were the major ones. In fact that, enables the adult labor to engage in income generating on off farm activities and children for schooling. Actually, this result is in line with Fikre Lemessa *et al.* (2016) who stated that the use of agro-chemicals were an assistance to eliminate labor cost and adult engaged to other off farm activities instead of labor intensive control of crop weeds, pests and diseases.

Table 13. Farmer's perception of agrochemical utilizations

Respondents category	Do you use agro-chemicals?		X ² (p-value)
	Yes	No	
Beekeepers	60(75%)	20(25%)	0.068
Non beekeepers	70(87.5%)	10(12.5%)	
Total	130(81.2%)	30(18.8%)	

4.5.3. Types and amount of agro-chemicals used by respondents

The types of agro-chemicals used by the respondents in the study area were Malathion 50%, Agrothoate 40%, 2, 4-D, Diazinon 60%, Pyriban 48%, Mancozeb 80% and Pallas 45 OD. According to the data collected from respondents, 87.5% and 97.5% of the interviewed beekeepers and non beekeepers respectively were using 2, 4-D for the control of weeds. Whereas 77.5% and 86.2% of beekeeping and non-beekeeping respondents respectively were using Agrothoate 40% for the control of Ecto-parasite and different pests while 35% and 22.5% of beekeeping and non-beekeeping respondents were Diazinon 60% used for different purpose respectively and Pallas 45 OD (57.5% and 61.2% of the beekeeping and non-beekeeping respondents) respectively were the least used agro-chemicals in the study area (Table 14). This result corresponds to the findings of Guesh Godifey (2015) and Dawit Melisie *et al.* (2016) who indicated that 2, 4-D, Agrothoate 40%, Malathion 50% and Mancozeb 80% were the major agro-chemicals used in different parts of Ethiopia. Moreover, statistical analysis did not show a significant difference between the beekeeping and non-beekeeping respondents on the utilization of Agrothoate 40% at $p < 0.05$ ($X^2 = 0.128$). This is because of the reason that majority of the respondents in the study area are *chat* growers thereby they are using this agrochemical for the control of *chat* pest. However, the use of other agro-chemicals by respondents did show a significant difference among the beekeeping and non-beekeeping respondents.

Table 14. Major types of agro-chemicals used by the respondents

Types of agro-chemicals	Agro-chemicals utilization				X ² (p-value)
	Beekeepers		Non beekeepers		
	N	%	N	%	
2,4-D	70	87.5	78	97.5	0.036
Agrothoate 40%	62	77.5	69	86.2	0.128
Malathion 50%	58	72.5	76	95	0.000
Pyriban 48%	51	63.8	65	81.2	0.007
Mancozeb 80%	31	38.8	51	63.8	0.003
Diazinon 60%	28	35	46	57.5	0.021
Pallas 45 OD	18	22.5	49	61.2	0.000

N= number of respondents and X²=chi square

More specifically, 92.5% of the respondents used 2,4-D with an average dose of 1.11 ± 0.016 liters per hectare (lit/ha) for weed control of *teff* and *wheat*. Agrothoate 40%, Pyriban 48% and Diazinon 60% were used at the rate of 1.16 ± 0.02 lit/ha, 1.29 ± 0.029 lit/ha, 1.07 ± 0.014 lit/ha respectively for the control of Aphides, Termite and African ball worms on *Chat*, *tomato*, *paper*, *cabbage* and *maize*. Similarly, respondents were using 1.2 ± 0.022 lit/ha of Malathion 50% for the control of *Chat*, *tomato*, *onion* and *cabbage* pests and diseases. Mancozeb 80% at the rate of 1.19 ± 0.023 lit/ha was used to treat late blight and powdery mildew of *wheat*, *tomato* and *potato*. Further, Pallas 45 OD at the rate of 1.05 ± 0.012 lit/ha was used by the respondents for the control of *teff* and wheat grass weeds (Table 15). However, it has been observed that all the rates used during the application of agro-chemicals for the control of crop weeds, pests and diseases were out of the recommended doses for each of the chemicals. In this case, it is clear that how much side effects that they could cause to non-target organisms including honeybees and our environment. Moreover, application of agro-chemicals below the recommended dose may lead to the development of resistant crop pest and disease strains and the chemical may not be as effective as expected to control the target. Our result is also with the same scenario with the findings of Dawit Melisie *et al.* (2016), who have reported that most farmers in Adami Tulu district of Oromia region were applying insecticides out of the recommended rate.

Table 15 Types and amount of agro-chemicals used for various field crops

Type of agro-chemicals	Frequency (%)	Mean ±SE	Type of crops	Used for the control of:
2,4-D	148(92.5%)	1.11±0.016	Wheat and Teff	Broad leaf weeds
Agrothoate 40%	131(81.87%)	1.16±0.021	Chat, tomato, paper and cabbage and maize	Aphides and African ball worms
Malathion 50%	134(83.5%)	1.2±0.022	Chat, tomato, onion and cabbage	Any worms
Pyriban 48%	116(72.5%)	1.29±0.029	onion and cabbage	Bollworms and insects
Mancozeb 80%	82(51.25%)	1.19±0.023	Wheat, Tomato and potato	Late blight, leaf spot and Downey mildew
Diazinon 60%	74(46.25%)	1.07±0.014	Teffe and maize	Termite and ball worms
Pallas 45 OD	67(41.87%)	1.05±0.012	Wheat and Teff	Any grass species

4.5.4. Trends of agrochemical use in the study area

Respondents pointed out that application of agro-chemicals has increased through time. Accordingly, the data revealed that application of 2,4-D, Agrothoate 40%, Malathion 50%, Pyriban 48%, Mancozeb 80%, Diazinon 60% and Pallas 45 OD has increased through years 2013 to 2017 (Fig.6). This implied that the use of agro-chemicals in the study area has become a common practice among the beekeeping and non-beekeeping respondents. This was supported by established good linear regression between number of respondents using agro-chemicals and period at $R^2 = 0.834$ (Fig 8). This relationship has also revealed the presence of positive and significant relationship between engagement in off farm activities and intensity of agrochemical use in the study area. This result was also in line with suggestions summarized by Jorge *et al.* (2009) who have explained that nominal expenditures on agro-chemicals increased steadily for most of the last half century worldwide. However, intensity of agrochemical use has declined during year 2015 due to low pest prevalence in the area in which occurrence of Eline was pinpointed as a major reason.




Figure 8. Trend of agrochemical application from the year 2013 to 2017

4.5.5. Sources of agro-chemicals

All the respondents were found to purchase agro-chemicals from various sources. Accordingly, 68.8% and 31.3% of the respondents confirmed that cooperatives and legal traders were potential sources of 2,4-D respectively. Similarly, Agrothoate 40%, Malathion 50%, Pyriban 48%, Mancozeb 80%, Diazinon 60% and Pallas 45 OD were found to be sourced from cooperatives and legal traders (Fig 9). However, it has been revealed that 15% of the respondents used Malathion 50% obtained from illegal traders. The major reason behind the illegal distribution Malathion 50% has been found that there are few numbers of licensed traders and cooperatives that are inaccessible to most of the agrochemical users in the study area. It has been also indicated that agro-chemicals are found illegally from open markets, shops, veterinary pharmacies and retailing farmers. This result is also in line with the findings of Kalayou Hiluf and Amare Ayalew (2015) and Fikre Lemessa (2016) who showed that most farmers are purchasing their agro-chemicals from open markets and unlicensed vendors or individuals. Moreover, purchasing of agro-chemicals from illegal sources could not ensure the user that agro-

chemicals are genuine products, effective against targets and to raise any compliant on the seller if something undesirable happened upon use. At this point, it is remarkable to note that respondents did not worry about side effects of agrochemical use rather their accessibilities in the study area.




Figure 9. Source of agro-chemicals used by the respondents

4.5.6. Time of agrochemical application

Agrochemical application is the practical delivery of different types of agro-chemicals and insecticides to their biological target groups. Efficiency and affectivity of agrochemical application in the field to properly control target biological groups highly depends on appropriate timing following chemical specific instructions which otherwise inflict undesirable damage on non-target organisms like honeybees and the general environment. According to the result of this survey, 80.6% and 95.6% of the respondents were found to apply 2,4-D at midday (Fig 10). Whereas, Agrothoate40%, Malathion 50%, Pyriban 48%, Mancozeb 80%, Diazinon 60% and Pallas 45 OD were applied to crops at any time of the day whenever 73.1%, 59.4%, 76.9%, 46.9%, 12.5% and 45% of the respondents did get time to apply and based on the interest of applicators (Fig 10). In this condition, it is very easy to understand that honeybees will receive bigger chance to be poisoned due to irregular time of applications practiced by respondents. This result is partially in line with findings of Sintayehu Fetene (2016) who indicated that farmers apply Agro-chemicals at different time of the day and overlapped with the active foraging time of the honeybees.




Figure 10. Timing of various agrochemical application

4.5.7. Respondents' awareness on agro-chemicals utilization and handling management

Farmers are serving as the main unit of agrochemical application. Hence, the degree of awareness on agrochemical utilization inherently affects their methods of application which could generate considerable

amount of agrochemical use ultimately to influence the health of honeybees. In this study, all the respondents were aware of the fact that agro-chemicals are useful not only for the control of crop pests but also causes environmental pollution. Furthermore, the data confirmed that 78.1% of the respondents do consider the fact that agro-chemicals could kill non-target organisms like honeybees. They further declared that agro-chemicals could directly and indirectly affect the wellbeing of the honeybees. Similarly, different authors reported that the decline of honeybees due to agrochemical effects on honeybees and plants that have useful function in the ecosystem in general and forage plants for bee in particular (Tadesse Amara and Asferachew Abate, 2008; Asamnew Tassew and Maria, 2016). This clearly indicates that even though, the farmers are aware of adverse effects of the agro-chemicals on the honeybees, they continued to use agro-chemicals without taking care of the side effects to minimize the death of honeybees. As regards to storage and handling of agro-chemicals before and after use, 90.62% of the respondents confirmed that agrochemical containers are hanged inside their homes and only 6.25% of them are keeping agro-chemicals safely out of reach. Shockingly, 3.12% of the respondents are keeping the agro-chemicals with their foodstuffs irregularly at home. Even though proper storage or disposal of empty agrochemical is one of the most important component in the safe use of agro-chemicals, 76.25%, 11.25% and 9.37% of the respondents confirmed that they are using the chemicals even if they are not needed, used them for unintended application and they dispose them anywhere on the ground respectively (Table 16). It is clear that this practice is not proper in the management of agro-chemicals and their ruminants as it could increase humans, animals and the environment exposure to the undesired effects of agro-chemicals. In general, we found that this result is in consistent with findings of Govinda (2014), Fikre Lemessa *et al.* (2016) and Hailay Abrha *et al.* (2016). However, it is wise to suggest that proper management of agro-chemicals before and after application could increase efficiency of treatments, quality life to humans and the environment, and decrease chance of non-target organisms to be exposed and affected by toxic elements.

Regarding to the management of empty agrochemical containers, we have found that 95%, 3.25% and 1.25% of the respondents indicated that they are using them for household purposes, left them anywhere in the farmland they burn them out respectively (Fig 17). However, it was confirmed that no one has buried empty agrochemical containers with their leftovers properly as instructed. On the other hand, the survey data revealed that most respondents (95.62%) did not know the use of protection and did not use protective clothes during agrochemical application (Table 16).

Furthermore; about 80% of the respondents never get trainings on how to use and importance of protection during application. In this regard, key informants of this study also witnessed that they have never received any instruction from either agrochemical suppliers or extension people on the how to use and why to use protective clothes during application. Consequently, it was, thus, confirmed that almost all (97.5%) of the respondents have never followed instructions during agrochemical application.

Table 16. Perception of farmers towards agro-chemicals and their sustainable use

Variables	N	Percentage
Adverse effects of agro-chemicals		
➤ Environment pollution	160	100
➤ Killing non-target species	125	78.1
Agro-chemicals storage practice		
➤ Hanging inside the house	145	90.62
➤ Locked up in safe place	5	3.12
➤ Along with food stuff	10	6.25
Left over Agro-chemicals solution		
➤ Stored and used for another application	18	11.25
➤ Pour in to bushes, rivers/stream	3	1.87
➤ Sell it to other farmers	2	1.25
➤ Apply even though it is not needed	122	76.25
➤ Disposed on the soil	15	9.37
Disposal of empty Agro-chemicals containers		
➤ Use them for house hold purpose	152	95
➤ Buried	-	-
➤ Left in the farm	6	3.75
➤ Burnt	2	1.25

Variables	N	Percentage
Do you use protective clothes while spraying agro-chemicals		
➤ Yes	7	4.38
➤ No	153	95.62
Health impacts		
➤ yes	147	91.9
➤ No	13	8.1
Training on use of Agro-chemicals		
➤ Yes	32	20
➤ No	128	80
Following the labeled instructions of the Agro-chemicals		
➤ Yes	4	2.25
➤ No	156	97.5

Regarding undesirable effects of agro-chemicals on honeybees and their produce, all respondents (80) have been affected negatively and have lost a total of 1556kg of honey with an average loss of 19.45 ± 4.0 kg honey per beekeeper per year due to improper application of agro-chemicals (Table18). Consequently, analysis of financial loss incurred indicated that respondents have lost 93360 ETB with an average of 1167 ± 245.42 ETB per beekeeper.

Table 18. Financial loss incurred due to unwise use of agro-chemicals in the study area

Factors	Number of respondents	Sum	Mean	Std. Deviation
Honey loss	80	1556	19.45	4.09
financial loss(ETB)	80	93360	1167	245.42

5: CONCLUSION AND RECOMMENDATION

In conclusion, the results of the present study show that poisons plants and chemical toxicities were among key causes of honeybee health problems in Sidama zone, SNNPR, Ethiopia. A total of ten plants were reported as having poisoning effect on honeybees by beekeepers in the study area. Among these plants *Lanthana camara*, *Euphorbia continifolia*, *Climatis flammula*, *Ranclus multifides*, *Discopodium penninervium*, *Climatisinte grifolia*, *Datura metal linn*, *Sesbania sesban*, *Phytolaca americana* and *Justitia schemperina* were the most commonly incriminated toxic plants. Similarly, agro-chemicals are considered as a powerful weapon or magic bullets in the study area in order to enhance the agriculture productivity. The utilization of agro-chemicals is increasing from time to time and mainly used to control weeds, pests, and diseases of crops and animals. Common agro-chemicals use in sidama zone was 2,4-D, Agrotheote40%, Malathion80%, Pyriban48%, Diazinone60%, Macozeb80% and Pallas 45 OD. It was expected that beekeepers had more information on toxicity of agro-chemicals to honeybees than non-beekeepers, but, there is no difference between beekeepers and non-beekeepers in agro-chemicals utilization in the study area. These agro-chemicals were illegal traded and available in open market, in shops along with food and other consumable items and veterinary drugs. Non beekeepers and beekeepers had no or little access to advices on how to use and handle the agro-chemicals. They stored left over agro-chemicals and empty agro-chemical containers with consumable items at home. Farmers were not following the instructions for application of agro-chemicals and they have been seen using over or under doses for application at any time they wanted. As the result of the improper use of agro-chemicals, honeybees were adversely affected. Decline of honeybee colonies and honey production have been registered due to various reasons of which agro-chemicals top in the listed.

According to the result of this study some of the suggested issues that require consideration by beekeepers and any development organizations are high lightened below:

- Creating awareness of the season of most poisonous plants are growing and cause problem on bee health.
- Experimental studies should be carried out to confirm the empirical knowledge of plant poisoning on honeybee. .
- Advise farmers to avoid the application of bee toxic agro-chemicals on blooming plants and it is a good idea to check for the presence of other blooming plants which might attract bees.
- A clear mechanism of working and chain of communication among institutions is very important to

minimize the prevailing risk imposed by improper use of agro-chemicals.

- The integrated efforts are very important to educate farmers on proper agro-chemical handling, management, utilization, appropriate safety precautions, effects of pesticide on honeybee health and Integrated Pest Management (IPM)

REFERENCES

- Abera Dereje. 2015. Survey of Toxic Plants in Livestock at Horro Guduru Wollega Zone, Journal of Biology, Agriculture and Healthcare, vol.5: pp.12-33.
- Addis Getu and Maleda Birhane. 2014. Chemical analysis of honey and major honey production challenges in and around Gonder, Ethiopia. *Aca. J. Nut.* 3(1)
- Alemtsehay Teklebirhan. 2011. Seasonal availability of common bee flora in relation to land use and colony performance in gegera watershed Atsbi wenberta district, Eastern zone of Tigray region, Ethiopia. pp.1-
- Asaminew Tassew and Maria W. 2016. Beekeeping dynamics and its driving forces of change in Ethiopian mixed farming system. Department of Animal Production and technology, College of Agriculture and Environmental Sciences, Bahir Dar University, Ethiopian.28: no.12
- Assemu Tesfa, Kerealem Ejigu and Adebabay Kebede. 2013. Assessment of current beekeeping management practice and honey bee floras of Western Amhara, Ethiopia. *Inter J Agri Biosci*, 2: 196-201.
- Atkins E.L., Greywood E.A. and Macdonald R. L. 1973. Toxicity of agro-chemicals and other agricultural chemicals on honeybees. University of California, Division of Agricultural Sciences *Leaflet.2287*.
- Atkins E.L., Kellum D. and Neuman K.J. 1975. Toxicity of agro-chemicals to honeybees. University of California, Division of Agricultural Sciences *Leaflet.2286*
- Atsbaha Hailemariam, Taye Tolemariam and Kebede Debele. 2015. Assessment of honey production system, constraints and opportunities in three selected Woredas of Tigray, *Basic Research Journal*, 4: 304–315
- Awraaris Getachew, Yemisirach Getachew, Dejen Assefa, Nuru Adgaba, Gebeyehu Ganga and Workneh Abebe. 2012. Honey production systems (*Apis mellifera* L.) in Kaffa, Sheka and Bench-Maji zones of Ethiopia. *Journal of Agricultural Extension and Rural Development*, 4(19): 528-541.
- Ayalew Kassaye. 2001. The honey bees (*Apis mellifera*) of Ethiopia. A morphometric study. M.Sc. Thesis, Agricultural University of Norway, ÅS, Norway.
- Bekele Tesfay. 2015. Beekeeping practices, factors affecting production, quality of honey and beeswax in bale zone, Oromia region. *African Journal of Rural Development*. 2 (2): 88
- Belets Gebremichael and Berhanu Gebremedhin. 2014. Adoption of improved box hive technology: Analysis of smallholder farmers in Northern Ethiopia. *International Journal of Agricultural Economics and Extension*, 2 (2): 077-082.
- Biruk Deribe, 2014. The Constraints of Honey Production in Beekeeping Cooperatives Case Study of Killite-Awlalo District, MSc Thesis presented to Mekelle University: Ethiopia.
- Bradbear N. 2003. Beekeeping and Sustainable Livelihood. Diversification booklet 1.FAO,65
- Bradbear N. 2009. Non-wood forest products: Bees and their role in forest livelihoods. A guide to the services provided by bees and the sustainable harvesting, processing and marketing of their products. *Agriculture, Forestry and Fisheries*. 3(1): 52-57
- Challa Kinate. 2010. Honey production, Marketing and Quality assessment in Gomma woreda, Southwestern, Ethiopia. *Basic Research Journal of Agricultural Science and* 4(10) : 304-315
- Chaubal P.D. and Deodikar G. B. 2017. Pollen Grains of Poisonous Plants, (August). doi: 10.1080/00173136309429114.
- Cornman R.S., Tarpay D.R., Chen Y., Jeffreys L., Lopez D and Pettis J.S. 2012. Pathogen webs in collapsing honeybee colonies. *PLoS One*. 7: 435- 462
- Craig I, Woods N, Dorr G.1998. A simple guide to predicting aircraft spray drift. *Crop Protection*; 17: 475–482.
- CSA (Central Statistical Agency of Ethiopia).2015. Agricultural Sample Survey report on livestock and livestock characteristics (private peasant holdings).Volume II, statistical bulletin 570, Addis Ababa, Ethiopia.
- CSA (Central Statistical Agency).2016. The Federal Democratic Republic of Ethiopia Central Statistical Agency Agricultural Sample Survey. Report on Crop Yield for Major Grain Crops. Private Small holder Farms, Main Season pp.1-16.
- Dawit Melisie, Tebkew Damte and Ashok K. 2016. Farmers' insecticide use practice and its effect on honeybees (*Apis mellifera*) foraging on onion flower in Adami Tullu district. *Glob. J. Pests Dis. Crop Prot.* 4: 139-145.
- Dereje Shibru, Getahun Asebe and Emanu Megersa. 2016. Identifying opportunities and constraints of beekeeping: the case of Gambella Zuria and Godere Weredas, Gambella Regional State, Ethiopia. *Entomol Ornithol Herpetol* 5: 2-6
- Desalegn Begna. 2015. Assessment of pesticides use and its economic impact on the apiculture subsector in selected districts of Amhara Region, Ethiopia, Holeta, Ethiopia. *Journal of Environmental Annual Toxicology*, 5: 267.

- Gezahegne Tadesse. 2001. *Beekeeping (in Amaharic)*, Mega Printer Enterprise, Addis Ababa, Ethiopia.
- Haftu Kebed and Gezu Tadesse. 2014. Survey on honey production system, challenges and opportunities in selected areas of Hadiya zone, Ethiopia, 6(6): 60-66
- Kerealem Ejigu, Tilahun Gebeyehu and Preston T.R. 2009. Constraints and prospects for apiculture research and development in Amhara region, Ethiopia. *Liv. Res. Rural Dev.* 21: no. 10.
- Kerealem Ejigu. 2005. Honeybee Production System, Opportunities and Challenges in Enebsesar Midir woreda (Amahara region) and Amaro special woreda (SNNPR), Ethiopia. M.Sc. Thesis, Alemaya University, Ethiopia.
- Kevan P.G. 2005. Pollinators as bio-indicators of the state of the environments: species, activity and diversity. *Agriculture, Ecosystems and Environment*
- Malede Birhan. 2013. Livestock Resource Potential and Constraints in Somali Regional State, Ethiopia. *Global Veterinaria*, 10(4): 432-438.
- MoARD. 2016. Livestock Development Master Plan Study. Phase I Report – Data Collection and Analysis, Volume N Apiculture. Addis Ababa, Ethiopia.
- Qaiser T., Ali M., Taj S. and Akmal N. 2013. Impact Assessment of Beekeeping in Sustainable Rural Livelihood. *Journal of Social Sciences*, 2(2), 82–90. Retrieved from [http://centreforexcellence.net/J/JSS/Vol2/No2/JSS2\(1\)Apr 2013.htm](http://centreforexcellence.net/J/JSS/Vol2/No2/JSS2(1)Apr 2013.htm)
- Tadesse Amare and Assferachew Abate. 2008. An assessment of the pesticide use, practice and hazards in the Ethiopian rift valley. 13
- Teklu Gebretsadik and Dinku Negash. 2016. Honeybee Production System, Challenges and Opportunities in Selected Districts of Gedeo Zone, Southern Nation, Nationalities and Peoples Regional State, Ethiopia. *International journal of research*, 4:12
- Tewodros Alemu, Eyassu Seifu and Amessalu Bezabih. 2015. Postharvest handling, opportunities and constraints to honey production in northern Ethiopia. *Liv. Res. Rural Dev.* 27 no. 5.
- Tewodros Alemu, Eyassu Seifu and Amessalu Bezabih. 2017. Honeybee production practices in Sekota district, northern Ethiopia, *African Journal of Rural Development*. 2:263-275
- Wolaye Kiros and Teklberhan Tsegay. 2017. Honey-bee production practices and hive technology preferences in Jimma and Illubabor Zone of Oromiya Regional State, Ethiopia. 9:31-43