

On Quasi-Similarity and Putnam- Fuglede Property of Operators In Hilbert Spaces

S.W. LUKETERO¹, P.G. RUGIRI^{1,2*} AND J.M. KHALAGAI³

SCHOOL OF MATHEMATICS, UNIVERSITY OF NAIROBI, P.O BOX 30197, 00100 NAIROBI KENYA

*Email of the corresponding author; githarapeterrugiri@yahoo.com

ABSTRACT

The problem of finding conditions under which two given operators say A and B have equal spectra has been considered by several authors. In this paper we show that quasi-similar operators A and B which satisfy the Putnam- Fuglede property have equal spectra and also equal essential spectra.

AMS subject classification: 47B47, 47A30, 47B20.

Key words and phrases: quasi-similarity, quasi-affinity and putnam-Fuglede property.

1.INTRODUCTION

Let H be a complex Hilbert space and $B(H)$ denote the Banach algebra of all bounded linear operators on H . An operators $A \in B(H)$ is said to be a quasiaffinity if A is both one-one and has dense range. The two operators A and B are said to be similar if there exist an invertible operator S such that $AS = SB$, while A and B are said to be quasi-similar if there exist quasi-affinities X and Y such that $AX = XB$ and $BY = YA$.

The concept of quasi-similarity and equality of spectra for a given pair of operators has been considered by a number of authors among them W.C. Clary [1] who showed that quasi-similar hyponormal operators have equal spectra. J.M. Khalagai and B. Nyamai [6] showed that if A and B quasi-similar operators with A dominant and B^* M-hyponormal then A and B have equal spectra. B.P.Dugga [4] showed that if $A_i, i = 1, 2$ are quasi-similar operators such that U_i is unitary in the polar decomposition $A_i = U_i |A_i|$ then A_1 and A_2 have equal spectra and also equal essential spectra. J.P Williams [8] and [9] showed that there are several cases which imply that two operators A and B have equal essential spectra under quasi-similarity. For example if A and B are both hyponormal or are both partial isometrics or are quasi-normal.

In this paper we prove results on equality of spectra and essential spectra for classes of operators that satisfy the Putnam-Fuglede property.

Theorem A (Putnam-Fuglede)

Let $A, B \in B(H)$ be normal operators. For any other operator X we have that $AX = XB$ implies $A^*X = XB^*$.

The following result by J.M. Khalagai and B.Nyamai [6] will be required.

Theorem B

Let A, B and X be operators such $AX=XB$ implies $A^*X = XB^*$. If X is either one-one or has dense range then A and B are normal operators.

The following result by R.G. Douglas [2] will also be required for the proofs of our results.

Theorem C

Let $A, B \in B(H)$ be quasi similar normal operators. Then A and B are unitarily equivalent and hence have equal spectra.

2.NOTATION AND TERMINOLOGY

Given an operator $A \in B(H)$ we denote the numerical range of A by $W(A)$.

Thus

$$W(A) = \{ \langle Ax, x \rangle : \|x\| = 1 \}$$

The spectrum of A is denoted by $\sigma(A)$. Thus $\sigma(A) = \{ \lambda \in \mathbb{C} : A - \lambda I \text{ is not invertible} \}$

Where \mathbb{C} is the complex number field. The commutator of A and B is denoted by $[A, B]$ where

$$[A, B] = AB - BA$$

An operator A is said to be dominant if to each $\lambda \in \mathbb{C}$ there corresponds a number $M_\lambda \geq 1$ such

$$\text{that } \|(A - \lambda)^*x\| \leq M_\lambda \|(A - \lambda)x\| \quad \forall x \in H$$

M-hypo-normal if $\exists M \geq M_\lambda$ for all λ in the definition of dominant operator .i.e $\|(A - \lambda)^*x\| \leq M \|(A - \lambda)x\| \quad \forall x \in H$

Hyponormal if $A^*A \geq AA^*$
 Quasinormal if $[A^*A, A] = 0$
 P-hypo-normal if for $0 < p \leq 1, (A^*A)^p \geq AA^*(A^*A)^{p-1}$
 Normal if $[A, A^*] = 0$
 Self adjoint if $A = A^*$
 Partial isometris if $A = AA^*A$
 Isometry if $A^*A = I$
 Unitary if $A^*A = AA^* = I$
 Fredholm if its range denoted by $\text{ran}A$ is closed and both null space, $\ker A$ and $\ker A^*$ are finite dimensional.
 The essential spectrum of A is denoted by $\sigma_e(A)$.
 Thus $\sigma_e(A) = \{ \lambda \in \mathbb{C} : A - \lambda I \text{ is not fredholm} \}$
 The following operator inclusions are proper.
 Normal \subset hyponormal \subset p-hypo-normal and Hypo-normal \subset M-hyponormal \subset Dominant.

3.RESULTS

Theorem 1

Let $A, B, X \in B(H)$ be operators such that:

$AX = XB$ where A and B satisfy Putnam –Fuglede property and X is a quasi-affinity.

Then we have:

- i. $\sigma(A) = \sigma(B)$
- ii. $\sigma(AA^*) = \sigma(BB^*)$
- iii. $\sigma(A^*A) = \sigma(B^*B)$

Proof

We note that $AX = XB$. (1) implies $A^*X = X^*B$. On taking adjoints we have
 $BX^* = X^*A$ (2).

Since X is a quasiaffinity it follows from (1) and (2) that A and B are quasi-similar. It now follows from both theorem B and C above that A and B are unitarily equivalent normal operators. Hence $\sigma(A) = \sigma(B)$.

Also using the equations $AX = XB, A^*X = X^*B, BX^* = X^*A$ and $B^*X^* = X^*A^*$ we get
 $A^*AX = A^*XB = XB^*B$ (3)

and

$$B^*BX^* = B^*X^*A = X^*A^*A \quad (4)$$

$$\text{Also } AA^*X = A X B^* = X B B^* \quad (5)$$

and

$$B B^* X^* = B X^* A^* = X^* A A^* \quad (6)$$

From (3) and (4) A^*A and B^*B are quasi similar positive operators. Hence $\sigma(A^*A) = \sigma(B^*B)$.

Also from (5) and (6) AA^* and BB^* are quasi-similar positive operators.

Hence $\sigma(AA^*) = \sigma(BB^*)$.

We note that from the results by J.P Williams [8] the following corollary is immediate.

Corollary 1

Let $A, B, X \in B(H)$ be operators such that $AX=XB$ where X is quasi-affinity then

$\sigma(A) = \sigma(B), \sigma(A^*A) = \sigma(B^*B)$ and $\sigma(AA^*) = \sigma(BB^*)$ under any one of the following conditions:

- i. A is dominant and B^* is m-hyponormal
- ii. A is dominant and B^* is p-hyponormal
- iii. A and B^* are p-hyponormal

Proof

All the classes of operators stated above are known to satisfy Putnam-Fuglede property.(see[3])

Hence results follow from theorem 1 above.

Corollary 2

Let $A, B, X \in B(H)$ be operators such that

$$AX=XB$$

where X is quasi-affinity then

$\sigma(A) = \sigma(B), \sigma(A^*A) = \sigma(B^*B)$ and $\sigma(AA^*) = \sigma(BB^*)$ under any one of the following conditions:

- i. A is dominant and B^* is m -hyponormal
- ii. A is dominant and B^* is p -hyponormal
- iii. A and B^* are p -hyponormal

Proof

We note that all the classes of operators stated above are known to satisfy Putnam-Fuglede property and exist quasi-affinities X and Y such that $AX=XB$ and $BY=YA$. We also note that in case of part,(iii),if in addition we have that in the polar decomposition $A=V|A|$ and $B=V|B|$.

U and V are unitary (see[4]) then $\sigma_e(A) = \sigma_e(B)$.

For an operator $B \in B(H)$ we say that B is consistent in invertibility (with respect to multiplication) or briefly that B is a CI operator if for each $A \in B(H)$, AB and BA are invertible or non-invertible together. Thus B is CI operator if $(\sigma(AB) = \sigma(BA))$. W.Gong and.Han [5]) proved among other results that an operator $B \in B(H)$ is CI operator iff $\sigma(B^*B) = \sigma(BB^*)$.

From this result the following corollary is immediate.

Corollary 3

Let $A, B, X \in B(H)$ be operators such that:

$AX = XB$ where A and B satisfy Putnam –Fuglede property and X is a quasi-affinity.

Then we have:

- i. $\sigma(A) = \sigma(B)$
- ii. $\sigma(AA^*) = \sigma(BB^*)$
- iii. $\sigma(A^*A) = \sigma(B^*B)$

Proof

In this case there exists two quasi-affinities say X and Y such that $AX=XB$ and $BY=YA$
 Thus the proof of theorem 1 can be traced to yield the result.

Corollary 4

Let $A, B, X \in B(H)$ be operators such that:

$AX = XB$ where A and B satisfy Putnam –Fuglede property and X is a quasi-affinity.

Then we have:

- i. $\sigma_e(A) = \sigma_e(B)$
- ii. $\sigma_e(AA^*) = \sigma_e(BB^*)$
- iii. $\sigma_e(A^*A) = \sigma_e(B^*B)$

Corollary 5

Let $A \in B(H)$ be quasi-similar operators which satisfy Putnam-Fuglede property. Then we have that B is a $C.I$ operator whenever A is and conversely.

Proof

We note that for such operators A and B .

$\sigma(AA^*) = \sigma(BB^*)$ and $\sigma(B^*B) = \sigma(A^*A)$.

Thus if A is CI operator then $\sigma(A^*A) = \sigma(AA^*) = \sigma(B^*B) = \sigma(BB^*)$. Hence A is a C.I operator.

In our next result we show that by imposing more stringent conditions on the intertwining we obtain equality of operators rather than equality of their spectra. In so doing we require the following result by I. H Sheth and J.M Khalegai [7]

Theorem D

Let $T, S, X \in B(H)$ be operators such that.

$TX = XS$ and $SX = XT$ with $O \notin W(X)$ where $S - T$ is normal.

Then $S = T$

We have the following result in this direction.

Theorem 2

Let $A, B, X \in B(H)$ be operators such that A and B satisfy Putnam-Fuglede property and $AX = XB$ with X self adjoint and $0 \notin W(X)$. then $A^*A = B^*B$ and $AA^* = BB^*$

Proof

We first note that $AX = XB \Rightarrow A^*X = XB^*$

Thus taking adjoints we also have $BX = XA$.

Thus using these equations we have: $A^*AX = A^*XB = XB^*B$ and

$$AA^*X = AXB^* = XBB^*$$

Now letting $T = A^*A$ and $S = B^*B$

we have that $S - T = B^*B - A^*A$ is normal.

Hence by theorem D above

$$B^*B = A^*A$$

Similarly if $T = AA^*$ and $S = BB^*$

then $BB^* - AA^*$ is normal.

Hence $BB^* = AA^*$

REMARKS

1. We note that $A = B$ implies $A^*A = B^*B$ and $AA^* = BB^*$

However the converse is not necessarily true.

2. We also note that in theorem 2 above we can replace the condition that A and B satisfy Putnam-Fuglede theorem by simply requiring that A and B are normal operators in view of theorem B and the fact that $0 \notin W(X)$.

REFERENCES

1. W.S. Clary, Equality of Spectra of quasisimilar hyponormal operators, Soc. 53(1975), 89-90.
2. R.G. Douglas, On the operator equation $S^*XT = X$ and related topics, Acta Sci.Maths (Szeged)30(1969),19-32.
3. B.P. Duggal, on dominant operators, Arch.Math. Vol46(1986),353-345.
4. _____, Quasi-similar p-hyponormal operators, Interger Equat Oper Th Vol 26(1996), 338-345
5. W. Gong and D. Han, Spectrum of products of operators and compact perturbations, Proc. Amer.Math Soc. Vol120 No3(1994),755-760.
6. J.M. Khalagai and B. Nyamai, on quasi-affinity and quasi-similarity of operators, Kenya J.Sci-Tech vol12No2 (1998), 14-17.
7. I.H. Sheth and J.M. Khalagai, on the operators equation $AH = KA$, Mathematics today VOL V/1987), 29-36.
8. J.P. Williams, Equality of essential spectra of quasi-similar quasinormal operators, J operator Theory 3(1980), 57-69.
9. _____, Quasi-similarity and hyponormal operators J operator Theory

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

