

# Comparative Analysis of Stratified Randomized Response Models for HIV Seroprevalence Surveys

Usman, A. & Abdulazeez, S.A.

Department of Mathematics, Statistics & Computer Science, Kaduna Polytechnic, Nigeria

## Abstract

Seroprevalence surveys of HIV pandemic are highly sensitive especially in Africa. The objective of this study is to reach research frontier to devise a two-way randomized response model (RRM) in stratification and use same to estimate HIV seroprevalence rates in a given population and compare results with the existing seroprevalence rates. The randomized response techniques (RRT) guarantees the anonymity of respondents in surveys aimed at determining the frequency of stigmatic, embarrassing or criminal behaviour where direct techniques for data collection may induce respondents to refuse to answer or give false responses. The motivation was to improve upon the existing RRM as well as to apply them to estimate HIV seroprevalence rates. Warner proposed the pioneering RRM for estimating the proportion of persons bearing a socially disapproved character. Quatember produced unified criteria for all RRTs, Kim and Warde proposed a stratified RRM and so many others. The proposed two-way RRM in stratification for HIV seroprevalence surveys was relatively more efficient than the Kim and Warde stratified estimator for a fixed sample size. The chosen design parameter was 0.7, using the criteria of Quatember who derived the statistical properties of the standardized estimator for general probability sampling and privacy protection. Furthermore, the model was used to estimate the HIV seroprevalence rate in a sampled population of adults 3,740 people aged 18 years and above attending a clinic in Kaduna, Nigeria using a sample size of 550. The findings revealed that HIV seroprevalence rate, as estimated by Model I, stood at 6.1% with a standard error of 0.0082 and a 95% confidence interval of [4.5%, 7.7%]. These results are consistent with that of Nigerian sentinel survey (2003) conducted by NACA, USAID and CDC which estimated the HIV seroprevalence in Kaduna State as 6.0%. Hence, the RRTs herein can serve as new viable methods for HIV seroprevalence surveys. Similarly, the result for model II show that, using the survey data, the model estimated the HIV seroprevalence rate as 8.74% with a standard error of 0.0134 and 95% confidence bands of [6.1%, 11.4%]. Accordingly, the sentinel projected seroprevalence rate, using the EPP Package, for the next ten years (2013) was 9.7%; very consistent with the 95% confidence interval. Hence, the RRTs herein can also serve as new viable methods for HIV seroprevalence surveys. Model II has a better chance of estimating HIV seroprevalence because it has higher privacy preservation.

## Key words

Randomized response techniques, stratified randomized response models, seroprevalence rates, design parameter, efficiency, sentinel surveys, stratified random sampling

## 1. Introduction

When sensitive topics are studied, respondents often react in ways that negatively affect the validity of the data. Such a threat to the validity of the results is the respondents' tendency to give socially desirable answers to avoid social embarrassment and to project a positive self-image (Rasinski, 1999). Warner (1965) reasoned that the reluctance of the respondents to reveal sensitive or probably harmful information would diminish when respondents could be convinced that their anonymity was guaranteed. Hence, Warner (1965) designed the first randomized response model (RRM). The crux of his method and all other RRTs that followed is that the meaning of the respondents' answers is hidden by a deliberate contamination of the data collection settings (Lee, 1993).

Studies with RRTs have been conducted in the areas of healthcare (Volicer & Volicer, 1982), on alcohol, drug abuse and sexual behaviour (Jarman, 1997), on child molestation (Fox and Tracy, 1986), on tax evasion (Houston & Tran, 2008), among others. Meta-analysis on 42 comparative studies showed that RRTs resulted in more valid population estimates than direct question-answer techniques (Lensvelt-Mulders et al., 2005). An advantage of using RRT when conducting sensitive research is that, the individual 'yes'-answer becomes meaningless as it is only a 'yes-answer' to the random device (Van der Hout, et al., 2002). The randomized response design is more effective than the direct question-answer design (Lensvelt-Mulders et al., 2005). The loss of efficiency in RR designs could be compensated when the results prove to be more valid (Kuk, 1990). When the loss in efficiency can be kept as small as possible the use of a RR design to study sensitive questions will become more profitable.

## 2. Methodology

In order apply the Straffied RRM; a study was conducted in Gwamna Awan General Hospital, Kaduna, Nigeria in November, 2011. With a carefully coordinated field work and sampling design on a population of 3,740

adults aged 18 years and above attending the Hospital using a sample size of 550 for each model. Furthermore, each model was used to estimate the HIV seroprevalence rate in the same population. Quatember (2009) both theoretically and empirically analyzed the effect of different design parameters on the performance of RRTs using different levels of privacy protection. Quatember (2009) suggested that 0.7 approximately works well for most RRM where the questions are regarded as highly sensitive. Hence, 0.7 is the chosen design parameter and deck of 50 cards as our random device throughout.

### 2.1 The Proposed HIV Seroprevalence Model I

The proposed HIV seroprevalence surveys Model I requires that a sample respondent in stratum  $h$  to answer an innocuous direct question and asked to use the random device  $R_{h1}$  if his/her answer to direct question is “yes”. If answer to the direct question is “no”, he/she is requested to use another random device  $R_{h2}$  twice. Both random devices  $R_{h1}$  and  $R_{h2}$  consist of two statements (i) “I am HIV positive” and (ii) “I am HIV negative”, presented with probabilities  $P_{h1}$  and  $(1 - P_{h1})$  respectively. Here the random device  $R_{h2}$  would to be answered twice. Hence, we can obtain the estimator of population proportion  $\pi_h$  in  $h$ th stratum based on the responses from  $R_{h1}$  as follows. The probability of a ‘yes’ response from the respondents using  $R_{h1}$  is given by:

$$\lambda_{h1} = P_{h1}\pi_h^* + (1 - P_{h1})\pi_{hy} = P_{h1}\pi_h^* + (1 - P_{h1}) \quad (1)$$

Also, the probability of a ‘no’ response from the respondents using  $R_{h1}$  is given by:

$$\lambda'_{h1} = P_{h1}(1 - \pi_h^*) + (1 - P_{h1})(1 - \pi_{hy}) = P_{h1}(1 - \pi_h^*) \quad (2)$$

Since the respondent using  $R_{h1}$  has already answered yes to the direct question,  $\pi_{hy} = 1$ .

Among those that answered ‘yes’ to the innocuous questions in stratum  $h$ ; suppose that  $n_{h1}$  report ‘yes’ and  $(n_h - n_{h1})$  report ‘no’, the likelihood of the sample in the same stratum is given below:

$$\xi = [P_{h1}\pi_h^* + (1 - P_{h1})]^{n_{h1}} \times [P_{h1}(1 - \pi_h^*)]^{n_h - n_{h1}} \quad (3)$$

We obtain the maximum likelihood estimate (MLE) of  $\pi_h^*$  as follows:

$$\therefore \pi_h^* = \frac{n_h P_{h1} - n_{h1} + n_{h1}}{n_h P_{h1}} \quad (4)$$

Hence, the unbiased estimators in terms of the responses of the respondents using  $R_{h1}$  is given by:

$$\hat{\pi}_{h1} = \frac{\hat{\lambda}_{h1} - (1 - P_{h1})}{P_{h1}} \quad (5)$$

Where; the proportion of ‘yes’ answers from  $R_{h1}$  in the sample is given as;

$$\hat{\lambda}_{h1} = \frac{n_{h1}}{n_h}$$

The variance of is obtained as follows:

$$\begin{aligned} \text{Var}(\hat{\pi}_{h1}) &= \left[ \frac{1}{P_{h1}} \right]^2 \text{Var}(\hat{\lambda}_{h1}) \\ &= \left[ \frac{1}{P_{h1}} \right]^2 \left( \frac{\hat{\lambda}_{h1}(1 - \hat{\lambda}_{h1})}{n_{h1}} \right) \\ \therefore \text{Var}(\hat{\pi}_{h1}) &= \frac{(1 - \pi_{h1})(P_{h1}\pi_{h1} + 1 - P_{h1})}{n_{h1}P_{h1}} \end{aligned} \quad (6)$$

The respondent, in  $h$ th stratum, giving a “no” answer to the question are to use  $R_{h2}$  twice to report two answers, where  $R_{h2}$  consists of the two statement of Warner’s RR method. To have the first response reported the probabilities of the two statements are  $P_{h2}$  and  $(1 - P_{h2})$  whereas to get the second response from the responses

these probabilities are  $P_{h2}^*$  and  $(1 - P_{h2}^*)$ . Two unbiased estimators based on the two set of responses from respondents using  $R_{h2}$  can be defined as follows:

$$\pi_{h12} = \frac{\hat{\lambda}_{h2} - (1 - P_{h2})}{(2P_{h2} - 1)} \quad (7)$$

$$\text{and } \pi_{h22} = \frac{\hat{\lambda}_{h2}^* - (1 - P_{h2}^*)}{(2P_{h2}^* - 1)} \quad (8)$$

$$\text{where; } \lambda_{h1} = P_{h2}\pi_h + (1 - P_{h1})(1 - \pi_h) = (2P_{h2} - 1)\pi_h + (1 - P_{h1}) \quad (9)$$

$$\lambda_{h2}^* = P_{h2}^*\pi_h + (1 - P_{h2}^*)(1 - \pi_h) = (2P_{h2}^* - 1)\pi_h + (1 - P_{h2}^*) \quad (10)$$

Which are the probabilities of ‘yes’ responses for the first and second use of  $R_{h2}$ . The variances of the estimators  $\hat{\pi}_{h12}$  and  $\hat{\pi}_{h22}$  are given by:

$$\text{Var}(\hat{\pi}_{h21}) = \frac{\lambda_{h1}(1 - \lambda_{h1})}{n_{h2}(2P_{h2} - 1)^2} = \frac{\pi_{h1}(1 - \pi_{h1})}{n_{h2}} + \frac{P_{h2}(1 - P_{h2})}{n_{h2}(2P_{h2} - 1)^2} \quad (11)$$

$$\text{and } \text{Var}(\hat{\pi}_{h22}) = \frac{\lambda_{h2}(1 - \lambda_{h2})}{n_{h2}(2P_{h2}^* - 1)^2} = \frac{\pi_{h2}(1 - \pi_{h2})}{n_{h2}} + \frac{P_{h2}^*(1 - P_{h2}^*)}{n_{h2}(2P_{h2}^* - 1)^2} \quad (12)$$

These were obtained from Warner’s RR model as given below. The first responses from respondents using  $R_{h2}$  can be defined as follows. The probability of a ‘yes’ response from the respondents using  $R_{h2}$  in the first response is given by:

$$\lambda_{h12} = P_{h1}\pi_h + (1 - P_{h1})(1 - \pi_h) \quad (13)$$

Also, the probability of a ‘no’ response from the respondents using  $R_{h2}$  in the first response is given by:

$$\lambda'_{h12} = P_{h1}(1 - \pi_h) + (1 - P_{h1})\pi_h \quad (14)$$

Among those that answered ‘no’ to the innocuous questions in stratum  $h$ ; suppose that  $n_{h2}$  report ‘yes’ and  $(n_h - n_{h2})$  report ‘no’ in first case, the likelihood of the sample in the same stratum is as follows:

$$\xi = [P_{h1}\pi_h + (1 - P_{h1})(1 - \pi_h)]^{n_{h2}} \times [P_{h1}(1 - \pi_h) + (1 - P_{h1})\pi_h]^{n_h - n_{h2}} \quad (15)$$

We also obtain the MLE of  $\pi_h$ , as follows:

$$\pi_{h12} = \frac{\hat{\lambda}_{h2} - (1 - P_{h2})}{(2P_{h2} - 1)} \quad (16)$$

Where; the proportion of ‘yes’ answers from  $R_{h1}$  in the sample is given as;

$$\hat{\lambda}_{h2} = \frac{n_{h2}}{n_h}$$

The variance of is obtained as follows:

$$\text{Var}(\pi_{h21}) = \left[ \frac{1}{n_h(2P_{h2} - 1)} \right]^2 \text{Var}(n_{h2}) \quad (17)$$

$$\text{Since; } \pi_{h21} = \frac{\frac{n_{h2}}{n_h} - (1 - P_{h2})}{(2P_{h2} - 1)} = \frac{n_{h2}}{n_h(2P_{h2} - 1)} + \frac{P_{h2} - 1}{2P_{h2} - 1}$$

$$\text{Then; } \text{Var}(\pi_{h21}) = \left[ \frac{1}{n_h(2P_{h2} - 1)} \right]^2 \text{Var}(n_{h2}) \\ = \frac{\text{Var}(X_{i2})}{n_h^2(2P_{h2} - 1)^2}$$

$$= \frac{[P_{h1}\pi_h + (1 - P_{h1})(1 - \pi_h)][P_{h1}(1 - \pi_h) + (1 - P_{h1})\pi_h]}{n_h(2P_{h2} - 1)^2}$$

Hence;  $Var(\hat{\pi}_{h21}) = \frac{\pi_h(1 - \pi_h)}{n_{h2}} + \frac{P_{h2}(1 - P_{h2})}{n_{h2}(2P_{h2} - 1)^2} = \frac{\lambda_{h1}(1 - \lambda_{h1})}{n_{h2}(2P_{h2} - 1)^2}$  (18)

Where;  $\lambda_{h1} = P_{h2}\pi_h + (1 - P_{h1})(1 - \pi_h) = (2P_{h2} - 1)\pi_h + (1 - P_{h1})$

The second response from  $R_{h2}$  have similar parameters; so that we have:

$$\pi_{h22} = \frac{\hat{\lambda}_{h2}^* - (1 - P_{h2}^*)}{(2P_{h2}^* - 1)}$$

and  $Var(\hat{\pi}_{h22}) = \frac{\pi_h(1 - \pi_h)}{n_{h2}} + \frac{P_{h2}^*(1 - P_{h2}^*)}{n_{h2}(2P_{h2}^* - 1)^2} = \frac{\lambda_{h2}(1 - \lambda_{h2})}{n_{h2}(2P_{h2}^* - 1)^2}$

where;  $\lambda_{h2}^* = P_{h2}^*\pi_h + (1 - P_{h2}^*)(1 - \pi_h) = (2P_{h2}^* - 1)\pi_h + (1 - P_{h2}^*)$

From Lanke (1976), to provide equal protection in  $R_{h1}$  and  $R_{h2}$  it can be shown that we must have either of the following:

$$P_{h2} = \frac{1}{2 - P_{h1}}$$

or  $P_{h2}^* = \frac{1}{2 - P_{h1}}$

With this restriction the variance of the estimators  $\hat{\pi}_{h12}$  and  $\hat{\pi}_{h22}$  become same. To estimate  $\pi_h$  from the information collected by the double use of  $R_{h2}$ , we defined an unbiased estimator as follows:

$$\hat{\pi}_{hP} = \lambda_1 \hat{\pi}_{h21} + \lambda_2 \hat{\pi}_{h22}$$

where;  $\lambda_1$  and  $\lambda_2$  are the weights assuming value 0.5 when  $Var(\hat{\pi}_{hP})$  is optimized.

Thus the  $\hat{\pi}_{hP}$  becomes:

$$\hat{\pi}_{hP} = \frac{\hat{\pi}_{h21} + \hat{\pi}_{h22}}{2}$$
 (19)

Its variance is given by:

$$Var(\hat{\pi}_{hP}) = \frac{Var(\hat{\pi}_{h21})}{2} = \frac{1}{2} \left[ \frac{\pi_h(1 - \pi_h)}{n_{h2}} + \frac{P_{h2}(1 - P_{h2})}{n_{h2}(2P_{h2} - 1)^2} \right]$$
 (20)

Since;  $Var(\hat{\pi}_{h21}) = Var(\hat{\pi}_{h22})$

and  $P_{h2} = 1 - P_{h2}^*$

An unbiased estimator in terms of all the information collected by both the random devices  $R_{h1}$  and  $R_{h2}$  in the  $h$ th stratum is defined as follows:

$$\hat{\pi}_{hP(tot)} = \pi_h = \frac{n_{h1}}{n_h} \hat{\pi}_{h1} + \frac{n_{h2}}{n_h} \hat{\pi}_{hP}$$
 (21)

As both the random devices  $R_{h1}$  and  $R_{h2}$  are independent, the variance of  $\hat{\pi}_{hP(tot)}$  under the restriction by Lanke (1976):

$$P_{h2} = \frac{1}{2 - P_{h1}}$$

Thus is given by:

$$Var(\hat{\pi}_{hP(tot)}) = \left( \frac{\lambda_h + 1}{2} \right) \frac{\pi_h(1 - \pi_h)}{n_h} + \frac{\lambda_h(1 - \pi_h)(1 - P_{h1})}{n_h P_{h1}} + \frac{(1 - \lambda_h)(1 - P_{h1})}{2n_h P_{h1}^2}$$
 (22)

where;  $\lambda_h = \frac{n_{h1}}{n_h}$

A stratified proportion estimator of the population proportion of the individuals with sensitive trait is defined as:

where;  $\hat{\pi}_{Sero} = \sum_{h=1}^L W_h \hat{\pi}_{hP(tot)}$  (23)

Its variance is given by:

$$Var(\hat{\pi}_{Sero}) = \sum_{h=1}^L \frac{W_h^2}{n_h} \left[ \pi_h (1 - \pi_h) \left( \frac{\lambda_h + 1}{2} \right) + \frac{\lambda_h (1 - \pi_h) (1 - P_{h1})}{P_{h1}} + \frac{(1 - \lambda_h) (1 - P_{h1})}{2P_{h1}^2} \right] \quad (24)$$

Its variance under the optimum allocation of total sample size into different strata is given by:

$$Var(\hat{\pi}_{Sero}) = \frac{1}{n} \left[ \sum_{h=1}^L W_h \left[ \left( \frac{\lambda_h + 1}{2} \right) \pi_h (1 - \pi_h) + \frac{\lambda_h (1 - P_{h1}) (1 - \pi_h)}{P_{h1}} + \frac{(1 - P_{h1}) (1 - \lambda_h)}{2P_{h1}^2} \right]^{\frac{1}{2}} \right]^2 \quad (25)$$

### 2.2 The Proposed HIV Seroprevalence Model II

The HIV seroprevalence surveys Model II requires that a sample respondent in stratum  $h$  to answer an innocuous direct question and asked to use the random device  $R_{h1}$  if his/her answer to direct question is “yes”. If answer to the direct question is “no”, he/she is requested to use another random device  $R_{h2}$ . The random device  $R_{h1}$  consists of two statements (i) “I am HIV positive” and (ii) “I am HIV negative”, presented with probabilities  $P_{h1}$  and  $(1 - P_{h1})$  respectively. Similarly, the random device  $R_{h2}$  consists of the two statements (i) “I am HIV positive” and (ii) “I am HIV negative”, presented with probabilities and  $P_{h2}$  and  $(1 - P_{h2})$  respectively. The probabilities of a ‘yes’ response from the respondents using  $R_{h1}$  and  $R_{h2}$  are respectively given by:

$$\lambda_{h1} = P_{h1} \pi_h + (1 - P_{h1}) \pi_{hy} = P_{h1} \pi_h + (1 - P_{h1}) \quad (26)$$

And

$$\lambda_{h2} = P_{h2} \pi_h + (1 - P_{h2}) \quad (27)$$

On the other hand, the probabilities of a ‘no’ response from the respondents using  $R_{h1}$  and  $R_{h2}$  are respectively given by:

$$\lambda'_{h1} = P_{h1} (1 - \pi_h) + (1 - P_{h1}) (1 - \pi_{hy}) = P_{h1} (1 - \pi_h) \quad (28)$$

And

$$\lambda'_{h2} = P_{h2} (1 - \pi_h) \quad (29)$$

Since the respondent using  $R_{h1}$  has already answered yes to the direct question,  $\pi_{hy} = 1$ .

Among those that answered ‘yes’ to the innocuous questions in stratum  $h$ ; suppose that  $n_{h1}$  report ‘yes’ and  $(n_h - n_{h1})$  report ‘no’, the likelihood of the sample in the same stratum is as follows:

$$\xi = [P_{h1} \pi_h + (1 - P_{h1})]^{n_{h1}} \times [P_{h1} (1 - \pi_h)]^{n_h - n_{h1}} \quad (30)$$

The natural log of the likelihood is given below:

$$\log \xi = n_{h1} \log [P_{h1} \pi_h + (1 - P_{h1})] + (n_h - n_{h1}) \log [P_{h1} (1 - \pi_h)] \quad (31)$$

To obtain the value of  $\pi_h$ , differentiate  $\log \xi$  w.r.t.  $\pi_h$  and equate to zero as follows:

$$\frac{\partial \log \xi}{\partial \pi_h} = \frac{n_{h1} P_{h1}}{P_{h1} \pi_h + (1 - P_{h1})} - \frac{(n_h - n_{h1}) P_{h1}}{P_{h1} (1 - \pi_h)} = 0 \quad (32)$$

$$\frac{n_{h1} P_{h1}}{P_{h1} \pi_h + (1 - P_{h1})} = \frac{(n_h - n_{h1}) P_{h1}}{P_{h1} (1 - \pi_h)}$$

$$n_{h1} P_{h1} (1 - \pi_h) = (n_h - n_{h1}) [P_{h1} \pi_h + (1 - P_{h1})]$$

$$\begin{aligned} n_{h1}P_{h1} - n_{h1}P_{h1}\pi_h &= n_hP_{h1}\pi_h + n_h - n_hP_{h1} - n_{h1}P_{h1}\pi_h - n_{h1} + n_{h1}P_{h1} \\ n_hP_{h1}\pi_h &= n_hP_{h1} - n_h + n_{h1} \\ \pi_h &= \frac{n_hP_{h1} - n_h + n_{h1}}{n_hP_{h1}} \end{aligned}$$

Hence, the unbiased estimators in terms of the responses of the respondents using  $R_{h1}$  is given by:

$$\hat{\pi}_{h1} = \frac{\hat{\lambda}_{h1} - (1 - P_{h1})}{P_{h1}} \quad (33)$$

Where the proportion of ‘yes’ answers from  $R_{h1}$  in the sample is  $\hat{\lambda}_{h1} = n_{h1} / n_h$ . The variance of  $\hat{\pi}_{h1}$  is obtained as follows:

$$\begin{aligned} Var(\hat{\pi}_{h1}) &= \left[ \frac{1}{P_{h1}} \right]^2 Var(\hat{\lambda}_{h1}) \\ &= \left[ \frac{1}{P_{h1}} \right]^2 \left( \frac{\hat{\lambda}_{h1}(1 - \hat{\lambda}_{h1})}{n_{h1}} \right) \\ &= \left[ \frac{1}{P_{h1}} \right]^2 \frac{[P_{h1}\pi_h + (1 - P_{h1})][P_{h1}(1 - \pi_h)]}{n_{h1}} \\ &= \frac{[P_{h1}\pi_h + (1 - P_{h1})](1 - \pi_h)}{n_{h1}P_{h1}} \end{aligned} \quad (34)$$

Hence;

$$Var(\hat{\pi}_{h1}) = \frac{(1 - \pi_h)(P_{h1}\pi_h + 1 - P_{h1})}{n_{h1}P_{h1}}$$

Similarly, the unbiased estimators in terms of the responses of the respondents using  $R_{h2}$  is given by:

$$\hat{\pi}_{h2} = \frac{\hat{\lambda}_{h2} - (1 - P_{h2})}{P_{h2}} \quad (35)$$

Where the proportion of ‘yes’ answers from  $R_{h2}$  in the sample is  $\hat{\lambda}_{h2} = n_{h2} / n_h$ . The variance of  $\hat{\pi}_{h2}$  is obtained as follows:

$$Var(\hat{\pi}_{h2}) = \frac{(1 - \pi_h)(P_{h2}\pi_h + 1 - P_{h2})}{n_{h2}P_{h2}}$$

In stratum  $h$  two randomization devices  $R_{h1}$  and  $R_{h2}$  are equally protective against the privacy of the respondents if  $P_{h1} = P_{h2} = P_h$ . Under this setting, the variances of the two unbiased estimators  $\hat{\pi}_{h1}$  and  $\hat{\pi}_{h2}$  become the same. We can also propose an estimator based on all the information collected in stratum  $h$  which we can use to estimate seroprevalence rates in stratum  $h$  as follows:

$$\hat{\pi}_h = \frac{n_{h1}}{n_h} \hat{\pi}_{h1} + \frac{n_{h2}}{n_h} \hat{\pi}_{h2} \quad (36)$$

Its variance is given by:

$$\begin{aligned} Var(\hat{\pi}_h) &= \left( \frac{n_{h1}}{n_h} \right)^2 Var(\hat{\pi}_{h1}) + \left( \frac{n_{h2}}{n_h} \right)^2 Var(\hat{\pi}_{h2}) \\ &= \left( \frac{n_{h1}}{n_h} \right)^2 \left[ \frac{(1 - \pi_h)(P_{h1}\pi_h + 1 - P_{h1})}{n_{h1}P_{h1}} \right] + \left( \frac{n_{h2}}{n_h} \right)^2 \left[ \frac{(1 - \pi_h)(P_{h2}\pi_h + 1 - P_{h2})}{n_{h2}P_{h2}} \right] \end{aligned} \quad (37)$$

$$= \left( \frac{n_{h1}}{n_h^2} \right) \left[ \frac{(1 - \pi_h)(P_{h1}\pi_h + 1 - P_{h1})}{P_{h1}} \right] + \left( \frac{n_{h2}}{n_h^2} \right) \left[ \frac{(1 - \pi_h)(P_{h2}\pi_h + 1 - P_{h2})}{P_{h2}} \right]$$

If we decide that  $P_{h1} = P_{h2} = P_h$  thus we get:

$$\begin{aligned} \text{Var}(\hat{\pi}_h) &= \left( \frac{n_{h1} + n_{h2}}{n_h^2} \right) \left[ \frac{(1 - \pi_h)(P_h\pi_h + 1 - P_h)}{P_h} \right] \\ &= \left( \frac{1}{n_h} \right) \left[ \frac{(1 - \pi_h)[P_h\pi_h + (1 - P_h)]}{P_h} \right] \end{aligned} \quad (38)$$

Hence;

$$\text{Var}(\hat{\pi}_h) = \frac{\pi_h(1 - \pi_h)}{n_h} + \frac{(1 - P_h)(1 - \pi_h)}{n_h P_h}$$

An unbiased stratified seroprevalence rates estimator is given by:

$$\hat{\pi}_{sero} = \sum_{h=1}^L W_h \hat{\pi}_h \quad (39)$$

where;  $W_h = N_h / N$  for is  $h = 1, 2, \dots, L$

$N_h$  is the total number of individuals in the stratum  $h$

$N$  is the total number of individuals in the population

obviously  $\sum_{h=1}^L W_h = 1$

Its variance is given by:

$$\text{Var}(\hat{\pi}_{sero}) = \sum_{h=1}^L \frac{W_h^2}{n_h} \left[ \pi_h(1 - \pi_h) + \frac{(1 - P_h)(1 - \pi_h)}{P_h} \right]^2 \quad (40)$$

Cochran (1977) established that the sampling fraction  $n_h / n$  is ignorable, then  $\text{Var}(\hat{\pi}_{sero})$  is minimized for a fixed total sample size  $n$  if:

$$n_h = \frac{n W_h \left[ \pi_h(1 - \pi_h) + \frac{(1 - P_h)(1 - \pi_h)}{P_h} \right]^{\frac{1}{2}}}{\sum_{h=1}^L W_h \left[ \pi_h(1 - \pi_h) + \frac{(1 - P_h)(1 - \pi_h)}{P_h} \right]^{\frac{1}{2}}} \quad (41)$$

where;  $n_h = n_{h1} + n_{h2}$

$$\sum_{h=1}^L n_h = n$$

Thus substituting the optimum value of  $n_h$  in (15) we get:

$$\text{Var}(\hat{\pi}_{sero}) = \frac{1}{n} \left[ \sum_{h=1}^L W_h \left[ \pi_h(1 - \pi_h) + \frac{(1 - P_h)(1 - \pi_h)}{P_h} \right]^{\frac{1}{2}} \right]^2 \quad (42)$$

### 3. Results

#### Computations from Model I

$$\hat{\pi}_{sero} = \sum_{i=1}^L W_h \hat{\pi}_{hP(tot)} \quad \text{where; } W_h = N_h / N \text{ for is } h = 1, 2, \dots, L$$

Its variance is given by:

$$Var(\hat{\pi}_{Sero}) = \sum_{h=1}^L \frac{W_h^2}{n_h} \left[ \pi_h(1-\pi_h) \left( \frac{\lambda_h+1}{2} \right) + \frac{\lambda_h(1-\pi_h)(1-P_{h1})}{P_{h1}} + \frac{(1-\lambda_h)(1-P_{h1})}{2P_{h1}^2} \right]$$

The computations for the model to estimate HIV seroprevalence rate give the following results:

$$\phi = \pi_h(1-\pi_h) \left( \frac{\lambda_h+1}{2} \right) + \frac{\lambda_h(1-\pi_h)(1-P_{h1})}{P_{h1}} + \frac{(1-\lambda_h)(1-P_{h1})}{2P_{h1}^2}$$

$$\hat{\pi}_{Sero} = \sum_{i=1}^L W_h \hat{\pi}_{hP(101)} = 0.0612$$

$$Var(\hat{\pi}_{Sero}) = \sum_{h=1}^L \frac{W_h^2}{n_h} \left[ \pi_h(1-\pi_h) \left( \frac{\lambda_h+1}{2} \right) + \frac{\lambda_h(1-\pi_h)(1-P_{h1})}{P_{h1}} + \frac{(1-\lambda_h)(1-P_{h1})}{2P_{h1}^2} \right]$$

$$Var(\hat{\pi}_{Sero}) = 0.000067$$

$$SE(\hat{\pi}_{Sero}) = \sqrt{Var(\hat{\pi}_{Sero})} = 0.0082$$

The 95% confidence interval for HIV seroprevalence rate using the two-way RR Model in stratification is given by:

$$(\hat{\pi}_{Sero}) \pm 1.96 \times SE(\hat{\pi}_{Sero}) = 0.0612 \pm 1.96 \times 0.0082 = [0.045, 0.077]$$

#### Computations from Model I

$$\phi = \pi_h(1-\pi_h) + (1-P_h)(1-\pi_h) / P_h$$

$$\hat{\pi}_{sero} = \sum_{h=1}^L W_h \hat{\pi}_h = 0.0874$$

$$Var(\hat{\pi}_{Sero}) = \sum_{h=1}^L \frac{W_h^2}{n_h} \left[ \pi_h(1-\pi_h) + \frac{(1-P_h)(1-\pi_h)}{P_h} \right]^2 = 0.00018$$

$$SE(\hat{\pi}_{Sero}) = \sqrt{Var(\hat{\pi}_{Sero})} = 0.0134$$

The 95% confidence interval for HIV seroprevalence rate is given by:

$$\hat{\pi}_{sero} \pm 1.96 \times SE(\hat{\pi}_{Sero}) = 0.0874 \pm 1.96 \times 0.0134 = [0.061, 0.114]$$

#### 4. Conclusion

This study was motivated by the fact that conventional data collection techniques usually cause evasive or untruthful responses when people are asked sensitive questions like their HIV serostatus. As a result, it is difficult to make accurate inferences from such unreliable data. Hence a two-way RR Model in stratification was devised using the work of Warner (1965), Arnab (2004), Quatember (2009), among others particularly for HIV seroprevalence surveys.

Furthermore, the model was used to estimate HIV seroprevalence rate in a small adult population using a sample size of 550 and a design parameter of 0.7. The result for model I show that, using the survey data, the model estimated the HIV seroprevalence rate as 6.1% with a standard error of 0.0082 and 95% confidence bands of [4.5%, 7.7%]. These estimates are for adults who are 18 years and above who attend a hospital. These results are consistent with that of Nigerian sentinel survey (2003) conducted by NACA, USAID and CDC which estimated the HIV seroprevalence in Kaduna State as 6.0%. Hence, the RRTs herein can serve as new viable methods for HIV seroprevalence surveys.

Similarly, the result for model II show that, using the survey data, the model estimated the HIV seroprevalence rate as 8.74% with a standard error of 0.0134 and 95% confidence bands of [6.1%, 11.4%]. These estimates are for adults who are 18 years and above who attend a hospital. Accordingly, the sentinel projected seroprevalence rate, using the EPP Package, for the next ten years (2013) was 9.7%; very consistent with the 95% confidence interval. Hence, the RRTs herein can also serve as new viable methods for HIV seroprevalence surveys. Model II has a better chance of estimating HIV seroprevalence because it has higher privacy preservation.

#### References

- Arnab, R. (2004). Optional randomized response techniques for complex survey designs. *Biometrical Journal* 46(1), 114-124.  
 Centre for Disease Control (1991). Pilot study of a household survey to determine HIV seroprevalence.



*Morbidity and Mortality Weekly Report* 40, 1-5.

Cochran, W.G. (1977). *Sampling techniques* (3rd ed.). New York: John Wiley and Sons.

Fox, J.A. and Tracy, P.E. (1986). *Randomized response: a method for sensitive surveys*. Beverly Hills, CA: Sage Publications Inc.

Houston, J. and Tran, Alfred. (2008). A survey of tax evasion using randomized response technique. *Journal of Statistical Planning and Inference* 148, 225-233.

Jarman, B. J. (1997). The Prevalence and Precedence of Socially Condoned Sexual aggression Within a Dating Context as Measured by Direct Questioning and Randomized response Technique. *Journal of Statistical Planning and Inference* 76, 229-238.

Kim, J.M. and Warde, W.D. (2005). A stratified Warner's randomized response model. *Journal of Statistical Planning and Inference* 120(2), 155-165.

Kuk, A.Y.C. (1990). Asking sensitive question indirectly. *Biometrika*, 77, 436-438.

Lee, R.M. (1993). *Doing research on sensitive topics*. Newbury Park, CA: Sage.

Lensvelt-Mulders, G.J.L.M.; Hox, J.J.; van der Heijden, P.G.M. and Maas, C.J.M. (2005). Meta-Analysis of Randomized Response Research: *Thirty-Five Years of Validation, Sociological Methods & Research* 33, 319-348.

Quatember, A. (2009). A standardization of randomized response strategies. *Survey Methodology*, 35(2), 143-152.

Rasinski, K. A., Willis, G. B., Baldwin, A. K., Yeh, W. and Lee, L. (1999). Methods of data collection, perception of risks and losses, and motivation to give truthful answers to sensitive survey questions. *Applied Cognitive Psychology* 22, 465-484.

Usman, A and Oshungade, I.O. (2012). A Two-way Randomized Response Technique in Stratification for Tracking HIV Seroprevalence. *Journal of Mathematical Theory and Modeling* 2(7), 86-97, 2012: *International Institute for Science, Technology and Education* (IISTE).

Usman, A and Oshungade, I.O. (2012). A Mixed-Stratified Randomized Response Technique Model for HIV Seroprevalence Surveys. *Research Journal of Mathematics and Statistics* 4(3), 70-75, 2012: *Maxwell Scientific Organization*.

Van der Hout, A., Van der Heijden, P.G.M and Gilchrist, R. (2002). A multivariate logistic regression model for randomized response data. *Quantitative Methods* 31, 25-41.

Volicer, B. J. and Volicer, L. (1982). Randomised response technique for estimating alcohol use and non compliance in hypertensives. *Journal of Studies in Alcohol* 43, 739-750.

Warner, S.L. (1965). Randomized response: a survey technique for eliminating evasive answer bias. *Journal of the American Statistical Association* 60, 63-69.

### Appendix Tables

**Table 1: Samples and Strata Sizes Model I**

Strata	Strata Description	$N_h$	$n_h$	$n_{h1}$	$n_{h21}$	$n_{h22}$	$W_h$
1	Married (Men/ Women)	1,285	189	32	42	38	0.344
2	Unmarried (Men/ Women)	2,020	297	56	55	63	0.540
3	Divorced/Separated/Widowed	435	64	12	10	11	0.116
Total		3,740	550	100	107	112	1.000

**Table 2: Summary of Results of the Random Devices Model I**

Strata	$\hat{\lambda}_{h1}$	$\hat{\pi}_{h1}$	$V(\hat{\pi}_{h1})$	$\hat{\lambda}_{h21}$	$\hat{\pi}_{h21}$	$V(\hat{\pi}_{h21})$	$\hat{\lambda}_{h22}$	$\hat{\pi}_{h22}$	$V(\hat{\pi}_{h22})$
1	0.376	0.109	0.0150	0.402	0.255	0.0358	0.365	0.163	0.0381
2	0.350	0.071	0.0083	0.401	0.253	0.0273	0.460	0.256	0.0246
3	0.343	0.061	0.0383	0.345	0.113	0.1412	0.379	0.198	0.0902

**Table 3: Summary of Computations Model I**

Strata	$\frac{n_{h1}}{n_h}$	$\hat{\pi}_{hP}$	$\frac{n_{h2}}{n_h}$	$\hat{\pi}_h$	$W_h \hat{\pi}_h$	$\frac{W_h^2}{n_h}$	$\hat{\pi}_h(1 - \hat{\pi}_h)$	$\phi$	$\sum_{h=1}^L \frac{W_h^2}{n_h} \phi$
1	0.169	0.209	0.201	0.060	0.0206	0.00063	0.056	0.037	0.000023
2	0.189	0.255	0.212	0.067	0.0362	0.00098	0.063	0.041	0.000040
3	0.188	0.156	0.172	0.038	0.0044	0.00021	0.037	0.019	0.000004
Total					0.0612				0.000067

**Table 4: Summary of Seroprevalence Results Model I**

N	n	$\hat{\pi}_{Sero}$	$Var(\hat{\pi}_{Sero})$	95% confidence interval	
				Lower limit	Upper limit
3,740	550	0.0610	0.000067	0.045	0.077

**Table 5: Samples and Strata Sizes Model II**

Strata	Strata Description	$N_h$	$n_h$	$n_{h1}$	$n_{h2}$	$W_h$
1	Married (Men/ Women)	1,285	189	35	38	0.344
2	Unmarried (Men/ Women)	2,020	297	57	58	0.540
3	Divorced/Separated/Widowed	435	64	11	9	0.116
Total		3,740	550	103	105	1.000

**Table 6: Summary of Results of the Random Devices Model II**

Strata	$\hat{\lambda}_{h1}$	$\hat{\pi}_{h1}$	$V(\hat{\pi}_{h1})$	$\hat{\lambda}_{h2}$	$\hat{\pi}_{h2}$	$V(\hat{\pi}_{h2})$	$\hat{\pi}_h$	$V(\hat{\pi}_h)$
1	0.365	0.093	0.0135	0.409	0.156	0.0130	0.098	0.0052
2	0.383	0.119	0.0085	0.392	0.131	0.0838	0.097	0.0033
3	0.324	0.034	0.0406	0.300	0.000	0.0476	0.011	0.0156

**Table 7: Summary of Computations Model II**

Strata	$W_h$	$\hat{\pi}_h$	$W_h \hat{\pi}_h$	$W_h^2 / n_h$	$\hat{\pi}_h(1 - \hat{\pi}_h)$	$\sum_{h=1}^L \frac{W_h^2}{n_h} \phi^2$
1	0.344	0.098	0.0337	0.00063	0.156	0.000056
2	0.540	0.097	0.0524	0.00098	0.131	0.000088
3	0.116	0.011	0.0013	0.00021	0.000	0.000036
Total	1.000		0.0874			0.000180

**Table 8: Summary of Seroprevalence Results Model II**

N	n	$\hat{\pi}_{Sero}$	$Var(\hat{\pi}_{Sero})$	95% confidence interval	
				Lower limit	Upper limit
3,740	550	0.0874	0.00018	0.061	0.113

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

## CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

**Prospective authors of journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

## MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

