

Common Fixed Points For Three Maps In Cone Metric Spaces

K.Prudhvi* and M.Rangamma

Department of Mathematics ,Osmania University, Hyderabad, Andhra Pradesh, India.

* E-mail of the corresponding author : prudhvikasani@rocketmail.com ; rangamma1999@gmail.com

Abstract

The existence of coincidence points and common fixed point theorem for three maps satisfying certain contractive conditions without exploiting the notation of continuity of any map involved therein cone metric space is proved . Our result extends and generalize some recent results .

Keywords: Cone metric space , Common Fixed Point , Coincidence point.

1. Introduction and preliminaries.

The study of common fixed points of mappings satisfying certain contractive conditions has been at the centre of vigorous research activity ,being the applications of fixed point very important in several areas of mathematics . In 2007, Huang and Zhang [5] generalized the concept of a metric space, replacing the set of real numbers by an ordered Banach space and obtained some fixed point theorems for mapping satisfying different contractive conditions. Subsequently Abbas and Jungck [1] and Abbas and Rhoades [2] have studied common fixed point theorems in cone metric spaces (see also,[5],[7] and the references mentioned there in). Jungck [6] defined a pair of self-mappings to be weakly compatible if they commute at their coincidence points. In recent years several authors have obtained coincidence point results for various class of mappings on a metric space , utilizing these concepts .For survey of coincidence point theory ,its applications ,comparison of different contractive conditions and related results ,we refer to [3],[4] and the references therein. The aim of this paper is to present coincidence points and common fixed point results for three mappings which satisfy generalized contractive conditions. These theorems generalize and extends the results of Abbas and Jungck [1]. In all that follows , E is a real Banach space, $N = \{1,2,3,\dots\}$ the set of all natural numbers . For the mappings $f,g :X \rightarrow X$,let $C(f,g)$ denotes set of coincidence points of f, g , that is $C(f,g) := \{z \in X : fz = gz\}$.

We recall some definitions of cone metric spaces and some of their properties [5] .

Definition 1.1. Let E be a real Banach Space and P a subset of E .The set P is called a cone if and only if :

- (a) P is closed , non empty and $P \neq \{0\}$;
- (b) $a,b \in R$, $a,b \geq 0, x,y \in P$ implies $ax + by \in P$;
- (c) $x \in P$ and $-x \in P$ implies $x = 0$.

Definition 1.2. Let P be a cone in a Banach Space E , define partial ordering ' \leq ' with respect to P by $x \leq y$ if and only if $y-x \in P$.We shall write $x < y$ to indicate $x \leq y$ but $x \neq y$ while $X \ll y$ will stand for $y - x \in \text{int}P$, where $\text{int}P$ denotes the interior of the set P . This Cone P is called an order cone .

Definition 1.3. Let E be a Banach Space and $P \subset E$ be an order cone .The order cone P is called normal if there exists $K > 0$ such that for all $x,y \in E$,

$$0 \leq x \leq y \text{ implies } \|x\| \leq K \|y\|.$$

The least positive number K satisfying the above inequality is called the normal constant of P .

Definition 1.4. Let X be a nonempty set of E .Suppose that the map $d: X \times X \rightarrow E$ satisfies :

- (d1) $0 \leq d(x,y)$ for all $x,y \in X$ and
 $d(x,y) = 0$ if and only if $x = y$;
 (d2) $d(x,y) = d(y,x)$ for all $x,y \in X$;
 (d3) $d(x,y) \leq d(x,z) + d(y,z)$ for all $x,y,z \in X$.

Then d is called a cone metric on X and (X,d) is called a cone metric space .

It is obvious that the cone metric spaces generalize metric spaces.

Example 1.5. ([5]) . Let $E = \mathbb{R}^2$, $P = \{ (x , y) \in E \text{ such that } : x, y \geq 0 \} \subset \mathbb{R}^2$,
 $X = \mathbb{R}$ and $d: X \times X \rightarrow E$ such that $d(x , y) = (|x - y| , \alpha |x - y|)$, where $\alpha \geq 0$ is a constant . Then (X,d)
 is a cone metric space.

Definition 1.6. Let (X,d) be a cone metric space .We say that $\{x_n\}$ is
 (i) a Cauchy sequence if for every c in E with $0 \ll c$, there is N such
 that for all $n , m > N$, $d(x_n, x_m) \ll c$;
 (ii) a convergent sequence if for any $0 \ll c$,there is an N such that for
 all $n > N$, $d(x_n, x) \ll c$, for some fixed x in X . We denote this
 $x_n \rightarrow x (n \rightarrow \infty)$.

A cone metric space X is said to be complete if every Cauchy sequence in X is convergent in X .

Definition 1.7. Let $f , g : X \rightarrow X$. Then the pair (f , g) is said to be (IT)-Commuting at
 $z \in X$ if $f(g(z)) = g(f(z))$ with $f(z) = g(z)$.

2 . Common fixed point theorem

In this section we obtain coincidence points and common fixed point theorems for three maps in cone metric spaces.

The following theorem extends and improves Theorem 2.1 of [1].

Theorem 2.1 . Let (X,d) be a cone metric space , and P a normal cone with normal constant K . Suppose mappings $f , g , h : X \rightarrow X$ satisfy :

$$d(fx , gy) \leq k d(hx , hy) \text{ for all } x , y \in X , \text{ with } x \neq y. \quad (2.1)$$

where $k \in [0,1)$ is a constant . If $f(X) \cup g(X) \subset h(X)$ and $h(X)$ is a complete subspace of X .Then the maps f , g and h have a coincidence point p in X . Moreover if (f , h) and (g , h) are (IT) – Commuting at p , then f , g and h have a unique common fixed point.

Proof. Suppose x_0 is an arbitrary point of X ,and define the sequence $\{y_n\}$ in X such that

$$\begin{aligned} y_{2n} &= fx_{2n} = hx_{2n+1} \\ \text{and } y_{2n+1} &= gx_{2n+1} = hx_{2n+2}, \text{ for all } n = 0,1,2,\dots \end{aligned}$$

By (2.1) ,we have

$$\begin{aligned} d(y_{2n} , y_{2n+1}) &= d(fx_{2n} , gx_{2n+1}) \\ &\leq k d(hx_{2n} , hx_{2n+1}) \\ &\leq k d(y_{2n-1} , y_{2n}) . \end{aligned}$$

Similarly , it can be shown that

$$d(y_{2n+1}, y_{2n+2}) \leq k d(y_{2n}, y_{2n+1}) .$$

Therefore ,for all n ,

$$d(y_{n+1}, y_{n+2}) \leq k d(y_n, y_{n+1}) \leq \dots \leq k^{n+1} d(y_0, y_1) .$$

Now, for any m > n ,

$$\begin{aligned} d(y_n, y_m) &\leq d(y_n, y_{n+1}) + d(y_{n+1}, y_{n+2}) + \dots + d(y_{m-1}, y_m) . \\ &\leq [k^n + k^{n+1} + \dots + k^{m-1}] d(y_1, y_0) \end{aligned}$$

$$\leq \frac{k^n}{1-k} d(y_1, y_0) .$$

From (1.3), we have

$$\|d(y_n, y_m)\| \leq \frac{k^n}{1-k} K \|d(y_1, y_0)\| ,$$

which implies that $d(y_n, y_m) \rightarrow 0$ as $n, m \rightarrow \infty$.

Hence $\{y_n\}$ is a Cauchy sequence, where $y_n = \{hx_n\}$.

Therefore $\{hx_n\}$ is a Cauchy sequence. Since $h(X)$ is complete, there exists q in $h(X)$ such that $hx_n \rightarrow q$ as $n \rightarrow \infty$. Consequently, we can find a point p in X such that $h(p) = q$. We shall show that $hp = fp = gp$.

Note that $d(hp, fp) = d(q, fp)$. Let us estimate $d(hp, fp)$

We have , by the triangle inequality

$$\begin{aligned} d(hp, fp) &\leq d(hp, hx_{2n+2}) + d(hx_{2n+2}, fp) \\ &= d(q, hx_{2n+2}) + d(fp, gx_{2n+1}) . \end{aligned}$$

By the contractive condition , we get.

$$\begin{aligned} d(fp, gx_{2n+1}) &\leq kd(hp, hx_{2n+1}) \\ &= kd(q, hx_{2n+1}) \rightarrow 0 \text{ as } n \rightarrow \infty . \text{ Since } \{hx_n\} \rightarrow q \text{ as } n \rightarrow \infty . \end{aligned}$$

Therefore , for large n , we get

$$d(hp, fp) \leq d(q, hx_{2n+2})$$

$\leq d(q, q) = 0$.
 Which leads to

$$d(hp, fp) = 0 \text{ and hence } hp = q = fp \text{ .} \quad (2.2)$$

$$\text{Similarly we can show } hp = q = gp \text{ .} \quad (2.3)$$

From (2.2) and (2.3) , it follows

$$q = hp = fp = gp \text{ , } p \text{ is a coincidence point of } f, g, h \text{ .} \quad (2.4)$$

Since (f, h) and (g, h) are (IT) – commuting at p .

We get by (2.4) and contractive condition ,

$$\begin{aligned} d(ffp, fp) &= d(ffp, gp) \leq kd(hfp, hp) \\ &< d(hfp, hp) = d(ffp, fp) \\ &= d(fhp, hp) = d(ffp, fp) \end{aligned}$$

$d(ffp, fp) < d(ffp, fp)$, a contradiction.

Therefore $ffp = fp$.

$$fp = ffp = fhp = hfp \text{ implies } ffp = hfp = fp = q.$$

$$\text{Therefore , } fp (=q) \text{ is a common fixed point of } f \text{ and } h \text{ .} \quad (2.5)$$

Similarly, we get $gp = ggp = ghp = hgp$, implies $ggp = hgp = gp = q$.

$$\text{Therefore , } gp = fp (=q) \text{ is a common fixed point of } g \text{ and } h \quad (2.6)$$

In view of (2.5) and (2.6) it follows f, g and h have a common fixed point namely q . The uniqueness of the common fixed point of q follows (2.1). Indeed ,

let q_1 be another common fixed point of f, g and h .

$$\text{Consider , } d(q, q_1) = d(fq, gq_1)$$

$$\leq k d(hq, hq_1) \text{ . As } k < 1 \text{ , it follows that}$$

$$d(q, q_1) = 0.$$

That is $q = q_1$.

Therefore f, g and h have a unique common fixed point.

Remark 2.2. If we take $g = f$ and $h = g$ in Theorem 2.1, then we obtain Theorem 2.1 of [1]. Also if we let $g = f$, $h = g$ and g is identity map on X in Theorem 2.1 , then we obtain Theorem 1 of [5] , that is , the

extension of Banach Fixed Point Theorem for cone metric spaces.

Acknowledgement : The authors are great full to the referees to careful reading of article.

References

- [1] M.Abbas and G.Jungck, common fixed point results for non commuting mappings without continuity in cone metric spaces. *J.math.Anal.Appl.* 341(2008) 416-420.
- [2] M.Abbas and B.E.Rhoades , Fixed and periodic point results in cone metric Spaces. *Appl.Math .Lett.*,22(2009),511-515.
- [3] V.Berinde ,A common fixed point theoremfor compactible quasi contractive self mappings in metric spaces ,*Appl.Math .Comput.*,213(2009),348-354.
- [4] B.Fisher ,Four mappings with a common fixed point *J.Univ .Kuwait Sci.*,(1981),131-139.
- [5] L.G.Huang, X.Zhang, cone metric spaces and fixed point theorems of contractive mappings, *J.Math.Anal.Appl.*332(2)(2007)1468-1476.
- [6] G.Jungck, common fixed points for non continuous non self maps on non-metric spaces, *Far East J.Math.Sci.(FJMS)*4(1996) 199-215.
- [7] S.Rezapour and Halbarani, some notes on the paper “cone metric spaces and fixed point theorem of contractive mappings “ ,*J.Math . Anal . Appl.*, 345(2008), 719-724.
- [8] S.L.Singh, Apichai Hematulin and R.P.Pant, new coincidence and common fixed point theorem, *Applied General Topology* 10(2009), no.1, 121-130.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

