

A Mathematical Model for Integrating Product of Two Functions

M.O. Oke

Department of Electrical & Electronic Engineering, Ekiti State University,

P.M.B 5363, Ado-Ekiti, Nigeria

E-mail: femioke91@gmail.com

Abstract

Integration by parts is a well-known method of integrating product of two functions. If the integrand involves a polynomial $f(x)$ of degree n and another function $g(x)$ that can be integrated at least $(n+1)$ times, then the solution to $\int f(x)g(x)dx$ is guaranteed after n routine applications of integration by parts method. This paper presents a mathematical model for integrating product of two functions without going through the routine application of integration by parts at each stage of integration, thus saving a lot of computational time. Some examples were considered to illustrate the effectiveness of the model. The model is found to be appropriate as it gives the same result with the well-known integration by parts method.

Keywords: Integrand; Mathematical model; Integration by parts; Differentiable functions

1. Introduction

Integration by parts is a method used for integrating product of two functions particularly when either function is not a derivative of the other (Stroud & Dexter 2007).

If $f(x)$ and $g(x)$ are two differentiable functions of x , then

$$\frac{d}{dx}(f \cdot g) = f \frac{dg}{dx} + g \frac{df}{dx} \quad (1)$$

(1) is the formula for the derivative of product of two functions as we can see in

Bhattacharyya (2009), Matthew & Alabi (2008), Dass (2009), Bajpai et al. (1981),

Richmond (1972) and Oke (2003), where $f = f(x)$ and $g = g(x)$.

Rearranging equation (1) we have:

$$f \frac{dg}{dx} = \frac{d}{dx}(f \cdot g) - g \frac{df}{dx} \quad (2)$$

Integrating equation (2) with respect to x , we have:

$$\int f dg = f \cdot g - \int g df \quad (3)$$

(3) is the formula for integration by parts as we can see in Kreyszig (1987), Thomas JR. & Finney (1984), Daniel (2000), Gupta (2009), Grossman (1985), and Goldstein et al. (2007).

Let us assume that $f(x)$ is a polynomial of degree n and let $g(x)$ be a function of x that can be integrated at least $(n+1)$ times. To evaluate $\int f(x)g(x)dx$ using the integration by parts method, we will need n applications of the formula in (3) above before we can get a solution. In this paper, we derived a mathematical model for integrating product of two functions. In applying the model, we don't need to go through the routine application of integration by parts each time we are integrating product of two functions. We only need to substitute the derivatives of $f(x)$ and the integrals of $g(x)$ in the newly derived model.

2. Materials and Methods

Let $f(x)$ and $g(x)$ be two functions of x , where $f(x)$ is a polynomial of degree n and $g(x)$ is a function of x that can be integrated at least $(n+1)$ times. Then:

$$\int f(x)g(x)dx = f(x) \int g(x)dx - \int (f'(x) \int g(x)dx)dx \quad (4)$$

$$= f(x) \int g(x)dx - f'(x) \iint g(x)(dx)^2 + \int (f''(x) \iint g(x)(dx)^2) dx \quad (5)$$

$$= f(x) \int g(x)dx - f'(x) \iint g(x)(dx)^2 + f''(x) \iiint g(x)(dx)^3 - \int (f'''(x) \iiint g(x)(dx)^3)dx \quad (6)$$

$$= f(x) \int g(x)dx - f'(x) \iint g(x)(dx)^2 + f''(x) \iiint g(x)(dx)^3 - f'''(x) \iiint \iint g(x)(dx)^4 + \int (f^{(4)}(x) \int \int \int \int g(x)(dx)^4)dx \quad (7)$$

Since $f(x)$ is a polynomial of degree n and $g(x)$ can be integrated at least $(n + 1)$ times, the procedure above will continue until the last stage where we now have:

$$\begin{aligned} \int f(x)g(x)dx &= f(x) \int g(x)dx - f'(x) \iint g(x)(dx)^2 + f''(x) \iiint g(x)(dx)^3 \\ &- f'''(x) \iiint \iint g(x)(dx)^4 + f^{(4)}(x) \iiint \iint \iint g(x)(dx)^5 - f^{(5)}(x) \iiint \iint \iint \iint g(x)(dx)^6 \\ &+ \dots \dots \dots + (-1)^{n-1} [f^{(n-1)}(x) \int \int \dots (n) \text{times} \dots \int g(x)(dx)^n] \\ &+ (-1)^n [\int (f^{(n)}(x) \int \int \dots (n) \text{times} \dots \int g(x)(dx)^n)dx] \end{aligned} \quad (8)$$

$$\begin{aligned} \int f(x)g(x)dx &= f(x) \int g(x)dx - f'(x) \iint g(x)(dx)^2 + f''(x) \iiint g(x)(dx)^3 \\ &- f'''(x) \iiint \iint g(x)(dx)^4 + f^{(4)}(x) \iiint \iint \iint g(x)(dx)^5 - f^{(5)}(x) \iiint \iint \iint \iint g(x)(dx)^6 \\ &+ \dots \dots \dots + (-1)^{n-1} [f^{(n-1)}(x) \int \int \dots (n) \text{times} \dots \int g(x)(dx)^n] \\ &+ (-1)^n [f^{(n)}(x) \int \int \dots (n + 1) \text{times} \dots \int g(x)(dx)^{n+1}] \end{aligned} \quad (9)$$

Putting all these in a more compact form, we have:

$$\int f(x)g(x)dx = \sum_{j=0}^n (-1)^j f^j(x) \int g(x)^{[j+1]} dx \quad (10)$$

where $f^j(x)$ is the j^{th} derivative of $f(x)$ and $\int g(x)^{[j+1]} dx$ represents the $(j + 1)^{\text{th}}$ integral of $g(x)$ with respect to x .

(10) above is the mathematical model for integrating product of two functions when the integrand involves a polynomial $f(x)$ of degree n and another function $g(x)$ which can be integrated at least $(n+1)$ times.

In order to apply the model to $\int f(x)g(x) dx$, we will find the derivatives of $f(x)$ up to the n^{th} derivative and the integrals of $g(x)$ up to the $(n + 1)^{\text{th}}$ integral and substitute all these into the formula in (10) above to get our final result directly.

3. Computational Examples

Example 1:

Let us consider $\int (x^4 + 5x^2 + 3x + 4) \sin 3x dx$

$$f(x) = x^4 + 5x^2 + 3x + 4 \text{ and } g(x) = \sin 3x$$

$$f^1(x) = 4x^3 + 10x + 3 ,$$

$$f^2(x) = 12x^2 + 10,$$

$$f^3(x) = 24x$$

$$\text{and } f^4(x) = 24.$$

From our notation:

$$\int g(x)^{[1]} dx = \int g(x) dx = \int \sin 3x dx = -\frac{1}{3} \cos 3x.$$

$$\int g(x)^{[2]} dx = \iint g(x) (dx)^2 = \iint g(x) dx dx = -\frac{1}{9} \sin 3x.$$

Similarly

$$\int g(x)^{[3]} dx = \frac{1}{27} \cos 3x.$$

$$\int g(x)^{[4]} dx = \frac{1}{81} \sin 3x$$

$$\int g(x)^{[5]} dx = -\frac{1}{243} \cos 3x.$$

Substituting all these into the formula in (10) above we have our final result directly as:

$$\begin{aligned} \int (x^4 + 5x^2 + 3x + 4) \sin 3x dx &= -\frac{1}{3} (x^4 + 5x^2 + 3x + 4) \cos 3x \\ &+ \frac{1}{9} (4x^3 + 10x + 3) \sin 3x + \frac{1}{27} (12x^2 + 10) \cos 3x - \frac{1}{81} (24x) \sin 3x - \frac{24}{243} \cos 3x + C. \end{aligned}$$

where C is the constant of integration.

Using the well-known integration by parts method to solve the problem, we have:

$$\int (x^4 + 5x^2 + 3x + 4) \sin 3x dx = -\frac{1}{3}(x^4 + 5x^2 + 3x + 4) \cos 3x + \frac{1}{3} \int (4x^3 + 10x + 3) \cos 3x dx$$

$$= -\frac{1}{3}(x^4 + 5x^2 + 3x + 4) \cos 3x + \frac{1}{9} (4x^3 + 10x + 3) \sin 3x$$

$$- \frac{1}{9} \int (12x^2 + 10) \sin 3x dx$$

$$= -\frac{1}{3}(x^4 + 5x^2 + 3x + 4) \cos 3x + \frac{1}{9} (4x^3 + 10x + 3) \sin 3x + \frac{1}{27} (12x^2 + 10) \cos 3x - \frac{1}{27} \int 24x \cos 3x dx$$

$$= -\frac{1}{3}(x^4 + 5x^2 + 3x + 4) \cos 3x + \frac{1}{9} (4x^3 + 10x + 3) \sin 3x + \frac{1}{27} (12x^2 + 10) \cos 3x - \frac{1}{81} 24x \sin 3x + \frac{1}{81} \int 24 \sin 3x dx$$

$$\int (x^4 + 5x^2 + 3x + 4) \sin 3x dx = -\frac{1}{3}(x^4 + 5x^2 + 3x + 4) \cos 3x$$

$$+ \frac{1}{9} (4x^3 + 10x + 3) \sin 3x + \frac{1}{27} (12x^2 + 10) \cos 3x - \frac{1}{81} (24x) \sin 3x - \frac{24}{243} \cos 3x + C$$

where C is the constant of integration.

This is the same with the result we got before.

Example 2:

Let us evaluate $\int (x^5 + 3x^4 + 2x) e^{2x} dx$.

$$f(x) = x^5 + 3x^4 + 2x \text{ and } g(x) = e^{2x}.$$

$$f^1(x) = 5x^4 + 12x^3 + 2,$$

$$f^2(x) = 20x^3 + 36x^2,$$

$$f^3(x) = 60x^2 + 72x,$$

$$f^4(x) = 120x + 72$$

$$\text{and } f^5(x) = 120.$$

From our notation:

$$\int g(x)^{[1]} dx = \frac{1}{2} e^{2x}$$

$$\int g(x)^{[2]} dx = \frac{1}{4} e^{2x}$$

$$\int g(x)^{[3]} dx = \frac{1}{8} e^{2x}$$

$$\int g(x)^{[4]} dx = \frac{1}{16} e^{2x}$$

$$\int g(x)^{[5]} dx = \frac{1}{32} e^{2x}$$

$$\int g(x)^{[6]} dx = \frac{1}{64} e^{2x}$$

Substituting all these into formula (10) above, we have our final result directly as:

$$\int (x^5 + 3x^4 + 2x)e^{2x} dx = \left[\frac{1}{2}(x^5 + 3x^4 + 2x) - \frac{1}{4}(5x^4 + 12x^3 + 2) + \frac{1}{8}(20x^3 + 36x^2) - \frac{1}{16}(60x^2 + 72x) + \frac{1}{32}(120x + 72) - \frac{15}{8} \right] e^{2x} + C.$$

where C is the constant of integration.

The integration by parts method, for this problem in example 2, gave us the same result after five routine applications.

Example 3:

Let us consider $\int x^2 \ln x dx$

$$f(x) = x^2 \text{ and } g(x) = \ln x.$$

$$f^1(x) = 2x \text{ and } f^2(x) = 2.$$

From our notation:

$$\int g(x)^{[1]} dx = x(\ln x - 1).$$

$$\int g(x)^{[2]} dx = \frac{x^2}{2} (\ln x - \frac{3}{2})$$

$$\int g(x)^{[3]} dx = \frac{x^3}{6} (\ln x - \frac{11}{6})$$

Putting all these in (10) above, we have our final result directly as:

$$\begin{aligned} \int x^2 \ln x dx &= x^3(\ln x - 1) - x^3 \left(\ln x - \frac{3}{2} \right) + \frac{x^3}{3} \left(\ln x - \frac{11}{6} \right). \\ &= \frac{x^3}{3} \left(\ln x - \frac{1}{3} \right) + C. \end{aligned}$$

where C is the constant of integration.

We also got the same result after two applications when we used the integration by parts method for the problem in example 3.

Example 4:

Let us consider $\int (x^5 + 3x^3 + 2x) \cos x dx$

$$f(x) = x^5 + 3x^3 + 2x \text{ and } g(x) = \cos x$$

$$f^1(x) = 5x^4 + 9x^2 + 2,$$

$$f^2(x) = 20x^3 + 18x,$$

$$f^3(x) = 60x^2 + 18,$$

$$f^4(x) = 120x$$

$$\text{and } f^5(x) = 120.$$

From our notation:

$$\int g(x)^{[1]} dx = \sin x$$

$$\int g(x)^{[2]} dx = -\cos x$$

$$\int g(x)^{[3]} dx = -\sin x$$

$$\int g(x)^{[4]} dx = \cos x$$

$$\int g(x)^{[5]} dx = \sin x$$

$$\int g(x)^{[6]} dx = \cos x$$

Putting all these in (10) above, we have our final result directly as:

$$\int (x^5 + 3x^3 + 2x) \cos x dx = (x^5 + 3x^3 + 2x) \sin x + (5x^4 + 9x^2 + 2) \cos x - (20x^3 + 18x) \sin x - (60x^2 + 18) \cos x + 120x \sin x + 120 \cos x + C.$$

C is the constant of integration.

We got the same result for this example when we used the integration by parts method after five routine applications.

Conclusion

In this paper, a mathematical model for integrating product of two functions was presented for an integrand involving a polynomial of degree n. The advantage of the model is in the computational time that is saved. We don't need to go through the routine applications of integration by parts method each time we are integrating product of two functions. We only need to substitute the derivatives of $f(x)$ up to the n^{th} derivative and the integrals of $g(x)$ up to the $(n+1)^{\text{th}}$ integral in the model. The mathematical model is found to be appropriate as it gives the same result with the integration by parts method.

References

- Bajpai, A.C., Calus, I.M., Fairley, J.A. & Walker, D. (1981), "Mathematics for Engineers and Scientists", *John Wiley and Sons*.
- Bhattacharyya, B. (2009), "Mathematical Physics", *New Central Book Agency (P) Ltd*.
- Daniel, A.A. (2000), "Application of Integral Calculus to Selected Science and Engineering Problems", *Ada Journal of Science*, 8(2), 74-92.
- Dass, H.K. (2009), "Engineering Mathematics", *S. Chang & Company Ltd*.
- Goldstein, L.J., Lay, D.C., Schneider, D.I. & Asmar, N.H. (2007), "Calculus and its Applications", *Pearson Education International*.
- Grossman, S.I. (1985), "Applied Calculus", *Wadsworth Publishing Company*.
- Gupta, B.D. (2009), "Mathematical Physics", *Vikas Publishing House PVT Ltd*.
- Kreyszig, E. (1987) "Advanced Engineering Mathematics", *Wiley Eastern Limited*.
- Matthew, D.A. & Alabi, P.O. (2008) "An Overview of Integration Techniques for Solving Engineering Problems", *Doe Journal of Science and Technology*, 5(3), 80-98.
- Oke, M.O. (2003), "Calculus of One and Several Variables for Scientists and Engineers", *Petoa Educational Publishers*.
- Richmond, A.E. (1972), "Calculus for Electronics", *Mc Graw-Hill Books Company*.
- Stroud, K.A. & Dexter, J.B. (2007), "Engineering Mathematics", *Palgrave Macmillan Ltd*.
- Thomas JR, G.B. & Finney, R.L. (1984) "Calculus and Analytic Geometry", *Addison-Wesley Publishing Company*.

Dr. M.O. OKE was born at Are – Ekiti in Nigeria on 16th April, 1969. He is a senior lecturer in the Department of Electrical and Electronic Engineering of Ekiti State University. He graduated with a B.Sc. degree in Industrial Mathematics from the University of Benin, Nigeria in 1991. He got a Postgraduate Diploma in Electrical and Electronic Engineering from the University of Ado-Ekiti, Nigeria and finished with a distinction in 2002. He had his M.Sc. degree in Mathematics from the University of Ilorin, Nigeria in 2005 and his Ph.D degree in Mathematics at the University of Ilorin, Nigeria this year (2012). Dr. M.O. Oke became a member of Nigerian Mathematical Society in 1992, Mathematical Association of Nigeria in 1994, National Association of Mathematical Physics in 1995 and American Mathematical Society in 2011. Dr. M.O. OKE specializes in Mathematical Modeling, Optimization and Numerical Analysis.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

