

OF GHOST FIELDS, CELESTIAL MONSTERS AND HELLHOUNDS-A FORTY SEVEN STOREY MODEL

¹DR K N PRASANNA KUMAR, ²PROF B S KIRANAGI AND ³PROF C S BAGEWADI

ABSTRACT: *We study a consolidated system of event; cause and n Qubit register which makes computation with n Qubits. Model extensively dilates upon systemic properties and analyses the systemic behaviour of the equations together with other concomitant properties. Inclusion of event and cause, we feel enhances the "Quantum ness" of the system holistically and brings out a relevance in the Quantum Computation on par with the classical system, in so far as the analysis is concerned. Additional VARIABLES OF Space Time provide bastion for the quantum space time studies.*

INTRODUCTION:

EVENT AND ITS VINDICATION:

There definitely is a sense of compunction, contrition, hesitation, regret, remorse, hesitation and reservation to the **acknowledgement of** the fact that there is a personal **relation to** what **happens to** oneself. Louis de Broglie said that the events have already happened and it **shall disclose** to the people **based on** their level of consciousness. So there is destiny to start with! Say I am undergoing some seemingly insurmountable problem, which has hurt my sensibilities, susceptibilities and sentimentalities that I refuse to accept that that event was waiting for me to happen. In fact this is the statement of stoic philosophy which is referred to almost as bookish or abstract. Wound is there; it **had to happen** to me. So I was wounded. Stoics tell us that the wound **existed** before me; I was born **to embody** it. It is the question of consummation, consolidation, concretization, consubstantiation, that of this, that **creates an** "event" in us; thus you have **become a quasi cause for** this wound. For instance, my feeling to **become an** actor made me to behave with such perfectionism everywhere, that people's expectations rose and when I did not come up to them I **fell**; thus the 'wound' was waiting for me and "I" was waiting for the wound! One fellow professor used to say like you are searching for ideas, ideas also searching for you. Thus the wound **possesses in itself** a nature which is "impersonal and preindividual" in character, beyond general and particular, the collective and the private. It is the question **of becoming** universalistic and holistic in your outlook. Unless this fate had not befallen you, the "**grand design**" would not have taken place in its entire entirety. It had to happen. And the concomitant ramifications and pernicious or positive **implications**. Everything is in order **because the** fate befell you. It is not as if the wound had to get something that is best from me or that I am a chosen by God to face the event. As said earlier 'the grand design' would have **been altered**. And **it cannot alter**. You got to play your part and go; there is just no other way. The legacy must go on. You shall be torch bearer and you shall hand over the torch to somebody. This is the name of the game in totalistic and holistic way.

When it comes to ethics, I would say it makes no sense if any obstreperous, obstreperous, ululations, serenading, tintinnabulations are made for the event has happened to me. It means to say that you are unworthy of the fate that has befallen you. To feel that what happened to you was unwarranted and not autonomous, telling the world that you are aggressively iconoclastic, veritably resentful, and volitionally resentful, is choosing the cast of allegation aspersions and accusations at the Grand Design. What is

immoral is to invoke the name of god, because some event has **happened to** you. Cursing him is immoral. Realize that it is all "grand design" and you are playing **a part**. Resignation, renunciation, revocation is only one form of resentment. Willing the event is primarily **to release** the eternal truth; in fact you cannot release an event despite the fact everyone tries all ways and means they pray god; they prostrate for others destitution, poverty, penury, misery. But **releasing an** event is something like an "action at a distance" which only super natural power can do.

Here we are face to face with volitional intuition and repetitive transmutation. Like a premeditated skirmisher, **one quarrel** with one self, with others, with god, and finally the accuser **leaves** this world in despair. Now look at this sentence which was quoted by I think Bousquet "if there is a **failure of** will", "I will **substitute a** longing for death" for that shall be apotheosis, a perpetual and progressive glorification of the will.

EVENT AND SINGULARITIES IN QUANTUM SYSTEMS:

What is an event? Or for that matter an ideal event? An event **is a** singularity or rather a set of singularities or set of singular points **characterizing a** mathematical curve, a physical state of affairs, a psychological person or a moral person. Singularities are turning points and points of inflection: they are bottle necks, foyers and centers; they are points of fusion; condensation and boiling; points of tears and joy; sickness and health; hope and anxiety; they are so to say "sensitive" points; such singularities should not be confused or confounded, aggravated or exacerbated with personality of a system expressing itself; or the individuality and idiosyncrasies of a system which is designated with a proposition. They should also **not be fused** with the generalizational concept or universalistic axiomatic predications and postulation alcovishness, or the dipsomaniac flageolet dirge of a concept. Possible a concept could be signified by a figurative representation or a schematic configuration. "Singularity is essentially, pre individual, and has no personalized bias in it, or for that matter a prejudice or pre circumspection of a conceptual scheme. It is in this sense **we can define a** "singularity" as being neither affirmative nor non affirmative. It can be positive or negative; it can **create or destroy**. On the other hand it must be noted that singularity is different both in its thematic discursive from the run of the mill day to day musings and mundane drooling. They are in that sense "extra-ordinary".

Each singularity is a **source and resource**, the origin, reason and raison d'être of a mathematical series, it could be any series any type, and that is interpolated or extrapolated to the structural location of the **destination of** another singularity. This according to this standpoint, there are different. It can be positive or negative; it can create or destroy. On the other hand it must be noted that singularity is different both in its thematic discursive from the run of the mill day to day musings and mundane drooling. There are in that sense "extra-ordinary".

This according to the widely held standpoint, there are different, multifarious, myriad, series IN A structure. In the eventuality of the fact that we conduct an unbiased and prudent examination of the series belonging to different "singularities" we can come to indubitable **conclusions** that the "singularity" of one system is different from the "other system" in the subterranean realm and ceratoid dualism of comparison and contrast

EPR experiment derived that there exists a communications between two particles. We go a further step to say that there **exists a channel** of communication however slovenly, inept, clumpy, between the two singularities. It is also possible the communication exchange could be one of belligerence, cantankerousness, tempestuousness, astutely truculent, with ensorcelled frenzy. That does not matter. All we are telling is that singularities communicate with each other.

Now, how do find **the reaction** of systems to these singularities. You do the same thing a boss does for you. "Problematize" the events and see how you behave. I will resort to "pressure tactics". "intimidation of deriding report", or "cut in the increment" to make you undergo trials, travails and tribulations. I am happy to see if you improve your work; but may or may not be sad if you succumb to it and hang yourself! We do the same thing with systems. systems show conducive response, felicitous reciprocation or behave erratically with inner roil, eponymous radicalism without and with blitzy conviction say like a solipsist nature of bellicose and blustering particles, or for that matter coruscation, trepidational **motion in** fluid flows, or seemingly perfidious incendiaries in gormandizing fellow elementary particles, abnormal ebullitions, surcharges calumniation and unwarranted (you think so but the system does not!) unrighteous fulminations.

So the point that is made here is "like we problematize the "events" to understand the human behaviour we have to "problematize" the events of systems to understand their behaviour.

This statement is made in connection to the fact that there shall be **creation or destruction** of particles or

complete obliteration of the system (blackhole evaporation) or obfuscation of results. Some systems are like "inside traders" they will not put signature at all! How do you find they did it! Anyway, there are possibilities of a CIA finding out as they recently did! So we can do the same thing with systems too. This is accentuation, corroboration, fortification, fomentary notes to explain the various coefficients we have used in the model as also the dissipations called for

In the Bank example we have clarified that various systems are individually conservative, and their conservativeness extends holistically too. That one law is universal does not mean there is complete adjudication of **nonexistence of** totality or global or holistic figure. Total always exists and "individual" systems always exist, if we do not bring Kant in to picture! For the time being let us not! Equations would become more eneuretic and frenzied...

Various, myriad, series in a structure. In the eventuality of the fact that we conduct an unbiased and prudent examination of the series belonging to different "singularities" we can come to indubitable conclusions that the "singularity" of one system is different from the "other system" in the subterranean realm and ceratoid dualism of comparison and contrast.

CONSERVATION LAWS:

Conservation laws bears ample testimony, infallible observatory, and impeccable demonstration to the fact that the essential predications, character constitutions, ontological consonances remain unchanged with evolution despite the system's astute truculence, serenading whimsicality, assymetric disposition or on the other hand anachronistic dispensation, eponymous radicality, entropic entrepotishness or the subdued, relationally contributive, diverse parametrizational, conducive reciprocity to environment, unconventional behaviour, eneuretic nonlinear frenetic ness, ensorcelled frenzy, abnormal ebulliations, surcharged fulminations, or the inner roil. And that holds well with the evolution with time. We present a model of the generalizational conservation of the theories. A theory of all the conservation theories. That all conservation laws hold and there is no relationship between them is bête noir. We shall on this premise build a 36 storey model that deliberates on various issues, structural, dependent, thematic and discursive,

Note THAT The classification is executed on systemic properties and parameters. And everything that is known to us measurable. We do not know "intangible". Nor we accept or acknowledge that. All laws of conservation must hold. Hence the holistic laws must hold. Towards that end, interrelationships must exist. All science like law wants evidence and here we shall provide one under the premise that for all conservations laws to hold each must be interrelated to the other, lest the very conception is a fricative contretemps. And we live in "Measurement" world.

QUANTUM REGISTER:

Devices that **harness and explore** the fundamental axiomatic predications of Physics has wide ranging amplitudinal **ramification** with its essence of locus and focus on information processing that outperforms their classical counterparts, and for unconditionally secure communication. However, in particular, implementations **based on** condensed-matter systems face the challenge of short coherence times. Carbon materials, particularly diamond, however, are suitable **for hosting** robust solid-state quantum registers, **owing to** their spin-free lattice and weak spin-orbit **coupling**. Studies with the structurally notched criticism and schizoid fragments of manifestations of historical perspective of diamond hosting quantum register have borne ample testimony and, at differential and determinate levels have articulated the generalized significations and manifestations of quantum logic elements can **be realized** by exploring long-range magnetic dipolar coupling between individually addressable single electron spins **associated with** separate colour centres in diamond. The strong distance dependence of this coupling was used to characterize the separation of single qubits ($98 \pm 3 \text{ \AA}$) with accuracy close to the value of the crystal-lattice spacing. Coherent **control over** electron spins, **conditional** dynamics, selective readout as well as switchable **interaction** should rip open glittering façade for a prosperous and

scintillating irreducible affirmation of open the way towards a viable room-temperature solid-state quantum register. As both electron spins are optically addressable, this solid-state quantum device operating at ambient conditions provides a degree of control that is at present available only for a few systems at low temperature (See for instance P. Neumann, R. Kolesov, B. Naydenov, J. Bec F. Rempp, M. Steiner V. Jacques,, G. Balasubramanian,M, M. L. Markham,, D. J. Twitchen,, S. Pezzagna,, J. Meijer, J. Twamley, F. Jelezko & J. Wrachtrup)

CAUSE AND EVENT:

MODULE NUMBERED ONE

NOTATION :

G_{13} : CATEGORY ONE OF CAUSE

G_{14} : CATEGORY TWO OF CAUSE

G_{15} : CATEGORY THREE OF CAUSE

T_{13} : CATEGORY ONE OF EVENT

T_{14} : CATEGORY TWO OF EVENT

T_{15} :CATEGORY THREE OFEVENT

FIRST TWO CATEGORIES OF QUBITS COMPUTATION:

MODULE NUMBERED TWO:

=====

===

G_{16} : CATEGORY ONE OF FIRST SET OF QUBITS

G_{17} : CATEGORY TWO OF FIRST SET OF QUBITS

G_{18} : CATEGORY THREE OF FIRST SET OF QUBITS

T_{16} :CATEGORY ONE OF SECOND SET OF QUBITS

T_{17} : CATEGORY TWO OF SECOND SET OF QUBITS

T_{18} : CATEGORY THREE OF SECOND SET OF QUBITS

THIRD SET OF QUBITS AND FOURTH SET OF QUBITS:

MODULE NUMBERED THREE:

=====

==

G_{20} : CATEGORY ONE OF THIRD SET OF QUBITS

G_{21} :CATEGORY TWO OF THIRD SET OF QUBITS

G_{22} : CATEGORY THREE OF THIRD SET OF QUBITS

T_{20} : CATEGORY ONE OF FOURTH SET OF QUBITS

T_{21} :CATEGORY TWO OF FOURTH SET OF QUBITS

T_{22} : CATEGORY THREE OF FOURTH SET OF QUBITS

FIFTH SET OF QUBITS AND SIXTH SET OF QUBITS

: MODULE NUMBERED FOUR:

=====
=

G_{24} : CATEGORY ONE OF FIFTH SET OF QUBITS

G_{25} : CATEGORY TWO OF FIFTH SET OF QUBITS

G_{26} : CATEGORY THREE OF FIFTH SET OF QUBITS

T_{24} :CATEGORY ONE OF SIXTH SET OF QUBITS

T_{25} :CATEGORY TWO OF SIXTH SET OF QUBITS

T_{26} : CATEGORY THREE OF SIXTH SET OF QUBITS

SEVENTH SET OF QUBITS AND EIGHTH SET OF QUBITS:

MODULE NUMBERED FIVE:

=====
==

G_{28} : CATEGORY ONE OF SEVENTH SET OF QUBITS

G_{29} : CATEGORY TWO OF SEVENTH SET OF QUBITS

G_{30} :CATEGORY THREE OF SEVENTH SET OF QUBITS

T_{28} :CATEGORY ONE OF EIGHTH SET OF QUBITS

T_{29} :CATEGORY TWO OF EIGHTH SET OF QUBITS

T_{30} :CATEGORY THREE OF EIGHTH SET OF QUBITS

(n-1)TH SET OF QUBITS AND nTH SET OF QUBITS :

MODULE NUMBERED SIX:

=====

- G_{32} : CATEGORY ONE OF(n-1)TH SET OF QUBITS
- G_{33} : CATEGORY TWO OF(n-1)TH SET OF QUBITS
- G_{34} : CATEGORY THREE OF (N-1)TH SET OF QUBITS
- T_{32} : CATEGORY ONE OF n TH SET OF QUBITS
- T_{33} : CATEGORY TWO OF n TH SET OF QUBITS
- T_{34} : CATEGORY THREE OF n TH SET OF QUBITS

GLOSSARY OF MODULE NUMBERED SEVEN

=====

- G_{36} : CATEGORY ONE OF TIME
- G_{37} : CATEGORY TWO OF TIME
- G_{38} : CATEGORY THREE OF TIME
- T_{36} : CATEGORY ONE OF SPACE
- T_{37} : CATEGORY TWO OF SPACE
- T_{38} : CATEGORY THREE OF SPACE

=====

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
 $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}, (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
 $(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
 $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
 $(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

are Dissipation coefficients

CAUSE AND EVENT:

MODULE NUMBERED ONE

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} \quad 2$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} \quad 3$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} \quad 4$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} \quad 5$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad 6$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad 7$$

$$+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \quad 8$$

$$-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}$$

FIRST TWO CATEGORIES OF QUBITS COMPUTATION: 9

MODULE NUMBERED TWO:

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 10$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 11$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 12$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 13$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 14$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 15$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \quad 16$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \quad 17$$

THIRD SET OF QUBITS AND FOURTH SET OF QUBITS: 18

MODULE NUMBERED THREE

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 19$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 20$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 21$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 22$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 23$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 24$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor 25

FIFTH SET OF QUBITS AND SIXTH SET OF QUBITS 26

: MODULE NUMBERED FOUR

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 27$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 28$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 29$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 30$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 31$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 32$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor** 33

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor** 34

SEVENTH SET OF QUBITS AND EIGHTH SET OF QUBITS: 35

MODULE NUMBERED FIVE

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 36$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 37$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 38$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 39$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 40$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 41$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor} \quad 42$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor} \quad 43$$

44

n-1)TH SET OF QUBITS AND nTH SET OF QUBITS :

45

MODULE NUMBERED SIX:

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 46$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 47$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 48$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 49$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 50$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 51$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor} \quad 52$$

53

SPACE AND TIME:GOVERNING EQUATIONS:

The differential system of this model is now (SEVENTH MODULE)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 54$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 55$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{13})^{(1)} \boxed{-(b''_{16})^{(1)}(G, t)} & \boxed{-(b''_{36})^{(7)}(G_{39}, t)} & \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7)}(G_{39}, t)} & & \end{array} \right] T_{13}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{ccc} (b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} & \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & & \end{array} \right] T_{14}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} & \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & & \end{array} \right] T_{15}$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$, $\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$ ARE SEVENTH DETRITION COEFFICIENTS

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} & \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{15} \quad 63$$

Where $\boxed{-(b''_{13})^{(1)}(G, t)}$, $\boxed{-(b''_{14})^{(1)}(G, t)}$, $\boxed{-(b''_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3 64

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detritions coefficients for category 1, 2 and 3

SECOND MODULE CONCATENATION:

65

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[\begin{array}{c} (a'_{16})^{(2)} \boxed{+(a'_{16})^{(2)}(T_{17}, t)} \boxed{+(a'_{13})^{(1,1)}(T_{14}, t)} \boxed{+(a'_{20})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{36})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{16} \quad 66$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[\begin{array}{c} (a'_{17})^{(2)} \boxed{+(a'_{17})^{(2)}(T_{17}, t)} \boxed{+(a'_{14})^{(1,1)}(T_{14}, t)} \boxed{+(a'_{21})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{37})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{17} \quad 67$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{c} (a'_{18})^{(2)} \boxed{+(a'_{18})^{(2)}(T_{17}, t)} \boxed{+(a'_{15})^{(1,1)}(T_{14}, t)} \boxed{+(a'_{22})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a'_{38})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{18} \quad 68$$

Where $\boxed{+(a'_{16})^{(2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2)}(T_{17}, t)}$ are first augmentation coefficients for category 1, 2 and 3 69

$\boxed{+(a'_{13})^{(1,1)}(T_{14}, t)}$, $\boxed{+(a'_{14})^{(1,1)}(T_{14}, t)}$, $\boxed{+(a'_{15})^{(1,1)}(T_{14}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{20})^{(3,3,3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3,3,3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3 70

$\boxed{+(a'_{36})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a'_{37})^{(7,7,7)}(T_{37}, t)}$, $\boxed{+(a'_{38})^{(7,7,7)}(T_{37}, t)}$ ARE SEVENTH DETRITION COEFFICIENTS 71

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{c} (b'_{16})^{(2)} \boxed{-(b'_{16})^{(2)}(G_{19}, t)} \boxed{-(b'_{13})^{(1,1)}(G, t)} \boxed{-(b'_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b'_{28})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b'_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{36})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{16} \quad 72$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{c} (b'_{17})^{(2)} \boxed{-(b'_{17})^{(2)}(G_{19}, t)} \boxed{-(b'_{14})^{(1,1)}(G, t)} \boxed{-(b'_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b'_{29})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b'_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{37})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{17} \quad 73$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{c} (b'_{18})^{(2)} \boxed{-(b'_{18})^{(2)}(G_{19}, t)} \boxed{-(b'_{15})^{(1,1)}(G, t)} \boxed{-(b'_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b'_{30})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b'_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{38})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{18} \quad 74$$

where $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3 75

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$, $\boxed{-(b'_{14})^{(1,1)}(G, t)}$, $\boxed{-(b'_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$, $-(b''_{21})^{(3,3,3)}(G_{23}, t)$, $-(b''_{22})^{(3,3,3)}(G_{23}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$ are fourth detritions coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$ are fifth detritions coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$ are sixth detritions coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$, $-(b''_{36})^{(7,7)}(G_{39}, t)$ are seventh detrition coefficients

THIRD MODULE CONCATENATION:

$$\frac{dG_{20}}{dt} = \tag{76}$$

$$(a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a'_{16})^{(2,2,2)}(T_{17}, t) & + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) \end{array} \right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a'_{17})^{(2,2,2)}(T_{17}, t) & + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) \end{array} \right] G_{21} \tag{77}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) \end{array} \right] G_{22} \tag{78}$$

79

$+(a''_{20})^{(3)}(T_{21}, t)$, $+(a''_{21})^{(3)}(T_{21}, t)$, $+(a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t)$, $+(a''_{17})^{(2,2,2)}(T_{17}, t)$, $+(a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t)$, $+(a''_{14})^{(1,1,1)}(T_{14}, t)$, $+(a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

80

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$, $+(a''_{37})^{(7,7,7)}(T_{37}, t)$, $+(a''_{38})^{(7,7,7)}(T_{37}, t)$ are seventh augmentation coefficient

81

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} \boxed{-(b''_{20})^{(3)}(G_{23}, t)} & \boxed{-(b''_{36})^{(7,7,7)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ & \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} & \end{array} \right] T_{20} \quad 82$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} \boxed{-(b''_{21})^{(3)}(G_{23}, t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ & \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} & \end{array} \right] T_{21} \quad 83$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{22})^{(3)} \boxed{-(b''_{22})^{(3)}(G_{23}, t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ & \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \end{array} \right] T_{22} \quad 84$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3)}(G_{23}, t)}$ are first detritions coefficients for category 1, 2 and 3 85

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$ are second detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1)}(G, t)}, \boxed{-(b''_{15})^{(1,1,1)}(G, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}, \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$ are seventh detritions coefficients 86

=====

FOURTH MODULE CONCATENATION:

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} \boxed{+(a''_{24})^{(4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)} \\ & \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} & \end{array} \right] G_{24} \quad 87$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} \boxed{+(a''_{25})^{(4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)} \\ & \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} & \end{array} \right] G_{25} \quad 88$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ & & + (a''_{38})^{(7,7,7,7)}(T_{37}, t) \end{array} \right] G_{26} \quad 89$$

90

Where $(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{28})^{(5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{13})^{(1,1,1,1)}(T_{14}, t)$, $(a'_{14})^{(1,1,1,1)}(T_{14}, t)$, $(a'_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$(a'_{16})^{(2,2,2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2,2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$(a'_{20})^{(3,3,3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3,3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, and 3

$(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7,7,7)}(T_{37}, t)$, $(a'_{36})^{(7,7,7,7)}(T_{37}, t)$ ARE SEVENTH augmentation coefficients

92

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ & & - (b''_{36})^{(7,7,7,7)}(G_{39}, t) \end{array} \right] T_{24} \quad 93$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ & & - (b''_{37})^{(7,7,7,7)}(G_{39}, t) \end{array} \right] T_{25} \quad 94$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ & & - (b''_{38})^{(7,7,7,7)}(G_{39}, t) \end{array} \right] T_{26} \quad 95$$

Where $-(b''_{24})^{(4)}(G_{27}, t)$, $-(b''_{25})^{(4)}(G_{27}, t)$, $-(b''_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$, $-(b''_{29})^{(5,5)}(G_{31}, t)$, $-(b''_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$, $-(b''_{33})^{(6,6)}(G_{35}, t)$, $-(b''_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$, $-(b''_{14})^{(1,1,1,1)}(G, t)$, $-(b''_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$

are fifth detrition coefficients for category 1,2 and 3

$$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)}$$

are sixth detrition coefficients for category 1,2 and 3

$$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)}$$
 ARE SEVENTH DETRITION

COEFFICIENTS

97

FIFTH MODULE CONCATENATION:

98

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)} G_{29} - \left[\begin{array}{c} \boxed{(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)} \quad \boxed{+(a''_{24})^{(4,4)}(T_{25}, t)} \quad \boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{28} \quad 99$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)} G_{28} - \left[\begin{array}{c} \boxed{(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)} \quad \boxed{+(a''_{25})^{(4,4)}(T_{25}, t)} \quad \boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{29} \quad 100$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)} G_{29} - \left[\begin{array}{c} \boxed{(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)} \quad \boxed{+(a''_{26})^{(4,4)}(T_{25}, t)} \quad \boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{30} \quad 101$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5)}(T_{29}, t)}$ are first augmentation coefficients for category 1,2 and 3 102

And $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$ are second augmentation coefficient for category 1,2 and 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$ are third augmentation coefficient for category 1,2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$ are fifth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$ are sixth augmentation coefficients for category 1,2, 3

103

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)} T_{29} - \left[\begin{array}{c} \boxed{(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{24})^{(4,4)}(G_{23}, t)} \quad \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{28} \quad 104$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)} T_{28} - \left[\begin{array}{c} \boxed{(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} \quad \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{29} \quad 105$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)} \boxed{-(b''_{30})^{(5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ & & \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{30} \quad 106$$

where $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$ are first detrition coefficients for category 1, 2 and 3 107

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$ are sixth detrition coefficients for category 1, 2, and 3

SIXTH MODULE CONCATENATION

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} \boxed{+(a''_{32})^{(6)}(T_{33}, t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)} & \boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)} \\ & & \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{32} \quad 109$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} \boxed{+(a''_{33})^{(6)}(T_{33}, t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)} & \boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)} \\ & & \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{33} \quad 110$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} \boxed{+(a''_{34})^{(6)}(T_{33}, t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)} & \boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)} \\ & & \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{34} \quad 111$$

$\boxed{+(a''_{32})^{(6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33}, t)}$ are first augmentation coefficients for category 1, 2 and 3 112

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$, $\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ARE SEVENTH

AUGMENTATION COEFFICIENTS

113

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{cccc} (b'_{32})^{(6)} & -(b''_{32})^{(6)}(G_{35}, t) & -(b''_{28})^{(5,5,5)}(G_{31}, t) & -(b''_{24})^{(4,4,4)}(G_{27}, t) \\ -(b''_{13})^{(1,1,1,1,1,1)}(G, t) & -(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & & \end{array} \right] T_{32} \quad 114$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{cccc} (b'_{33})^{(6)} & -(b''_{33})^{(6)}(G_{35}, t) & -(b''_{29})^{(5,5,5)}(G_{31}, t) & -(b''_{25})^{(4,4,4)}(G_{27}, t) \\ -(b''_{14})^{(1,1,1,1,1,1)}(G, t) & -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & & \end{array} \right] T_{33} \quad 115$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{cccc} (b'_{34})^{(6)} & -(b''_{34})^{(6)}(G_{35}, t) & -(b''_{30})^{(5,5,5)}(G_{31}, t) & -(b''_{26})^{(4,4,4)}(G_{27}, t) \\ -(b''_{15})^{(1,1,1,1,1,1)}(G, t) & -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) & \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & & \end{array} \right] T_{34} \quad 116$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients for category 1, 2 and 3 117

$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$ ARE SEVENTH DETRITION COEFFICIENTS

118

SEVENTH MODULE CONCATENATION:

119

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[\begin{array}{cccc} (a'_{36})^{(7)} & + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(7)}(T_{17}, t) & + (a''_{20})^{(7)}(T_{21}, t) & + (a''_{24})^{(7)}(T_{23}, t)G_{36} & + \\ (a''_{28})^{(7)}(T_{29}, t) & + (a''_{32})^{(7)}(T_{33}, t) & + (a''_{13})^{(7)}(T_{14}, t) & & & \end{array} \right] G_{36} \quad 120$$

121

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[\begin{array}{cccc} (a'_{37})^{(7)} & + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{14})^{(7)}(T_{14}, t) & + (a''_{21})^{(7)}(T_{21}, t) & + \\ (a''_{17})^{(7)}(T_{17}, t) & + (a''_{25})^{(7)}(T_{25}, t) & + (a''_{33})^{(7)}(T_{33}, t) & + (a''_{29})^{(7)}(T_{29}, t) & \end{array} \right] G_{37} \quad 122$$

$$\frac{dG_{38}}{dt} = \tag{123}$$

$$(a_{38})^{(7)}G_{37} - \tag{124}$$

$$\left[\begin{aligned} & \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) \right] + \underbrace{\left[(a''_{15})^{(7)}(T_{14}, t) \right]}_{\dots} + \left[(a''_{22})^{(7)}(T_{21}, t) + (a''_{18})^{(7)}(T_{17}, t) \right] + \\ & \left[(a''_{26})^{(7)}(T_{25}, t) + (a''_{34})^{(7)}(T_{33}, t) + (a''_{30})^{(7)}(T_{29}, t) \right] \end{aligned} \right] G_{38} \tag{125}$$

$$\frac{dT_{36}}{dt} = \tag{126}$$

$$\begin{aligned} & (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) \right] - (b''_{16})^{(7)}((G_{19}), t) - (b''_{13})^{(7)}((G_{14}), t) - \\ & (b''_{20})^{(7)}((G_{231}), t) - (b''_{24})^{(7)}((G_{27}), t) - (b''_{28})^{(7)}((G_{31}), t) - \\ & (b''_{32})^{(7)}((G_{35}), t) \end{aligned} \tag{126}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \tag{127}$$

$$\begin{aligned} & \left[(b'_{36})^{(7)} - (b''_{37})^{(7)}((G_{39}), t) \right] - (b''_{17})^{(7)}((G_{19}), t) - (b''_{19})^{(7)}((G_{14}), t) - \\ & (b''_{21})^{(7)}((G_{231}), t) - (b''_{25})^{(7)}((G_{27}), t) - (b''_{29})^{(7)}((G_{31}), t) - \\ & (b''_{33})^{(7)}((G_{35}), t) \end{aligned} \tag{127}$$

Where we suppose

(A) $(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0,$
 $i, j = 13, 14, 15$

(B) The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

(C) $\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)}$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants

and $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, T)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

(D) $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

(E) There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{dT_{38}}{dt} = \left[(b_{38})^{(7)}T_{37} - \left[(b_{38})^{(7)} - \boxed{(b_{38}'')^{(7)}((G_{39}), t)} - \boxed{(b_{18}'')^{(7)}((G_{19}), t)} - \boxed{(b_{20}'')^{(7)}((G_{14}), t)} - \boxed{(b_{22}'')^{(7)}((G_{23}), t)} - \boxed{(b_{26}'')^{(7)}((G_{27}), t)} - \boxed{(b_{30}'')^{(7)}((G_{31}), t)} - \boxed{(b_{34}'')^{(7)}((G_{35}), t)} \right] T_{38} \right] \quad 128$$

$+ (a_{36}'')^{(7)}(T_{37}, t) = \text{First augmentation factor}$ 134

(1) $(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$ 135

(F) (2) The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded. 136

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$: 137

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 138$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 139$$

(G) (3) $\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)}$ 140

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 141$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 142

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16, 17, 18$

They satisfy Lipschitz condition: 143

$$|(a_i'')^{(2)}(T_{17}, t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 144$$

$$|(b_i'')^{(2)}((G_{19}), t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t} \quad 145$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}, t)$ and $(a_i'')^{(2)}(T_{17}, t) \cdot (T_{17}, t)$. And (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the SECOND augmentation coefficient would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$: 147

(H) (4) $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 148

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$: 149

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 150$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 151$$

Where we suppose 152

$$(I) \quad (5) \quad (a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 153$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 154$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)} \quad 155$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$: 156

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 157

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T'_{21} - T_{21}| e^{-(\hat{M}_{20})^{(3)}t} \quad 158$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} \|G'_{23} - G_{23}\| e^{-(\hat{M}_{20})^{(3)}t} \quad 159$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the THIRD augmentation coefficient, would be absolutely continuous. 160

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 161

(J) (6) $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants 162

$(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22,$ 163

satisfy the inequalities 164

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \quad 165$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \quad 166$$

167

Where we suppose 168

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 169$$

(L) (7) The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

170

$$(M) \quad (8) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition:

171

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} |(G_{27})' - (G_{27})| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(4)}(T'_{25}, t)$ and $(a''_i)^{(4)}(T_{25}, t)$. (T'_{25}, t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a''_i)^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 4$ then the function $(a''_i)^{(4)}(T_{25}, t)$, the **FOURTH augmentation coefficient WOULD** be absolutely continuous.

172

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$:

173

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

174

(P) (9) There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose 176

(S) $(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30$ 177

(R) (10) The functions $(a''_i)^{(5)}, (b''_i)^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

178

(S) (11) $\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)}$
 $\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition: 179

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(5)}(T'_{29}, t)$ and $(a''_i)^{(5)}(T_{29}, t)$. (T'_{29}, t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a''_i)^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 5$ then the function $(a''_i)^{(5)}(T_{29}, t)$, the FIFTH **augmentation coefficient** attributable would be absolutely continuous. 180

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 181

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants
 $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 182

(S) There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose 183

$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34$ 184

(12) The functions $(a''_i)^{(6)}, (b''_i)^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

185

$$(13) \lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

186

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} \|(G_{35})' - (G_{35})\| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T'_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T'_{33}, t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 6$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the SIXTH **augmentation coefficient** would be absolutely continuous.

187

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

188

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

189

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

190

Theorem 1: if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

191

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

192

193

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

194

195

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

Definition of $G_i(0), T_i(0)$:

196

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

197

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

198

Definition of $G_i(0), T_i(0)$:

199

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 200

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 201$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 202$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 203$$

By 204

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13}(s_{(13)}, s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)} \quad 205$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)} \quad 206$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)} \quad 207$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)} \quad 208$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)} \quad 209$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

210

if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$$

By

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

211

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)}, \quad 212$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad 213$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad 214$$

By 215

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16}(s_{(16)}, s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + a''_{17}(s_{(16)}, s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)} \quad 216$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + a''_{18}(s_{(16)}, s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)} \quad 217$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G(s_{(16)}, s_{(16)})) \right) T_{16}(s_{(16)}) \right] ds_{(16)} \quad 218$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G(s_{(16)}, s_{(16)})) \right) T_{17}(s_{(16)}) \right] ds_{(16)} \quad 219$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G(s_{(16)}, s_{(16)})) \right) T_{18}(s_{(16)}) \right] ds_{(16)} \quad 220$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

221

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)}, \quad 222$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad 223$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad 224$$

By 225

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20}(s_{(20)}, s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + a''_{21}(s_{(20)}, s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)} \quad 226$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + a''_{22}(s_{(20)}, s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)} \quad 227$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G(s_{(20)}, s_{(20)})) \right) T_{20}(s_{(20)}) \right] ds_{(20)} \quad 228$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G(s_{(20)}, s_{(20)})) \right) T_{21}(s_{(20)}) \right] ds_{(20)} \quad 229$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)} \quad 230$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 231

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)}, \quad 232$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad 233$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} \quad 234$$

By 235

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)} \quad 236$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)} \quad 237$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)} \quad 238$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)} \quad 239$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)} \quad 240$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 241

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)}, \quad 242$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad 243$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad 244$$

By 245

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)} \quad 247$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)} \quad 248$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)} \quad 249$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)} \quad 250$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)} \quad 251$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

252

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i \leq (\hat{P}_{32})^{(6)}, T_i \leq (\hat{Q}_{32})^{(6)}, \quad 253$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad 254$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad 255$$

By 256

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)} \quad 257$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)} \quad 258$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)} \quad 259$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)} \quad 260$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)} \quad 261$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

262

(a) The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 263
 .Indeed it is obvious that

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)}t} - 1 \right)$$

From which it follows that 264

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)}t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$ 265

(b) The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 266
 .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = (1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 267$$

From which it follows that 268

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$ 269

(a) The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 270
 .Indeed it is obvious that

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} = (1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 271

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0 \right) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$ 272

(b) The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 273
 .Indeed it is obvious that

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} = (1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 274

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0 \right) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

(c) The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 275
 .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} = (1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that 276

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{((\hat{P}_{28})^{(5)} + G_{29}^0)}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

(d) The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 277
 .Indeed it is obvious that

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)}t \right) G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that 278

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{((\hat{P}_{32})^{(6)} + G_{33}^0)}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem1

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

279

280

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 281

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have 282

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0 \right) e^{-\frac{((\hat{P}_{13})^{(1)} + G_j^0)}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 283$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[((\hat{Q}_{13})^{(1)} + T_j^0) e^{-\frac{((\hat{Q}_{13})^{(1)} + T_j^0)}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 284$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying GLOBAL EQUATIONS into itself 285

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 286

$$d \left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote 287

Definition of \tilde{G}, \tilde{T} :

$$(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a'_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) - (a'_{13})^{(1)}(T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} &\leq \frac{1}{(\bar{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \end{aligned} \quad 288$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a'_{13})^{(1)}$ and $(b'_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ and $(\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ . 289

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(1)}$ and $(b'_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 290

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0 \quad 291$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\bar{M}_{13})^{(1)})_1$, and $((\bar{M}_{13})^{(1)})_3$: 292

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < ((\bar{M}_{13})^{(1)}) \text{ it follows } \frac{dG_{14}}{dt} \leq ((\bar{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\bar{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\bar{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\bar{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\bar{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 293

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$ then $T_{14} \rightarrow \infty$. 294

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14}')^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to 295

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$$

If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

296

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

297

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

298

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

299

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying

300

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

301

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote

302

Definition of $\widetilde{G}_{19}, \widetilde{T}_{19}$: $(\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

303

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a_{16}')^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a_{16}'')^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \end{aligned}$$

$$G_{16}^{(2)} |(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)})| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 304

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)}t} \leq & \quad 305 \\ \frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + & \\ (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) & \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 306

Remark 1: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as 307
 not conformal with the reality, however we have put this hypothesis, in order that we can postulate
 condition necessary to prove the uniqueness of the solution bounded by
 $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ .

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it
 suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on
 (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 308

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 309

Remark 3: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < (\widehat{M}_{16})^{(2)} \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)} \quad 310$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 4: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is 311
 analogous with the preceding one. An analogous property is true if G_{17} is bounded from below.

Remark 5: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then 312
 $T_{17} \rightarrow \infty$.

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 313

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 314
 unbounded. The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b'_{18})^{(2)}((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

315

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 316

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[(\widehat{P}_{20})^{(3)} + ((\widehat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{20})^{(3)} \quad 317$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[((\widehat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{20})^{(3)} \right] \leq (\widehat{Q}_{20})^{(3)} \quad 318$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i into itself 319

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 320

$$d \left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 321

$$\underline{\text{Definition of } \widetilde{G}_{23}, \widetilde{T}_{23} : ((\widetilde{G}_{23}), (\widetilde{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))}$$

It results 322

$$\begin{aligned} |\widetilde{G}_{20}^{(1)} - \widetilde{G}_{20}^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} \} ds_{(20)} \end{aligned} \quad 323$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{20})^{(3)} t} &\leq \\ \frac{1}{(\widehat{M}_{20})^{(3)}} & \left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + \right. \end{aligned} \quad 324$$

$$(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}d\left(\left((G_{23})^{(1)}, (T_{23})^{(1)}; (G_{23})^{(2)}, (T_{23})^{(2)}\right)\right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a'_{20})^{(3)}$ and $(b'_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ and $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ . 325

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on (G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 326

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 327

Remark 3: if G_{20} is bounded, the same property have also G_{21} and G_{22} . indeed if

$$G_{20} < ((\widehat{M}_{20})^{(3)})_1 \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 4: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 328

Remark 5: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$ then $T_{21} \rightarrow \infty$. 329

Definition of $(m)^{(3)}$ and ε_3 : 330

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b'_i)^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 331

$$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t \rightarrow \infty} ((b'_{22})^{(3)}((G_{23})(t), t)) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

332

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$ and to choose

333

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[(\hat{P}_{24})^{(4)} + ((\hat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\hat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{24})^{(4)}$$

334

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[((\hat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)} \right] \leq (\hat{Q}_{24})^{(4)}$$

335

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying IN to itself

336

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric

337

$$d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

$$\underline{\text{Definition of}} \left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) : \left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}\left(\left(G_{27}\right), \left(T_{27}\right)\right)$$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_{24}^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

338

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \\ \frac{1}{(\bar{M}_{24})^{(4)}} \left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\hat{A}_{24})^{(4)} + \right. \\ &\left. (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)} \right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right); \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) \end{aligned}$$

339

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$ and $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 340

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 341

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1$, $((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 342

Remark 3: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < ((\widehat{M}_{24})^{(4)})_1 \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 4: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 343

Remark 5: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b''_i)^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 344

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b''_i)^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 345

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is}$$

unbounded. The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions ANALOGOUS inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

346

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} < 1$ and to choose 347

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[(\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \quad 348$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 349$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i into itself 350

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 351

$$d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

Definition of $(\widehat{G_{31}}, \widehat{T_{31}}) : (\widehat{G_{31}}, \widehat{T_{31}}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{-(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

352

$$|(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} \leq \quad 353$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}} \left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)} \right) d \left(((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (35,35,36) the result follows

Remark 1: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$ and $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ . 354

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 355

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1$, $((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 356

Remark 3: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)} ((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 4: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 357

Remark 5: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(5)} ((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 358

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)} ((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)} \quad \text{359}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 360

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$$

If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5}$$

By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

361

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} < 1$ and to choose

362

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{32})^{(6)}$$

363

$$\frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)}$$

364

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i into itself 365

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 366

$$d\left((G_{35})^{(1)}, (T_{35})^{(1)}, (G_{35})^{(2)}, (T_{35})^{(2)} \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

$$\underline{\text{Definition of}} \quad (\widehat{G_{35}}, \widehat{T_{35}}) : \quad (\widehat{G_{35}}, \widehat{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$$

It results

$$|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} ds_{(32)} +$$

$$\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{-(\bar{M}_{32})^{(6)}s_{(32)}} +$$

$$(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} +$$

$$G_{32}^{(2)} | (a'_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a'_{32})^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$ 367

From the hypotheses it follows

$$(1) \quad (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \\ i, j = 13, 14, 15$$

(2) The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$(3) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \\ \lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants
 and $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T'_{14}, t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$:

(V) $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$:

(W) There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$|(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\hat{M}_{32})^{(6)}t} \leq$$

$$\frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d \left(((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)}) \right)$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

NOTE: SIMILAR ANALYSIS FOLLOWS FOR MODULE SEVEN

Remark 1: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ and $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ . 369

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 370

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1$, $((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 371

Remark 3: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 4: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 372

Remark 5: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \rightarrow \infty$. 373

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

374

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to

375

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b_{34}'')^{(6)}((G_{35})(t), t(t), t) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

- (e) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying 37,35,36 into itself .Indeed it is obvious that 376

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$\left(1 + (a_{36})^{(7)} t \right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

377

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[((\hat{P}_{36})^{(7)} + G_{37}^0) e^{-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$

378

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose

$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[(\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{36})^{(7)}$$

379

380

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying 381
 GLOBAL EQUATIONS AND ITS CONCOMITANT CONDITIONALITIES into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric

$$d\left((G_{39})^{(1)}, (T_{39})^{(1)}, (G_{39})^{(2)}, (T_{39})^{(2)} \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

Definition of $(\widetilde{G}_{39}), (\widetilde{T}_{39})$:

$$((\widetilde{G}_{39}), (\widetilde{T}_{39})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$$

It results

$$|\widetilde{G}_{36}^{(1)} - \widetilde{G}_{36}^{(2)}| \leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} +$$

$$\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} +$$

$$(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} +$$

$$G_{36}^{(2)} | (a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} & |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} \leq \\ & \frac{1}{(\bar{M}_{36})^{(7)}} \left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\hat{A}_{36})^{(7)} + \right. \\ & \left. (\hat{P}_{36})^{(7)} (\hat{K}_{36})^{(7)} \right) d \left(((G_{39})^{(1)}, (T_{39})^{(1)}); (G_{39})^{(2)}, (T_{39})^{(2)} \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (37,35,36) the result follows

Remark 1: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ and $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$ respectively of \mathbb{R}_+ . 385

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(7)}$ and $(b'_i)^{(7)}$, $i = 36, 37, 38$ depend only on T_{37} and respectively on (G_{39}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$

From CONCATENATED GLOBAL EQUATIONS it results

$$\begin{aligned} G_i(t) & \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0 \\ T_i(t) & \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \quad \text{for } t > 0 \end{aligned}$$

Definition of $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$ and $((\bar{M}_{36})^{(7)})_3$: 387

Remark 3: if G_{36} is bounded, the same property have also G_{37} and G_{38} . indeed if

$$G_{36} < ((\bar{M}_{36})^{(7)}) \text{ it follows } \frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\bar{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)} ((\bar{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\bar{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)} ((\bar{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If G_{37} or G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 7: If G_{36} is bounded, from below, the same property holds for G_{37} and G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 388

Remark 5: If T_{36} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \rightarrow \infty$. 389

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 390

$$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2\varepsilon_7} \right)$, $t = \log \frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t \rightarrow \infty} ((b_{38}'')^{(7)}((G_{39})(t), t)) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

$$-(\sigma_2)^{(2)} \leq -(a_{16}')^{(2)} + (a_{17}')^{(2)} - (a_{16}'')^{(2)}(T_{17}, t) + (a_{17}'')^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} \quad 391$$

$$-(\tau_2)^{(2)} \leq -(b_{16}')^{(2)} + (b_{17}')^{(2)} - (b_{16}'')^{(2)}((G_{19}), t) - (b_{17}'')^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 392$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 393

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 394

$$(a) \text{ of the equations } (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0 \quad 395$$

$$\text{and } (b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \text{ and} \quad 396$$

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 397

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 398

$$\text{roots of the equations } (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0 \quad 399$$

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 400$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 401

(b) If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 402

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 403$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 404$$

and $\boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 405$$

and analogously 406

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 407$$

Then the solution satisfies the inequalities 408

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined 409

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 410$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 411$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 412$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 413$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 414$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 415

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 416

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 417$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

418

Behavior of the solutions

419

If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

(a) $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$:

420

(b) By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:-

421

(c) If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

422

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution satisfies the inequalities

$$G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(s_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined

423

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(s_1)^{(3)}t}$$

424

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((s_1)^{(3)} - (p_{20})^{(3)} - (s_2)^{(3)})} \right) \left[e^{((s_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(s_2)^{(3)}t} \right] + G_{22}^0 e^{-(s_2)^{(3)}t} \leq G_{22}(t) \leq$$

425

$$\frac{(a_{22})^{(3)}G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})} [e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 426$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 427$$

$$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})} [e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 428$$

$$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} [e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 429

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

430

431

432

If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

(d) $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 433

(e) By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 434

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ 435

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

(f) If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

and
$$(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$

and analogously

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$

and
$$(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$ are defined respectively

Then the solution satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions

If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

(g) $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 455

(h) By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 456

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

(i) If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 457

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}$, if $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$ are defined respectively

Then the solution satisfies the inequalities 458

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 459$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 460$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((s_1)^{(5)} - (a'_{30})^{(5)})} \right) \left[e^{(s_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \quad 461$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 462$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 463$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 464$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})} \left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 465

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions 466

If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

(j) $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 467

(k) By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 468

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

(l) If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

470

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{a_{32}^0}{a_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

471

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$ are defined respectively

Then the solution satisfies the inequalities

472

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 473$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \right) \leq G_{34}(t) \leq \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \quad 474$$

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 475$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 476$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 477$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 478

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

479

If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$$(m) \quad (\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)} \text{ four constants satisfying}$$

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 480

(n) By $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$ and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and 481

Definition of $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 482

By $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the

roots of the equations $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

(o) If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 483

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

are defined respectively

Then the solution satisfies the inequalities 484

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined

485

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 486$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \right) \leq G_{38}(t) \leq \quad 487$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 488$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 489$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 490$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 491

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

From GLOBAL EQUATIONS we obtain

492

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - \\ (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:-
$$v^{(7)} = \frac{G_{36}}{G_{37}}$$

It follows

$$- \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq \\ - \left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

(a) For $0 < \left(v_0 \right)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}, \quad (C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

493

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

(b) If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$ we find like in the previous case,

494

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (\bar{v}_1)^{(7)}$$

(c) If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain

495

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)} ((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}) t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a'_{36})^{(7)} = (a'_{37})^{(7)}$, then $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$ if in addition $(v_0)^{(7)} = (v_1)^{(7)}$ then $v^{(7)}(t) = (v_0)^{(7)}$ and as a consequence $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$ **this also defines $(v_0)^{(7)}$ for the special case .**

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}$, then $(\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then

$(u_1)^{(7)} = (\bar{u}_1)^{(7)}$ if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(v_1)^{(7)}$ and $(\bar{v}_1)^{(7)}$, **and definition of $(u_0)^{(7)}$.**

We can prove the following 496

If $(a''_i)^{(7)}$ and $(b''_i)^{(7)}$ are independent on t , and the conditions 496A

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0 \quad \text{496B}$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0 \quad \text{496C}$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0, \quad \text{497C}$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0, \quad \text{497D}$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0 \quad \text{497E}$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0 \quad \text{497F}$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0 \quad \text{497G}$$

with $(p_{36})^{(7)}, (r_{37})^{(7)}$ as defined are satisfied, then the system WITH THE SATISFACTION OF THE FOLLOWING PROPERTIES HAS A SOLUTION AS DERIVED BELOW.

. 497

Particular case : 498

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition

$(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

499

From GLOBAL EQUATIONS we obtain

500

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-

$$v^{(3)} = \frac{G_{20}}{G_{21}}$$

501

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

502

From which one obtains

(a) For $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

503

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

(b) If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case,

504

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

(c) If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain

505

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (C)^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}(\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}]t}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}(\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}]t}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

506

: From GLOBAL EQUATIONS we obtain

507

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

508

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

$$(d) \text{ For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_0)^{(4)}]t}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_0)^{(4)}]t}}, \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

509

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

(e) If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 510

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} (v_1)^{(4)} - (v_2)^{(4)}] t}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} (v_1)^{(4)} - (v_2)^{(4)}] t}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}} \leq (\bar{v}_1)^{(4)}$$

(f) If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 511

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}} \leq (v_0)^{(4)}$$

512

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a_{24}''^{(4)}) = (a_{25}''^{(4)})$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .** 513

Analogously if $(b_{24}''^{(4)}) = (b_{25}''^{(4)})$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \right) - (a''_{29})^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

(g) For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}, \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

516

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}, \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

(h) If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

517

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}} \leq (\bar{v}_1)^{(5)}$$

(i) If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

518

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}} \leq (v_0)^{(5)}$$

519

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of $(u_0)^{(5)}$.**

we obtain

520
521

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

(j) For $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_0)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_0)^{(6)}]t}}, \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

522

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}, \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

523

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

(k) If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

524

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (\bar{v}_1)^{(6)}$$

(l) If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

525

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

526
527

We can prove the following

528

Theorem 3: If $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ are independent on t , and the conditions

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0 ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined, then the system

529

If $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ are independent on t , and the conditions

530.

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

531

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

532

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0 ,$$

533

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

534

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined are satisfied , then the system

If $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ are independent on t , and the conditions

535

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0 ,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined are satisfied, then the system

If $(a''_i)^{(4)}$ and $(b''_i)^{(4)}$ are independent on t , and the conditions 536

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined are satisfied, then the system

If $(a''_i)^{(5)}$ and $(b''_i)^{(5)}$ are independent on t , and the conditions 537

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined satisfied, then the system

If $(a''_i)^{(6)}$ and $(b''_i)^{(6)}$ are independent on t , and the conditions 538

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined are satisfied, then the system 539

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 540$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 541$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 542$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 543$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 544$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 545$$

has a unique positive solution, which is an equilibrium solution for the system 546

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 547$$

$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0$	548
$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0$	549
$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0$	550
$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0$	551
$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0$	552
has a unique positive solution , which is an equilibrium solution for	553
$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0$	554
$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0$	555
$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0$	556
$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0$	557
$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0$	558
$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0$	559
has a unique positive solution , which is an equilibrium solution	560
$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0$	561
$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0$	563
$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0$	564
$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0$	565
$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0$	566
$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0$	567
has a unique positive solution , which is an equilibrium solution for the system	568
$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0$	569
$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0$	570
$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0$	571
$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0$	572
$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0$	573

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 574$$

has a unique positive solution , which is an equilibrium solution for the system 575

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 576$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 577$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 578$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 579$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 580$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 584$$

has a unique positive solution , which is an equilibrium solution for the system 582

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 583$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 584$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 585$$

586

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 587$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 588$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 589$$

has a unique positive solution , which is an equilibrium solution for the system (79 to 36) 560

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Definition and uniqueness of T_{37}^* :-

561

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

(f) By the same argument, the equations 92,93 admit solutions G_{36}, G_{37} if

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - [(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^* such that $\varphi(G_{37}^*) = 0$

562

Finally we obtain the unique solution of 89 to 97

G_{37}^* given by $\varphi((G_{39})^*) = 0, T_{37}^*$ given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

563

Definition and uniqueness of T_{21}^* :-

564

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{21})$ being increasing, it follows

that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]} \quad 565$$

Definition and uniqueness of T_{25}^* :- 566

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]} \quad 567$$

Definition and uniqueness of T_{29}^* :- 568

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]} \quad 568$$

Definition and uniqueness of T_{33}^* :- 569

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]} \quad 569$$

(g) By the same argument, the equations 92,93 admit solutions G_{13}, G_{14} if 570

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - \\ [(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

(h) By the same argument, the equations 92,93 admit solutions G_{16}, G_{17} if 571

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - \\ [(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 571

(i) By the same argument, the concatenated equations admit solutions G_{20}, G_{21} if 572

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

573

(j) By the same argument, the equations of modules admit solutions G_{24}, G_{25} if

574

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

(k) By the same argument, the equations (modules) admit solutions G_{28}, G_{29} if

575

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

(l) By the same argument, the equations (modules) admit solutions G_{32}, G_{33} if

578

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

579

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

580

581

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

Finally we obtain the unique solution of 89 to 94

582

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

583

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 584

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad \text{585}$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad \text{586}$$

Obviously, these values represent an equilibrium solution 587

Finally we obtain the unique solution 588

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 589

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]} \quad \text{590}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 591

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^*)]} \quad \text{592}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 593

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^*)]} \quad \text{594}$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

595

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(1)}$ and $(b'_i)^{(1)}$ belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i \tag{596}$$

$$\frac{\partial(a'_{14})^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)}, \quad \frac{\partial(b'_i)^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 597

$$\frac{dG_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})G_{13} + (a_{13})^{(1)}G_{14} - (q_{13})^{(1)}G_{13}^*T_{14} \tag{598}$$

$$\frac{dG_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})G_{14} + (a_{14})^{(1)}G_{13} - (q_{14})^{(1)}G_{14}^*T_{14} \tag{599}$$

$$\frac{dG_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})G_{15} + (a_{15})^{(1)}G_{14} - (q_{15})^{(1)}G_{15}^*T_{14} \tag{600}$$

$$\frac{dT_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})T_{13} + (b_{13})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*G_j \tag{601}$$

$$\frac{dT_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})T_{14} + (b_{14})^{(1)}T_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*G_j \tag{602}$$

$$\frac{dT_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})T_{15} + (b_{15})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*G_j \tag{603}$$

If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(2)}$ and $(b'_i)^{(2)}$ belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable 604

Denote 605

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i \tag{606}$$

$$\frac{\partial(a'_{17})^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b'_i)^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \tag{607}$$

taking into account equations (global)and neglecting the terms of power 2, we obtain 608

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \tag{609}$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \tag{610}$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \tag{611}$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*G_j \tag{612}$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*G_j \tag{613}$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*G_j \quad 614$$

If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Denote

Definition of G_i, T_i :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b''_i)^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

616

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

617

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 618$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 619$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 6120$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^*G_j \quad 621$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^*G_j \quad 622$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^*G_j \quad 623$$

If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

624

Denote

Definition of G_i, T_i :-

625

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial(b''_i)^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

626

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^*T_{25} \quad 627$$

$$\frac{dG_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^*T_{25} \quad 628$$

$$\frac{dG_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^*T_{25} \quad 629$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*G_j \quad 630$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*G_j \quad 631$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*G_j \quad 632$$

633

If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Denote

Definition of G_i, T_i :- 634

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b'_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 635

$$\frac{dG_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})G_{28} + (a_{28})^{(5)}G_{29} - (q_{28})^{(5)}G_{28}^*T_{29} \quad 636$$

$$\frac{dG_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})G_{29} + (a_{29})^{(5)}G_{28} - (q_{29})^{(5)}G_{29}^*T_{29} \quad 637$$

$$\frac{dG_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^*T_{29} \quad 638$$

$$\frac{dT_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*G_j \quad 639$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*G_j \quad 640$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*G_j \quad 641$$

If the conditions of the previous theorem are satisfied and if the functions $(a'_i)^{(6)}$ and $(b'_i)^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable 642

Denote

Definition of G_i, T_i :- 643

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b'_i)^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations(global) and neglecting the terms of power 2, we obtain 644

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 645$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^* T_{33} \quad 646$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^* T_{33} \quad 647$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^* G_j \quad 648$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^* G_j \quad 649$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^* G_j \quad 650$$

Obviously, these values represent an equilibrium solution of 79,20,36,22,23, 651

If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ and $(b_i'')^{(7)}$ belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of G_i, T_i :- 652

$$G_i = G_i^* + G_i \quad , \quad T_i = T_i^* + T_i \quad 653$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}} (T_{37}^*) = (q_{37})^{(7)} \quad , \quad \frac{\partial (b_i'')^{(7)}}{\partial G_j} ((G_{39})^{**}) = s_{ij}$$

Then taking into account equations(SOLUTIONAL) and neglecting the terms of power 2, we obtain 654

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^* T_{37} \quad 655$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^* T_{37} \quad 657$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 658$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^*G_j \quad 659$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^*G_j \quad 660$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^*G_j \quad 661$$

2. 662

The characteristic equation of this system is

$$\begin{aligned} & \left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)} \right) \left\{ \left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)} \right) \right. \\ & \left[\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)} \right) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right] \\ & \left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)} \right) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^* \right) \\ & + \left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)} \right) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^* \right) \\ & \left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)} \right) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^* \right) \\ & \left(\left((\lambda)^{(1)} \right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)} \right) (\lambda)^{(1)} \right) \\ & \left(\left((\lambda)^{(1)} \right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)} \right) (\lambda)^{(1)} \right) \\ & + \left(\left((\lambda)^{(1)} \right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)} \right) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\ & + \left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)} \right) \left((a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^* \right) \\ & \left. \left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)} \right) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \right\} = 0 \\ & + \\ & \left((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)} \right) \left\{ \left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)} \right) \right. \end{aligned}$$

$$\left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right]$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^*$$

$$+ \left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)} \right) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^*$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^*$$

$$\left((\lambda)^{(2)} \right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)}$$

$$\left((\lambda)^{(2)} \right)^2 + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)}$$

$$+ \left((\lambda)^{(2)} \right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18}$$

$$+ (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \left((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right)$$

$$\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \} = 0$$

+

$$\left((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \left\{ (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \right\}$$

$$\left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right]$$

$$\left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^*$$

$$+ \left((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^*$$

$$\left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^*$$

$$\left((\lambda)^{(3)} \right)^2 + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)}$$

$$\left((\lambda)^{(3)} \right)^2 + \left((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)}$$

$$+ \left((\lambda)^{(3)} \right)^2 + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22}$$

$$+ (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \left((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^* \right)$$

$$\left(\left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right) \} = 0$$

+

$$\begin{aligned} & \left((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)} \right) \{ (\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \} \\ & \left[\left((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\ & \left((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \\ & + \left((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)} \right) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\ & \left((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \\ & \left((\lambda)^{(4)} \right)^2 + \left((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\ & \left((\lambda)^{(4)} \right)^2 + \left((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} \right) (\lambda)^{(4)} \\ & + \left((\lambda)^{(4)} \right)^2 + \left((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\ & + \left((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) \left((a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^* \right) \\ & \left((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)} \right) \{ (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \} \\ & \left[\left((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\ & \left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \\ & + \left((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)} \right) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\ & \left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \} \end{aligned}$$

$$\begin{aligned} & \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} \right) (\lambda)^{(5)} \\ & \left(((\lambda)^{(5)})^2 + (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} \right) (\lambda)^{(5)} \\ & + \left(((\lambda)^{(5)})^2 + (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} \right) (\lambda)^{(5)} (q_{30})^{(5)} G_{30} \\ & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^*) \\ & \left(\left((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)} \right) \{ (\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)} \} \\ & \left[\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\ & \left(\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \right) \\ & + \left((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)} \right) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\ & \left(\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\ & \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} \\ & \left(((\lambda)^{(6)})^2 + (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\ & + \left(((\lambda)^{(6)})^2 + (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\ & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\ & \left(\left((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \{ (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \} \\ & \left[\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) S_{(37),(37)} T_{37}^* + (b_{37})^{(7)} S_{(36),(37)} T_{37}^* \right) \\
 & + \left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)}) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \right) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) S_{(37),(36)} T_{37}^* + (b_{37})^{(7)} S_{(36),(36)} T_{36}^* \right) \\
 & \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left(((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & + \left(((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) \left((a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\
 & \left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) S_{(37),(38)} T_{37}^* + (b_{37})^{(7)} S_{(36),(38)} T_{36}^* \right) \} = 0
 \end{aligned}$$

REFERENCES

=====

(1) A HAIMOVICI: "On the growth of a two species ecological system divided on age groups". Tensor, Vol 37 (1982), Commemoration volume dedicated to Professor Akitsugu Kawaguchi on his 80th birthday

(2) FRTJOF CAPRA: "The web of life" Flamingo, Harper Collins See "Dissipative structures" pages 172-188

(3) HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [http://www.eolss.net

(4) MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204, doi:10.1029/2005JD006097

(5) STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" J. Atmos. Sci., 53, 980-1006

(6) FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010

(7) HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [http://www.eolss.net

(8) MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204,

doi:10.1029/2005JD006097

(8A) STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, “Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus” *J. Atmos. Sci.*, 53, 980-1006

(8B) FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), “Precipitation-generated oscillations in open cellular cloud fields” *Nature*, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010

-
- (9)^{a b c} Einstein, A. (1905), "Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?", *Annalen der Physik* **18**:
639 Bibcode 1905AnP...323..639E,DOI:10.1002/andp.19053231314. See also the English translation.
- (10)^{a b} Paul Allen Tipler, Ralph A. Llewellyn (2003-01), *Modern Physics*, W. H. Freeman and Company, pp. 87–88, ISBN 0-7167-4345-0
- (11)^{a b} Rainville, S. et al. World Year of Physics: A direct test of $E=mc^2$. *Nature* 438, 1096-1097 (22 December 2005) | doi: 10.1038/4381096a; Published online 21 December 2005.
- (12)[^] In F. Fernflores. The Equivalence of Mass and Energy. Stanford Encyclopedia of Philosophy
- (13)[^] Note that the relativistic mass, in contrast to the rest mass m_0 , is not a relativistic invariant, and that the velocity is not a Minkowski four-vector, in contrast to the quantity dx^μ , where dx^μ is the differential of the proper time. However, the energy-momentum four-vector is a genuine Minkowski four-vector, and the intrinsic origin of the square-root in the definition of the relativistic mass is the distinction between $d\tau$ and dt .
- (14)[^] Relativity DeMystified, D. McMahon, Mc Graw Hill (USA), 2006, ISBN 0-07-145545-0
- (15)[^] Dynamics and Relativity, J.R. Forshaw, A.G. Smith, Wiley, 2009, ISBN 978-0-470-01460-8
- (16)[^] Hans, H. S.; Puri, S. P. (2003). *Mechanics* (2 ed.). Tata McGraw-Hill. p. 433. ISBN 0-07-047360-9., Chapter 12 page 433
- (17)[^] E. F. Taylor and J. A. Wheeler, **Spacetime Physics**, W.H. Freeman and Co., NY. 1992. ISBN 0-7167-2327-1, see pp. 248-9 for discussion of mass remaining constant after detonation of nuclear bombs, until heat is allowed to escape.
- (18)[^] Mould, Richard A. (2002). *Basic relativity* (2 ed.). Springer. p. 126. ISBN 0-387-95210-1., Chapter 5 page 126
- (19)[^] Chow, Tail L. (2006). *Introduction to electromagnetic theory: a modern perspective*. Jones & Bartlett Learning. p. 392. ISBN 0-7637-3827-1., Chapter 10 page 392
- (20)[^] [2] Cockcroft-Walton experiment
- (21)^{a b c} Conversions used: 1956 International (Steam) Table (IT) values where one calorie

$\equiv 4.1868 \text{ J}$ and one BTU $\equiv 1055.05585262 \text{ J}$. Weapons designers' conversion value of one gram TNT $\equiv 1000$ calories used.

(22)[^] Assuming the dam is generating at its peak capacity of 6,809 MW.

(23)[^] Assuming a 90/10 alloy of Pt/Ir by weight, a C_p of 25.9 for Pt and 25.1 for Ir, a Pt-dominated average C_p of 25.8, 5.134 moles of metal, and 132 J.K^{-1} for the prototype. A variation of ± 1.5 picograms is of course, much smaller than the actual uncertainty in the mass of the international prototype, which are ± 2 micrograms.

(24)[^] [31] Article on Earth rotation energy. Divided by c^2 .

(25)^{^ a b} Earth's gravitational self-energy is 4.6×10^{-10} that of Earth's total mass, or 2.7 trillion metric tons. Citation: *The Apache Point Observatory Lunar Laser-Ranging Operation (APOLLO)*, T. W. Murphy, Jr. *et al.* University of Washington, Dept. of Physics (132 kB PDF, here.).

(26)[^] There is usually more than one possible way to define a field energy, because any field can be made to couple to gravity in many different ways. By general scaling arguments, the correct answer at everyday distances, which are long compared to the quantum gravity scale, should be *minimal coupling*, which means that no powers of the curvature tensor appear. Any non-minimal couplings, along with other higher order terms, are presumably only determined by a theory of quantum gravity, and within string theory, they only start to contribute to experiments at the string scale.

(27)[^] G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Physical Review D* 14:3432–3450 (1976).

(28)[^] A. Belavin, A. M. Polyakov, A. Schwarz, Yu. Tyupkin, "Pseudoparticle Solutions to Yang Mills Equations", *Physics Letters* 59B:85 (1975).

(29)[^] F. Klinkhammer, N. Manton, "A Saddle Point Solution in the Weinberg Salam Theory", *Physical Review D* 30:2212.

(30)[^] Rubakov V. A. "Monopole Catalysis of Proton Decay", *Reports on Progress in Physics* 51:189–241 (1988).

(31)[^] S.W. Hawking "Black Holes Explosions?" *Nature* 248:30 (1974).

(32)[^] Einstein, A. (1905), "Zur Elektrodynamik bewegter Körper." (PDF), *Annalen der Physik* 17: 891–921, Bibcode 1905AnP...322...891E, DOI:10.1002/andp.19053221004. English translation.

- (33)[△] See e.g. Lev B. Okun, *The concept of Mass*, Physics Today **42** (6), June 1969, p. 31–36, http://www.physicstoday.org/vol-42/iss-6/vol42no6p31_36.pdf
- (34)[△] Max Jammer (1999), *Concepts of mass in contemporary physics and philosophy*, Princeton University Press, p. 51, ISBN 0-691-01017-X
- (35)[△] Eriksen, Erik; Vøyenli, Kjell (1976), "The classical and relativistic concepts of mass", *Foundations of Physics* (Springer) **6**: 115–124, Bibcode 1976FoPh....6..115E, DOI:10.1007/BF00708670
- (36)[△] ^{a b} Janssen, M., Mecklenburg, M. (2007), *From classical to relativistic mechanics: Electromagnetic models of the electron.*, in V. F. Hendricks, et al., , *Interactions: Mathematics, Physics and Philosophy* (Dordrecht: Springer): 65–134
- (37)[△] ^{a b} Whittaker, E.T. (1951–1953), 2. Edition: *A History of the theories of aether and electricity, vol. 1: The classical theories / vol. 2: The modern theories 1900–1926*, London: Nelson
- (38)[△] Miller, Arthur I. (1981), *Albert Einstein's special theory of relativity. Emergence (1905) and early interpretation (1905–1911)*, Reading: Addison–Wesley, ISBN 0-201-04679-2
- (39)[△] ^{a b} Darrigol, O. (2005), "The Genesis of the theory of relativity." (PDF), *Séminaire Poincaré* **1**: 1–22
- (40)[△] Philip Ball (Aug 23, 2011). "Did Einstein discover $E = mc^2$?" Physics World.
- (41)[△] Ives, Herbert E. (1952), "Derivation of the mass-energy relation", *Journal of the Optical Society of America* **42** (8): 540–543, DOI:10.1364/JOSA.42.000540
- (42)[△] Jammer, Max (1961/1997). *Concepts of Mass in Classical and Modern Physics*. New York: Dover. ISBN 0-486-29998-8.
- (43)[△] Stachel, John; Torretti, Roberto (1982), "Einstein's first derivation of mass-energy equivalence", *American Journal of Physics* **50** (8): 760–763, Bibcode1982AmJPh..50..760S, DOI:10.1119/1.12764
- (44)[△] Ohanian, Hans (2008), "Did Einstein prove $E=mc^2$?", *Studies In History and Philosophy of Science Part B* **40** (2): 167–173, arXiv:0805.1400, DOI:10.1016/j.shpsb.2009.03.002

(45)^ Hecht, Eugene (2011), "How Einstein confirmed $E=mc^2$ ", *American Journal of Physics* **79** (6): 591–600, Bibcode 2011AmJPh..79..591H, DOI:10.1119/1.3549223

(46)^ Rohrlich, Fritz (1990), "An elementary derivation of $E=mc^2$ ", *American Journal of Physics* **58** (4): 348–349, Bibcode 1990AmJPh..58..348R, DOI:10.1119/1.16168

(47) (1996). *Lise Meitner: A Life in Physics*. California Studies in the History of Science. **13**. Berkeley: University of California Press. pp. 236–237. ISBN 0-520-20860-

(48)^ UIBK.ac.at

(49)^ J. J. L. Morton; *et al.* (2008). "Solid-state quantum memory using the ^{31}P nuclear spin". *Nature* **455** (7216): 1085–1088. Bibcode 2008Natur.455.1085M.DOI:10.1038/nature07295.

(50)^ S. Weisner (1983). "Conjugate coding". *Association of Computing Machinery, Special Interest Group in Algorithms and Computation Theory* **15**: 78–88.

(51)^ A. Zeilinger, *Dance of the Photons: From Einstein to Quantum Teleportation*, Farrar, Straus & Giroux, New York, 2010, pp. 189, 192, ISBN 0374239665

(52)^ B. Schumacher (1995). "Quantum coding". *Physical Review A* **51** (4): 2738–2747. Bibcode 1995PhRvA..51.2738S. DOI:10.1103/PhysRevA.51.2738.

Acknowledgments:

The introduction is a collection of information from various articles, Books, News Paper reports, Home Pages Of authors, Journal Reviews, Nature 's Letters, Article Abstracts, Research papers, Abstracts Of Research Papers, Stanford Encyclopedia, Web Pages, Ask a Physicist Column, Deliberations with Professors, the internet including Wikipedia. We acknowledge all authors who have contributed to the same. In the eventuality of the fact that there has been any act of omission on the part of the authors, we regret with great deal of compunction, contrition, regret, trepidation and remorse. As Newton said, it is only because erudite and eminent people allowed one to piggy ride on their backs; probably an attempt has been made to look slightly further. Once again, it is stated that the references are only illustrative and not comprehensive

First Author: ¹**Mr. K. N.Prasanna Kumar** has three doctorates one each in Mathematics, Economics, Political Science. Thesis was based on Mathematical Modeling. He was recently awarded D.litt. for his work on 'Mathematical Models in Political Science'--- Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Corresponding [Author:drknpkumar@gmail.com](mailto:drknpkumar@gmail.com)

Second Author: ²**Prof. B.S Kiranagi** is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150 publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent publication history.-- UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

Third Author: ³**Prof. C.S. Bagewadi** is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. Several Books on Differential Geometry, Differential Equations are coauthored by him--- Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India

=====

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

