

On The Class of (K-N) Quasi-Normal Operators On Hilbert Space

Dr. Salim D. M and Ahmed M.K.

AL-Mustansiriya University/College of Education/ Department of mathematics

Abstract

In this work, we introduce another class of normal operator which is (K-N) quasi-normal operator and given some properties of this concept as well as discussion the relation between this operator with another types of normal operators.

Keyword: (K-N) quasi-normal

1- Introduction and Terminologies

Let H be complex Hilbert space, and $B(H)$ the space of all bounded linear operator from H in to H , the quasi-normal operator was introduced at first by A. Brown [2] in 1953 and given some properties of this operator, but this concept was generalized by researchers such as D. Senthilkumar and others introduced new types of quasi-normal operator is said to be K-quasi-normal operator with relationships between these types of operators. But in 2011 O. Ahmed [5], given another class of quasi-normal operator, which is n-power quasi-normal operator.

In this search, we introduce another generalize of quasi-normal operator which (K-N)quasi-normal operator and modification the result appear in [1] and [3] about this concept. As well as, given some basic properties of this operator with relation between (K-N) quasi-normal operator and another classes of quasi normal operators.

2- Basic Concepts

Here, we recall fundamental concepts of this work and in first we give the definition of normal operator.

Definitions (2.2),[4]:

- (1) An operator $T : H \rightarrow H$ is said to be normal operator if and only if $TT^* = T^*T$.
- (2) An operator $T : H \rightarrow H$ is said to be quasi normal operator if T and T^*T are commute.

Next, we recall the generalized of normal operator by the following definition.

Definition (2.4), [4]:

An operator $T : H \rightarrow H$ is said to be n-power quasi normal operator if and only if $T^n T^* T = T^* T T^n$.

Also, we give the definition of (K-N) quasi-normal operator, this definition is generalized to definition appear in [4].

Definition (2.5):


Let T be a bounded operator from a complex Hilbert space H to it self, then T is said to be (K-N) quasi normal operator if satisfy the condition $T^K (T^*T) = N(T^*T)T^K$, where K is positive integer and N is bounded operator from a complex Hilbert space H to it self.

Next, can be introduce the relation between (K-N) quasi normal operator and other classes by the following remark.

Remarks (2.6):

- 1- Its clearly that if $K=1$, we get T is (N) quasi normal operator, and if $N=I$, we get T is n-power- quasi normal, and we get T is quasi normal if $N=I$ and $K=1$.
- 2- Every (N) quasi normal is (K-N) quasi normal

To illustrate this remarks, we will introduce the following digram.



3- Properties of (K-N) quasi normal operator

The following theorem give some properties of (K-N) quasi-normal operator.

Theorem (3.1):

Let $T \in B(H)$ is an operator if C is commutes with U and V , and $C^2T^K = NC^2T^K$ then T is (K-N) quasi normal.

Where, $B^2 = TT^*$, $C^2 = T^*T$, $U = \text{Re}T = \frac{T + T^*}{2}$ and $V = \text{Im}T = \frac{T - T^*}{2i}$

Proof:

Since $CU = UC$, $BV = VB$ so, $C^2U = UC^2$, $B^2V = VB^2$ thus $C^2U^K = U^KC^2$, $B^2V^K = V^KB^2$ then

$$C^2T^K + C^2(T^K)^* = T^KC^2 + (T^K)^*C^2$$

$$C^2T^K - C^2(T^K)^* = T^KC^2 - (T^K)^*C^2$$

This gives $T^KC^2 = C^2T^K$

$T^K (T^*T) = (T^*T)T^K$, and by using the condition $B^2T^K = NB^2T^K$ so we get:
 $T^K (T^*T) = N(T^*T)T^K$ then, T is (K-N) quasi normal.

more properties give by the following theorem.

Theorem(3.2):

If $T \in B(H)$ is an operator such that $C^2U^K = \frac{1}{N}U^K C^2$, $C^2V^K = \frac{1}{N}V^K C^2$ then T is (K-N) quasi normal.

Proof:

Since $C^2U^K = \frac{1}{N}U^K C^2$, $C^2V^K = \frac{1}{N}V^K C^2$ then we have

$$C^2(U + iV)^K = \frac{1}{N}(U + iV)^K C^2 \text{ and we have } C^2T^K = \frac{1}{N}T^K C^2 \text{ therefore;}$$

$$(T^*T)T^K = \frac{1}{N}T^K (T^*T) \text{ so, } T^K (T^*T) = N(T^*T)T^K \text{ then, } T \text{ is (K-N) quasi normal.}$$

The operation on (K-N) quasi normal have been given by the following theorem.

Theorem(3.3):

Let T_1, T_2 be two (K-N) quasi normal from H to H , such that $T_1^K T_2^* = T_2^K T_1^* = T_1^* T_2 = T_2^* T_1 = 0$ then $T_1 + T_2$ is (K-N) quasi normal.

Proof:

$$\begin{aligned} (T_1 + T_2)^K [(T_1 + T_2)^* (T_1 + T_2)] &= (T_1 + T_2)^K [(T_1^* + T_2^*)(T_1 + T_2)] \\ &= (T_1 + T_2)^K (T_1^*T_1 + T_1^*T_2 + T_2^*T_1 + T_2^*T_2) \\ &= (T_1 + T_2)^K (T_1^*T_1 + T_2^*T_2) \\ &= (T_1^K + T_2^K)(T_1^*T_1 + T_2^*T_2) \\ &= T_1^K T_1^* T_1 + T_1^K T_2^* T_2 + T_2^K T_1^* T_1 + T_2^K T_2^* T_2 \\ &= T_1^K T_1^* T_1 + T_2^K T_2^* T_2 \\ &= N((T_1 T_1^*) T_1^K) + N((T_2 T_2^*) T_2^K) \end{aligned}$$

Hence $T_1 + T_2$ is (K-N) quasi normal.

From above theorem, we can get the corollary its proof easy can be omitted it.

Corollary (3.4):

Let T_1, T_2 be two (K-N) quasi normal, such that $T_1^K T_2^* = T_2^K T_1^* = T_1^* T_2 = T_2^* T_1 = 0$ then $T_1 - T_2$ is (K-N) quasi normal.

Theorem(3.5):

Let T_1 be (K-N) quasi normal operator and T_2 (K-power) quasi normal operator. Then there product $T_1 T_2$ is (K-N) quasi normal operator if the following conditions are satisfied

- (i) $T_1 T_2 = T_2 T_1$
- (ii) $T_1 T_2^* = T_2^* T_1$

Proof:

$$\begin{aligned}
 (T_1 T_2)^K (T_1 T_2)^* (T_1 T_2) &= (T_1^K T_2^K)(T_2^* T_1^*)(T_1 T_2) \\
 &= (T_1^K T_2^K)(T_1^* T_2^*)(T_1 T_2) \\
 &= T_1^K (T_2^K T_1^*)(T_2^* T_1) T_2 \\
 &= T_1^K (T_1^* T_2^K)(T_1 T_2^*) T_2 \\
 &= T_1^K T_1^* (T_2^K T_1) T_2^* T_2 \\
 &= (T_1^K T_1^* T_1)(T_2^K T_2^* T_2) \\
 &= N((T_1^* T_1) T_1^K)(T_2^* T_2) T_2^K \\
 &= N(T_1^* T_1 (T_1^K T_2^*) T_2 T_2^K) \\
 &= N(T_1^* T_1 T_2^* T_1^K T_2 T_2^K) \\
 &= N[T_1^* (T_1 T_2^*)(T_1^K T_2) T_2^K] \\
 &= N[(T_1^* T_2^*)(T_1 T_2)(T_1^K T_2^K)] \\
 &= N[(T_2 T_1)^* (T_1 T_2)(T_1 T_2)^K] \\
 &= N[(T_1 T_2)^* (T_1 T_2)(T_1 T_2)^K]
 \end{aligned}$$

Hence, the product $T_1 T_2$ is (K-N) quasi normal operator.

4- Reference :

- [1] A. A. S. Jibril, " On n-Power Normal operators ", The journal for science and engineering, Vol.33, pp. 247-251, 2008.
- [2] A. Brown, " On a class of operators", Proc. Amer. Math. Soc, 723-728, 1953.
- [3] D. Senthilkumar and others, " k- QUASI-NORMAL OPERATORS", International journal of math. And comp., Vol. 15, issue No. 2, 2012.
- [4] E. A. Kuffi and H. A. Satar, " A new linear normal operators and it's relations", Jornal of college of education, AL-Mustansiriya university, Baghdad, Iraq,2014.
- [5] O. Ahmed, " On the class of n-Power quasi-normal operators on Hilbert space", Bulletin of math. Analysis and App., ISSN 1821-1292, Vol. 3, PP. 213-228, 2011.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

