

## ON EQUALITY OF SPECTRA AND ESSENTIAL SPECTRA OF OPERATORS IN HILBERT SPACES

S.W. LUKETERO<sup>1</sup>, P.G. RUGIRI<sup>2\*</sup> AND J.M. KHALAGAI<sup>3</sup>

SCHOOL OF MATHEMATICS, UNIVERSITY OF NAIROBI, P.O BOX 30197, 00100  
NAIROBI KENYA

### ABSTRACT:

It is a well known fact in operator theory that for any operator  $A$ , the essential spectrum of  $A$  is contained in the spectrum of  $A$ . The task of finding conditions under which the two are equal has been the subject matter in a number of several research papers. In this paper we show that quasisimilar pure dominant operators have their essential spectra equal to their spectra provided one of the intertwining quasiaffinities is compact.

AMS subject classification 47B47, 47A30, 47B20 key words and phrases: pure dominant and essential spectrum.

### 1. INTRODUCTION

Let  $H$  be a complex Hilbert space and  $B(H)$  denote the Banach algebra of all bounded linear operators on  $H$ . An operator  $A \in B(H)$  is said to be a quasiaffinity if  $A$  is both one-one and has dense range. Two operators  $A$  and  $B$  are said to be similar if there is an invertible operator  $S$  such that  $AS = SB$ , while  $A$  and  $B$  are said to be quasisimilar if there exist quasiaffinities  $X$  and  $Y$  such that  $AX = XB$  and  $BY = YA$ .

The concept of quasisimilarity and equality of spectra has been considered by a number of authors among them S. Clary [1] who showed that quasisimilar hyponormal operators have equal spectra. J.M. Khalagai and B. Nyamai [2] showed that if  $A$  and  $B$  are quasisimilar with  $A$  dominant and  $B$   $M$ -hyponormal then  $B$  have equal spectra. B.P. Duggal [3] Showed that if  $A_i$ ,  $i = 1, 2$  are quasi-similar  $P$  – hyponormal such that  $U_i$ ,  $i = 1, 2$  is unitary in the polar decomposition  $A_i = U_i |A_i|$  then  $A_1$  and  $A_2$  have not only same spectra but also same essential spectra.

The Problem of looking for conditions under which the essential spectrum is equal to spectrum of a given operator has also been considered by a number of authors. In particular J.P. Williams [5] apart from showing that there are several cases under which quasisimilar operators  $A$  and  $B$  have equal essential spectra also proved the following result on equality of spectrum and essential spectrum of a given operator.

### Theorem A [5]

Suppose that  $T$  is a pure dominant operator,  $K$  is a compact operator having dense range and  $KT = TK$ . Then spectrum of  $T$  is equal to essential spectrum of  $T$ .

It is at this point that we also pick up the quest of delving into this theory in this paper.

### Definition 1

Let  $x \in \mathcal{H}$ . We define  $\rho_T(x)$  to be the set of complex numbers  $\alpha$  for which there exists a neighbourhood  $V_\alpha$  of  $\alpha$  with  $u$  analytic on having values in  $H$  such that  $(zI - T)u(z) = x$  on  $V_\alpha$ .

This set is open and contains the complement of the spectrum  $\sigma(T)$  of  $T$ . The function  $u$  is called a local resolvent of  $T$  on  $V_\alpha$ . By definition the local spectrum of  $T$  at  $x$  denoted by  $\sigma(T, x)$  is the complement of  $\rho_T(x)$  and so is a compact subset of  $\sigma(T)$ . We say that  $T$  has the single valued extension property (in short SVEP) if  $(zI - T)u(z) = 0$  implies  $u = 0$  for any analytic function  $u$  defined on any domain  $D$  of a complex plane with values in  $H$ .

An operator  $T \in H$  is said to satisfy Dunford's property (C) if for each closed subset  $F$  of the complex plane the corresponding local spectrum subspace  $H_T(F) = \{x \in H : \sigma(T, x) \subset F\}$  is closed.

### Theorem B [5]

Suppose  $A$  and  $B$  are dominant operators satisfying Dunford's property (C) and are quasisimilar with at least one of the implementing quasiaffinities compact, then  $A$  and  $B$  have equal spectra and also equal essential spectra.

## 2. NOTATION AND TERMINOLOGY

Given an operator  $A \in B(H)$  the spectrum of  $A$  is denoted by  $\sigma(A)$ .

Thus  $\sigma(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is not invertible}\}$  where  $\mathbb{C}$  is the complex number field. The commutator of  $A$  and  $B$  is denoted by  $[A, B]$  where  $[A, B] = AB - BA$ .

We denote range of  $A$  and Kernel of  $A$  by  $\text{ran } A$  and  $\text{Ker } A$ . An operator  $A$  is said to be dominant if to each  $\lambda \in \mathbb{C}$  there corresponds a number  $M_\lambda \geq 1$  such that  $\|(A - \lambda)^* x\| \leq M_\lambda \|(\lambda - A)x\| \forall x \in H$ . But  $A$  is said to be pure dominant if there exists no non-trivial reducing subspace of  $A$  on which the restriction of  $A$  is normal.  $A$  is  $M$ -hyponormal if  $\exists M > M_\lambda$  for all  $\lambda$  in the definition of dominant operator such that  $\|(A - \lambda)^* x\| \leq M \|(\lambda - A)x\| \forall x \in H$ .

Also  $A$  is:

Hyponormal if  $A^*A \geq AA^*$

Quasinormal if  $[A^*A, A] = 0$

$P$ -hyponormal if for  $0 < p \leq 1, (A^*A)^p \geq (AA^*)^p$

Normal if  $[A, A^*] = 0$

Self adjoint if  $A = A^*$

Partial isometry if  $A = A A^* A$

Isometry  $A^* A = I$

Unitary if  $A^* A = A A^* = I$

Fredholm if  $\text{ran} A$  is closed and both  $\text{Ker} A$  and  $\text{Ker} A^*$  are finite dimensional.

The essential spectrum of  $A$  is denoted by  $\sigma_e(A)$ . Thus  $\sigma_e(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is not fredholm}\}$

An operator  $A$  is said to be compact if it maps the unit ball into a compact set. Equivalently  $A$  is compact if and only if  $\lim_{n \rightarrow \infty} A x_n = 0$  for each sequence  $(x_n)$  in  $H$  weakly converging to 0.

Note that the following result about properties of compact operators can easily be verified.

#### LEMMA C [2]

(i) If  $A$  is compact then so is  $A^*$

(ii) If  $A$  is compact and  $B$  is bounded then  $AB$  and  $BA$  are also compact.

We also have the following inclusion of classes of operators.

Normal  $\subset$  hyponormal  $\subset$  p-hyponormal

and Hyponormal  $\subset$  M-hyponormal  $\subset$  Dominant.

### 3. RESULTS

#### Theorem 1

Let  $A, B \in B(H)$  be quasisimilar pure dominant operators with at least one of the intertwining quasiaffinities compact. Then we have:

$$\sigma_e(A) = \sigma(A)$$

$$\sigma_e(B) = \sigma(B)$$

Proof: Since  $A$  and  $B$  are quasisimilar there exist two quasiaffinities  $X$  and  $Y$  such that

$$AX = XB \text{ and } BY = YA$$

We also have that either  $X$  or  $Y$  is compact implies  $XY$  and  $YX$  are compact operators each with dense range. It can also be verified easily that  $[A, XY] = 0$  and  $[B, YX] = 0$ . Now from theorem A above we have  $\sigma_e(A) = \sigma(A)$  and  $\sigma_e(B) = \sigma(B)$

Hence the result.

### Corollary 1

Let  $A, B \in B(H)$  be quasiinvertible operator's with either  $A$  or  $B$  compact. If  $AB$  and  $BA$  are pure dominant operators then we have  $\sigma_e(AB) = \sigma(AB)$  and  $\sigma_e(BA) = \sigma(BA)$

**Proof:** We first note that quasi invertibility is the same as quasiaffinity. Thus  $A$  and  $B$  are quasiaffinities and  $AB$  and  $BA$  are also compact quasiaffinities which are quasisimilar since we have:

$$(AB)A = A(BA) \text{ and}$$

$$(BA)B = B(AB)$$

Hence by theorem 1 above  $\sigma_e(AB) = \sigma(AB)$  and  $\sigma_e(BA) = \sigma(BA)$

### Corollary 2

Let  $A$  and  $B$  be quasisimilar pure dominant operators which satisfy Dunford's condition (C) with at least one of the implementing quasiaffinities compact. Then we have

$$\sigma_e(A) = \sigma(A) = \sigma(B) = \sigma_e(B)$$

**Proof:** The proof follows easily from both theorems  $B$  and theorem 1.

### Corollary 3

Let  $A, B \in B(H)$  be quasiinvertible operators with either  $A$  or  $B$  compact. If  $AB$  and  $BA$  are pure dominant operators satisfying Dunford's property (C) then we have

$$\sigma_e(AB) = \sigma(AB) = \sigma(BA) = \sigma_e(BA)$$

Remark; *Then its clear*  $\sigma_e(AB) = \sigma_e(BA)$

**Proof:** It follows easily from both theorem B and corollary 1 above.

### Theorem 2

Let  $A$  be a pure dominant operator and  $B$  be such that  $AX = XB$  implies  $A^*X = XB^*$  where  $X$  is a compact quasiaffinity, then  $\sigma_e(A) = \sigma(A)$ .

**Proof:** Since  $AX = XB$  implies  $A^*X = XB^*$  it can easily be verified that

$$[A, XX^*] = 0 \text{ and}$$

$$[B, X^*X] = 0$$

Where  $XX^*$  is compact with dense range. Hence by theorem A above  $\sigma_e(A) = \sigma(A)$

### Corollary 4

If  $A$  is a pure dominant operator such that

$$AX = XA^* \text{ and } A^*X = XA$$

Where  $X$  is a compact quasiaffinity then  $\sigma_e(A) = \sigma(A)$

**Proof:** In this case  $[A, XX^*] = 0$  where  $XX^*$  is compact with dense range, hence the result follows from theorem 2 above.

## REFERENCES

1. Clary, S., Equality of spectra of hyponormal operators, proc. Anaer.soc.53(1975), 88 – 90.
2. Diendonne, J., History of Functional Analysis (North-Holland Amsterdam 1981)
3. Duggal, B.P., Quasisimilar  $p$ -hyponormal operators, Equat oper. Vol 26(1996), 318 – 345)
4. Khalagai, J. M. and Nyamai B, On quasiaffinity and quasisimilarity of operators, Kenya J. Sci. Tech Vol 12 No 2 (1998), 14-17
5. Williams, J. P., Quasisimilarity and hyponormal operators J. Oper Theory 5 (1981), 127-139.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

### CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

**Prospective authors of journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

### MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

### IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

