

A Common Fixed Point Theorem for Two Random Operators using Random Mann Iteration Scheme

S. Saluja

Department of Mathematics, Government J. H. P. G. College Betul (MP) India-460001

Devkrishna Magarde

Department of Mathematics, Patel College of Science and Technology, Bhopal(MP) India-462044

Email:-devmagarde@gmail.com

Alkesh Kumar Dhakde

Department of Mathematics, IES College of Technology, Bhopal(MP) India-462044

Abstract: In this paper, we proved that if a random Mann iteration scheme is defined by two random operators is convergent under some contractive inequality the limit point is a common fixed point of each of two random operators in Banach space.

Keywords: Mann iteration, fixed point, measurable mappings, Banach space.

AMS Subject Classification: 47H10, 47H40.

1.Introduction and Preliminaries:

Kasahara [8] had shown that if an iterated sequence defined by using a continuous linear mapping is convergent under certain assumption, then the limit point is a common fixed point of each of two non-linear mappings. Ganguly [6] arrived at same conclusion by taking the same contractive condition and using the sequence of Mann iteration [9].

In this note, it is proved that if a random Mann iteration scheme is defined by two operators is convergent under some contractive inequality the limit point is a common fixed point of each of two random operators in a Banach space.

The study of random fixed point has been an active area of contemporary research in mathematics. Random iteration scheme has been elaborately discussed by Choudhury ([1], [2], [3], [4]). Looking to the immense applications of iterative algorithms in signal processing and image reconstruction, it is essential to venture upon random iteration.

We first review the following concepts, which are essential for our study.

Throughout this paper (Ω, Σ) denotes a measurable space and X stands for a separable Banach space. C is a nonempty subset of X .

A mapping $f : \Omega \rightarrow C$ is said to be measurable if $f^{-1}(B \cap C) \in \Sigma$ for every Borel subset B of X .

A mapping $F : \Omega \times C \rightarrow C$, is said to be a random operator, if $F(., x) : \Omega \rightarrow C$ is measurable for every $x \in C$.

A measurable mapping $g : \Omega \rightarrow C$ is said to be a random fixed point of the random operator $F : \Omega \times C \rightarrow C$, if $F(t, g(t)) = g(t)$ for all $t \in \Omega$.

A random operator $F : \Omega \times C \rightarrow C$ is said to be continuous if, for fixed $t \in \Omega$, $F(t, .) : C \rightarrow C$ is continuous.

Definition 1 (Random Mann Iteration scheme): Let $S, T : \Omega \times C \rightarrow C$ be two random operators on a nonempty convex subset C of a separable Banach space X . Then the sequence $\{x_n\}$ of random Mann iterates associates with S or T is defined as follows:

- (1) Let $x_0 : \Omega \rightarrow C$ be any given measurable mapping.
- (2) $x_{n+1}(t) = (1 - c_n)x_n + c_n S(t, x_n(t))$ for $n > 0, t \in \Omega$, or
- (3) $x_{n+1}(t) = (1 - c_n)x_n + c_n T(t, x_n(t))$ for $n > 0, t \in \Omega$,

where c_n satisfies:

- (4) $c_0 = 1$ for $n = 0$
- (5) $0 < c_n \leq 1$ for $n > 0$

$$(6) \quad \lim_{n \rightarrow \infty} c_n = h > 0$$

2.Main Result:

Theorem 1: Let $S, T : \Omega \times C \rightarrow C$, where C is a nonempty closed convex subset of a separable Banach space X , be two continuous random operators which satisfy the following inequality: for all $x, y \in C$ and $t \in \Omega$,

$$(7) \quad \|S(t, x) - T(t, y)\| \leq \alpha \max\{\|x(t) - y(t)\|, \|x(t) - T(t, y(t))\|, \|y(t) - S(t, x(t))\|\} \\ + \beta \max\{\|x(t) - S(t, x(t))\|, \|y(t) - T(t, y(t))\|\} \\ + \gamma \max\{\|x(t) - S(t, x(t))\| + \|x(t) - y(t)\|, \\ \|y(t) - T(t, y(t))\| + \|x(t) - y(t)\|\}$$

where $\alpha, \beta, \gamma \geq 0$ and $\alpha + \beta + \gamma < 1$.

If the sequence $\{x_n(t)\}$ of random Mann iterates associated with S or T satisfying (1)-(6) converges, then it converges to a common random fixed point of both S and T .

Proof: We may assume that the sequence $\{x_n(t)\}$ defined by (2) is pointwise convergent, that is, for all $t \in \Omega$,

$$(8) \quad x(t) = \lim_{n \rightarrow \infty} x_n(t)$$

Since X is a separable Banach space, for any continuous random operator

$A : \Omega \times C \rightarrow C$, and any measurable mapping $f : \Omega \rightarrow C$, the mapping $x(t) = A(t, f(t))$ is measurable mapping [7].

Since $x(t)$ is measurable and C is convex, it follows that $\{x_n(t)\}$ constructed in the random iteration from (2)-(6) is a sequence of measurable mappings. Hence being limit of measurable mapping sequence is also measurable. Now for $t \in \Omega$, from (2), (6) and (7) we obtain

$$\|x(t) - T(t, x(t))\| \leq \|x(t) - x_{n+1}(t)\| + \|x_{n+1}(t) - T(t, x(t))\| \\ \leq \|x(t) - x_{n+1}(t)\| + \|(1 - c_n)x_n(t) + c_n S(t, x_n(t)) - T(t, x(t))\| \\ \leq \|x(t) - x_{n+1}(t)\| + (1 - c_n)\|x_n(t) - T(t, x(t))\| \\ + c_n\|S(t, x_n(t)) - T(t, x(t))\|$$

$$(9) \quad \|x(t) - T(t, x(t))\| \leq \|x(t) - x_{n+1}(t)\| + (1 - h)\|x_n(t) - T(t, x(t))\| \\ + c_n [\alpha \max\{\|x_n(t) - x(t)\|, \|x_n(t) - T(t, x(t))\|, \\ \|x(t) - S(t, x_n(t))\|\} + \beta \max\{\|x_n(t) - S(t, x_n(t))\|, \\ \|x(t) - T(t, x(t))\|\} + \gamma \max\{\|x_n(t) - S(t, x_n(t))\| + \\ \|x_n(t) - x(t)\|, \|x(t) - T(t, x(t))\| + \|x_n(t) - x(t)\|\}]$$

Now $c_n(S(t, x_n(t)) - x_n(t)) = c_n S(t, x_n(t)) - c_n x_n(t) = x_{n+1}(t) - x_n(t)$

Implies that $\|S(t, x_n(t)) - x_n(t)\| \leq \frac{1}{c_n} \|x_{n+1}(t) - x_n(t)\|$

This shows that for $t \in \Omega$, $S(t, x_n(t)) - x_n(t) \rightarrow 0$ as $n \rightarrow \infty$ and so

$S(t, x_n(t)) - x(t) \rightarrow 0$ as $n \rightarrow \infty$ as S is continuous random operator and x is a measurable mapping. Consequently from (9) on taking limit as $n \rightarrow \infty$ we obtain

$$\|x(t) - T(t, x(t))\| \leq 0 + (1 - h)\|x(t) - T(t, x(t))\| + c_n [\alpha \max\{0, \|x(t) - T(t, x(t))\|, 0\} \\ + \beta \max\{0, \|x(t) - T(t, x(t))\|\} + \gamma \max\{0, \|x(t) - T(t, x(t))\|\}]$$

$$\leq (1 - h + h\alpha + h\beta + h\gamma)\|x(t) - T(t, x(t))\|$$

implies that $T(t, x(t)) = x(t)$ for all $t \in \Omega$ (since $\alpha + \beta + \gamma < 1$) as T is continuous random operator and x is measurable.

Therefore,

$$\begin{aligned} \|S(t, x(t)) - x(t)\| &= \|S(t, x(t)) - T(t, x(t))\| \\ &\leq \alpha \max\{\|x(t) - x(t)\|, \|x(t) - T(t, x(t))\|, \|x(t) - S(t, x(t))\|\} \\ &\quad + \beta \max\{\|x(t) - S(t, x(t))\|, \|x(t) - T(t, x(t))\|\} \\ &\quad + \gamma \max\{\|x(t) - S(t, x(t))\| + \|x(t) - x(t)\|, \\ &\quad\quad\quad \|x(t) - T(t, x(t))\| + \|x(t) - x(t)\|\} \end{aligned}$$

$$\begin{aligned} \|S(t, x(t)) - x(t)\| &\leq \alpha \max\{\|x(t) - x(t)\|, \|x(t) - x(t)\|, \|x(t) - S(t, x(t))\|\} \\ &\quad + \beta \max\{\|x(t) - S(t, x(t))\|, \|x(t) - x(t)\|\} \\ &\quad + \gamma \max\{\|x(t) - S(t, x(t))\| + \|x(t) - x(t)\|, \\ &\quad\quad\quad \|x(t) - x(t)\| + \|x(t) - x(t)\|\} \\ &\leq \alpha \max\{0, 0, \|x(t) - S(t, x(t))\|\} + \beta \max\{\|x(t) - S(t, x(t))\|, 0\} \\ &\quad + \gamma \max\{\|x(t) - S(t, x(t))\|, 0\} \\ &\leq (\alpha + \beta + \gamma)\|x(t) - S(t, x(t))\| \end{aligned}$$

Since $\alpha + \beta + \gamma < 1$ implies that $S(t, x(t)) = x(t)$.

Uniqueness:-Let $v(t)$, $v(t) \neq x(t)$ is another common fixed point of S and T , then, using (7), we have

$$\begin{aligned} \|x(t) - v(t)\| &\leq \alpha \max\{\|x(t) - v(t)\|, \|x(t) - T(t, v(t))\|, \|v(t) - S(t, x(t))\|\} \\ &\quad + \beta \max\{\|x(t) - S(t, x(t))\|, \|v(t) - T(t, v(t))\|\} \\ &\quad + \gamma \max\{\|x(t) - S(t, x(t))\| + \|x(t) - v(t)\|, \\ &\quad\quad\quad \|v(t) - T(t, v(t))\| + \|x(t) - v(t)\|\} \\ &\leq \alpha \max\{\|x(t) - v(t)\|, \|x(t) - v(t)\|, \|v(t) - x(t)\|\} \\ &\quad + \beta \max\{\|x(t) - x(t)\|, \|v(t) - v(t)\|\} \\ &\quad + \gamma \max\{\|x(t) - x(t)\| + \|x(t) - v(t)\|, \\ &\quad\quad\quad \|v(t) - v(t)\| + \|x(t) - v(t)\|\} \\ &\leq (\alpha + \gamma)\|x(t) - v(t)\| \end{aligned}$$

$$\Rightarrow x(t) = v(t) \text{ as } \alpha + \gamma < 1$$

This complete the proof.

References:

[1] Choudhury B. S., Convergence of a random iteration scheme to a random fixed point, *J. Appl. Math. Stoc. Anal.*, 8(1995), 139-142.
 [2] Choudhury B. S., Random Mann iteration scheme, *Appl. Math. Lett.*, 16 (2003), 93-96.
 [3] Choudhury B. S., Ray M., Convergence of an iteration leading to a solution of a random operator equation, *J. Appl. Math. Stoc. Anal.*, 12 (1999), 161-168.
 [4] Choudhury B. S., Upadhyay A., An iteration leading to a solution and fixed point of operators, *Soochow J. Math.*, 25 (1999), 394-400.
 [5] Ciric Lj., Quasi-contractions in Banach space, *Publ. Inst. Math.*, 21(35) (1977), 41-48.

- [6] Ganguly A. K., On common fixed point of two mappings, *Mathematics Seminar Notes.*, 8 (1980), 343-345.
- [7] Himmelberg C.J., Measurable relations, *Fund.Math.*, LXXXVII (1975), 53-71.
- [8] Kasahara S., Fixed point iterations using linear mappings, *Mathematics Seminar Notes.*, 6 (1978), 87-90.
- [9] Rhoades B. E., Extensions of some fixed point theorems of Ciric, Maiti and Pal, *Mathematics Seminar Notes.*, 6 (1978), 41-46.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

